
a
Catabolic and osmotic modules rare or undetected in FSM
% of genomes encoding
RGM (n = 97)
FSM (n = 95)
Leprosy clade, three tissue-derived genomes
82%
1%
0%
47%
1%
0%
19%
0%
0%
59%
0%
0%
75%
8%
0%
ureC
narG
catA
ectC
kdpB
b
Habitat of clade-dominant metagenomes
Clade-dominant grid cells (%)
15%
14%
54%
21%
28%
12%
21%
11%
12%
12%
56%
SGM
65 cells
RGM
273 cells
FSM
202 cells
freshwater
soil/sediment
wastewater
drinking-water
marine/saline
host-assoc.
aquaculture
other
c
Detection in public metagenome runs
Lake/reservoir
River/stream
Estuary/brackish
Marine
Ground/drinking water
Sediment
Soil/peat/wetland
Wastewater
Host-associated
FSM
RGM (+ sister RGM)
0.22
0.03
0.46
Runs with the clade detected (%)
9 of 13 habitat classes shown (Table S11i)
d
Aquatic against other biomes
Genomes with biome metadata (%)
26
16
51
34
19
75
SGM
n=42
RGM
n=85
FSM
n=94
aquatic
other biomes (terrestrial, anthropogenic, host)