

# DYNAMIC ČECH COMPLEXES, SHEAVES AND PERSISTENT COHOMOLOGY

JIAQI TANG

June 11, 2026

## ABSTRACT

We propose the concepts of dynamic Čech complexes, dynamic Čech sheaves and persistent dynamic cohomology. We also prove the differentiability of dynamic complex flows, as well as the stability and convergence of persistent dynamic cohomology. To address the indecomposability of persistent dynamic cohomology modules, we introduce the dynamic topological feature measure.

**Keywords:** Algebraic Topology; Persistent Homology and Applications; Topological Data Analysis

## CONTENTS

|     |                                                                  |    |
|-----|------------------------------------------------------------------|----|
| 1   | INTRODUCTION                                                     | 1  |
| 2   | PRELIMINARIES                                                    | 3  |
| 2.1 | COMPLEXES AND PERSISTENT HOMOLOGY                                | 3  |
| 2.2 | ČECH COHOMOLOGY AND SHEAVES                                      | 9  |
| 3   | DYNAMIC ČECH COMPLEXES AND DYNAMIC ČECH SHEAVES                  | 13 |
| 3.1 | THEORY OF DYNAMIC ČECH COMPLEXES                                 | 13 |
| 3.2 | CONSTRUCTION OF THE DYNAMIC ČECH SHEAF                           | 23 |
| 4   | PERSISTENT DYNAMIC COHOMOLOGY THEORY                             | 30 |
| 4.1 | FUNDAMENTAL THEORY OF PERSISTENT DYNAMIC COHOMOLOGY              | 30 |
| 4.2 | CONVERGENCE, STABILITY, AND DYNAMIC TOPOLOGICAL FEATURE MEASURES | 33 |
| 4.3 | STATISTICS AND ANALYSIS OF DYNAMIC TOPOLOGICAL FEATURES          | 36 |
| 5   | CONCLUSIONS AND OUTLOOK                                          | 39 |
| A   | PROOF OF STABILITY THEOREM 4.9 AND THEOREM 4.13                  | 40 |
| B   | PROOF OF CONVERGENCE THEOREM 4.10 AND ITS COROLLARIES            | 43 |
| B.1 | CONVERGENCE IN THE DYNAMIC INTERPOLATION DISTANCE                | 43 |
| B.2 | OTHER CONVERGENCE RESULTS                                        | 43 |

# 1 INTRODUCTION

In modern data science, the emergence of high-dimensional data poses severe challenges to traditional analytical methods. Although data generated from fields such as sensor detection, gene sequencing, and image feature extraction is presented in high-dimensional form, its intrinsic structure is often constrained by underlying geometric rules—namely, the generated high-dimensional observational data is typically embedded in a low-dimensional manifold structure[CG09]. Topological Data Analysis (TDA) has emerged as a result, and its core problem can be formulated as follows: given a finite point set  $X_n = \{x_1, \dots, x_n\}$  in a high-dimensional observational space  $\mathbb{R}^D$ , it is essentially a random sample of an unknown  $d$ -dimensional compact Riemannian manifold  $(M, g)$  under a certain probability distribution  $\mu$ , where  $d \ll D$ , i.e.,  $X_n \stackrel{\text{i.i.d.}}{\sim} \mu$ .

The core task of TDA is to construct a mapping  $\mathcal{C}$  from the power set  $2^{\mathbb{R}^D}$  to a certain discrete label set  $\mathcal{Y}$ , such that for a sample  $X'$  from an unknown manifold  $M'$  and the true label  $y$  of  $M'$ , we have

$$\mathbb{P}(\mathcal{C}(X') \neq y) < \varepsilon, \quad \varepsilon > 0.$$

Moreover, this mapping  $\mathcal{C}$  can preserve the topological invariants of  $M$ , thus being robust to noise and redundancy.

To recover the topology of  $M$  from discrete samples  $X'$ , Persistent Homology, systematically proposed by Edelsbrunner et al.[ELZ02], has become a core tool. Its approach involves constructing a Filtered Simplicial Complex and studying the evolution of its homology groups. This method has been successfully applied to numerous fields, ranging from molecular structure analysis to cosmology research[Was16].

Traditional TDA mostly relies on the Vietoris-Rips complex, which is widely used due to its high computational efficiency[Ott+17]. However, the Čech complex provides a more accurate topological approximation through its cover-based geometric construction. Carlsson[Car14] pointed out that the direct connection between the Čech complex and the Nerve Lemma gives it significant theoretical advantages. In recent years, with the improvement of computational capabilities, the Čech method has regained attention, especially in applications requiring high-precision topological features.

As a powerful tool for describing local-global relationships, Sheaf Theory has been introduced into TDA in recent years. Robinson systematically elaborated on the applications of Sheaf Theory in signal processing in Topological Signal Processing[Rob14]. More relevantly, the work of de Silva et al.[SG07] explored the sheaf-theoretic structure of multi-parameter persistent homology, laying the foundation for the in-depth application of Sheaf Theory in TDA.

Topological analysis of dynamic data is a current research hotspot. Existing methods for handling dynamic data are mostly based on discretization and static approximation. Let  $I = [0, T]$  be the time parameter interval, and the dataset be  $\{X_n(t)\}_{t \in I}$ . The mainstream approach is to construct discrete time slices  $\{t_0, t_1, \dots, t_m\}$ , establish a static complex for each  $t_k$ , and then analyze its persistent homology[PH13]. Although multi-parameter persistent homology provides a more general theoretical framework, it suffers from high computational complexity and faces the fundamental challenge of indecomposability, while failing to fully utilize the geometric properties of the Čech complex.

Despite progress in research, there remain the following essential deficiencies at the mathematical level:

1. **Lack of a continuous dynamic theory:** Existing methods fail to define an intrinsic geometric structure of complexes that continuously depends on parameters. Their ana-

lytical objects are discrete sequences  $\{\mathcal{K}_{t_k}\}_{k=1}^m$ , rather than a continuous “simplicial flow”  $\{\mathcal{C}(t)\}_{t \in I}$ .

2. **Separation of Čech theory and Sheaf Theory:** The strong geometric expressiveness of the Čech complex and the excellent ability of Sheaf Theory to describe local-global relationships have not been unified within a dynamic framework. Specifically, there is a lack of a sheaf-theoretic structure to unify the intuitively “dynamic Čech complex”, making it impossible to establish a local-global correspondence principle for dynamic topological features.
3. **Challenge of indecomposability for multi-parameter persistence modules:** Bi-parameter persistence modules are indecomposable in general cases[BS25], and traditional barcode decomposition methods fail, posing a fundamental theoretical obstacle to dynamic topological analysis.

To address the above issues, this paper aims to construct a unified framework for dynamic Čech cohomology and Sheaf Theory on manifolds. The core lies in the mathematical construction at the following three levels:

1. **Dynamic Čech Complex:** For the first time, a rigorous definition of the parameterized Čech complex is given. Considering a continuous parameter space  $T$ , a family of Čech complexes  $\mathcal{C}(t)$  varying with  $t \in T$  is defined, thereby forming a complex-valued functor  $\mathcal{C}: T \rightarrow \text{Simp}$ , and its continuity and differentiability theories are established.
2. **Dynamic Čech Sheaf:** A dynamic Čech sheaf  $\mathcal{D}_{\Phi, \epsilon}$  is constructed on the product space  $M \times T$ . Through dynamic star neighborhoods and local consistency conditions, a local-global correspondence for the continuous evolution of topological features is established.
3. **Persistent Dynamic Cohomology:** The Persistent Dynamic Cohomology module  $\mathbb{PDH}^p(\Phi)$  is defined. By introducing the dynamic topological feature measure  $\mu_{\Phi}^p$  to characterize the distribution of topological features in the scale-time parameter space, the indecomposability problem of multi-parameter persistence modules is circumvented, and a stability theorem under the dynamic interpolation distance is proven.

Furthermore, for the discrete data collected, we first reconstruct it into continuous data through continuous interpolation or continuous models, and then apply the dynamic TDA method to be developed in this paper to obtain stronger conclusions compared with traditional TDA (e.g., stability, differentiability and predictability).

This paper is organized as follows:

In section 2, we review the preliminary knowledge related to simplicial complexes, persistent homology, Čech cohomology, sheaf theory, some fundamental definitions and propositions used throughout this paper.

In section 3, we construct dynamic Čech complexes and dynamic Čech sheaves, and analyze their basic properties.

In section 4, we construct the framework of persistent dynamic cohomology and resolve the indecomposability of two-parameter persistence modules via dynamic topological feature measures. We derive the stability theorem 4.9 and convergence theorem 4.10 for the dynamic case by referring to traditional persistent cohomology, and present relevant statistical tools.

Appendix A presents the proof of stability theorem 4.9 and theorem 4.13.

Appendix B presents the proof of convergence theorem 4.10 and its corollaries.

## 2 PRELIMINARIES

In this section, we shall give some notions and fundamental results about simplicial complexes, persistent homology, Čech cohomology and sheaf theory, which serve as the theoretical basis for the subsequent research.

We refer the readers [EH21; GQ20; Oud15; Jr08; Han25; KS14; BCM25; Fri25; Hat02] for more details.

### 2.1 COMPLEXES AND PERSISTENT HOMOLOGY

*Definition 2.1* (Abstract Simplicial Complex). An **abstract simplicial complex**  $K$  consists of a finite set  $V$  (called the vertex set) together with a family  $\Sigma$  of subsets of  $V$  (called the simplex set) satisfying:

1. For any  $v \in V$ , we have  $\{v\} \in \Sigma$ .
2. If  $\sigma \in \Sigma$  and  $\tau \subseteq \sigma$ , then  $\tau \in \Sigma$ .

Sometimes, we write  $\sigma \in K$  means  $\sigma \in \Sigma$ .

*Definition 2.2* ( $p$ -Simplex). Let  $K$  be a simplicial complex. The dimension of a simplex  $\sigma \in \Sigma$  is defined as  $\dim \sigma = |\sigma| - 1$ . A  **$p$ -simplex** of  $K$  is a simplex in  $K$  of dimension  $p$ , i.e., a  $\sigma \in \Sigma$  such that  $|\sigma| = p + 1$ . In particular, 0-simplices are called vertices, 1-simplices are called edges, 2-simplices are called triangular faces, and 3-simplices are called tetrahedra.

After introducing the concept of simplicial complexes, we proceed to present two commonly used complexes: the VR complex and the Čech complex.

*Definition 2.3* (Vietoris–Rips Complex). Given a metric space  $(X, d)$  and a radius  $\varepsilon > 0$ , the **Vietoris–Rips complex** of a point set  $X_n \subset X$  is defined as:

$$\mathcal{VR}_\varepsilon(X_n) := \{\sigma \subseteq X_n \mid \forall x, y \in \sigma, d(x, y) \leq \varepsilon\}.$$

*Definition 2.4* (Čech Complex). Given a metric space  $(X, d)$  and a radius  $\varepsilon > 0$ , the **Čech complex** of a point set  $X_n \subset X$  is defined as:

$$\mathcal{C}_\varepsilon(X_n) := \left\{ \sigma \subseteq X_n \mid \bigcap_{x \in \sigma} B(x, \varepsilon) \neq \emptyset \right\}.$$

where  $B(x, \varepsilon)$  denotes the open ball centered at  $x$  with radius  $\varepsilon$ .

**Proposition 2.5** (Inclusion Relations Between Complexes). *For a point set  $X_n$  in the same metric space and any  $\varepsilon > 0$ , we have:*

$$\mathcal{C}_\varepsilon(X_n) \subseteq \mathcal{VR}_{2\varepsilon}(X_n) \subseteq \mathcal{C}_{2\varepsilon}(X_n).$$

*Proof.* Let  $\sigma \in \mathcal{C}_\varepsilon(X_n)$ , we can take any point  $z \in \bigcap_{x \in \sigma} B(x, \varepsilon)$ . Then for all  $x, y \in \sigma$ , we have

$$d(x, z) < \varepsilon, d(y, z) < \varepsilon \Rightarrow d(x, y) \leq d(x, z) + d(z, y) < 2\varepsilon \Rightarrow \sigma \in \mathcal{VR}_{2\varepsilon}(X_n).$$

Otherside,  $\sigma \in \mathcal{VR}_{2\varepsilon}(X_n) \iff \forall x, y \in \sigma \text{ s.t. } d(x, y) < 2\varepsilon$ . For any vertex  $x_0 \in \sigma$ , for every  $x \in \sigma$ ,  $d(x, x_0) < 2\varepsilon \Rightarrow x_0 \in B(x, 2\varepsilon)$ . Thus  $x_0 \in \bigcap_{x \in \sigma} B(x, 2\varepsilon)$ . Hence  $\sigma \in \mathcal{C}_{2\varepsilon}(X_n)$ .  $\square$

Besides, some fundamental definitions regarding chain complex, subcomplex and filtered complex are necessary for subsequent discussion.

**Definition 2.6** (Chain Complex). A **chain complex**  $(C_\bullet, \partial_\bullet)$  is a sequence of abelian groups (or modules)  $C_p$  ( $p \in \mathbb{Z}$ ) together with group homomorphisms (called boundary operators)  $\partial_p: C_p \rightarrow C_{p-1}$ :

$$\cdots \longrightarrow C_{p+1} \xrightarrow{\partial_{p+1}} C_p \xrightarrow{\partial_p} C_{p-1} \longrightarrow \cdots$$

such that  $\partial_p \circ \partial_{p+1} = 0$  for all  $p$ .

**Definition 2.7** (Homology Group). The  **$p$ -th homology group** of a chain complex  $(C_\bullet, \partial_\bullet)$  is defined as:

$$H_p(C_\bullet) := \ker \partial_p / \partial_{p+1} C_{p+1}$$

An element  $c \in \ker \partial_p$  is called a  $p$ -cycle, and an element  $c \in \text{im } \partial_{p+1}$  is called a  $p$ -boundary.

**Definition 2.8** (Simplicial Chain Complex). Given a simplicial complex  $K$ , its **simplicial chain complex**  $C_\bullet(K)$  is defined as follows:

- $C_p(K)$  is the free abelian group generated by all  $p$ -simplices in  $K$ .
- The boundary homomorphism  $\partial_p: C_p(K) \rightarrow C_{p-1}(K)$  is defined on the generator  $[v_0, \dots, v_p]$  by:

$$\partial_p([v_0, \dots, v_p]) = \sum_{i=0}^p (-1)^i [v_0, \dots, \widehat{v_i}, \dots, v_p].$$

**Proposition 2.9.** *The simplicial chain complex  $(C_\bullet(K), \partial_\bullet)$  satisfies  $\partial_p \circ \partial_{p+1} = 0$ .*

*Proof.* Let  $[v_0, \dots, v_{p+1}]$  be a  $(p+1)$ -simplex. By definition,

$$\begin{aligned} \partial_{p+1}[v_0, \dots, v_{p+1}] &= \sum_{i=0}^{p+1} (-1)^i [v_0, \dots, \widehat{v_i}, \dots, v_{p+1}], \\ \partial_p \partial_{p+1}[v_0, \dots, v_{p+1}] &= \sum_{i=0}^{p+1} (-1)^i \partial_p [v_0, \dots, \widehat{v_i}, \dots, v_{p+1}] \\ &= \sum_{i=0}^{p+1} \left( \sum_{0 \leq j < i \leq p+1} (-1)^{i+j} [v_0, \dots, \widehat{v_j}, \dots, \widehat{v_i}, \dots, v_{p+1}] \right. \\ &\quad \left. + \sum_{0 \leq i < j \leq p+1} (-1)^{i+j+1} [v_0, \dots, \widehat{v_i}, \dots, \widehat{v_j}, \dots, v_{p+1}] \right) = 0 \end{aligned}$$

□

**Definition 2.10** (Subcomplex). Let  $K = (V_K, \Sigma_K)$  be an abstract simplicial complex. A pair  $L = (V_L, \Sigma_L)$  is a **subcomplex** of  $K$ , written  $L \subseteq K$ , if  $L$  is an abstract simplicial complex such that  $V_L \subseteq V_K$  and  $\Sigma_L \subseteq \Sigma_K$ .

**Definition 2.11** (Filtered Complex). Let  $K$  be a simplicial complex. A filtration of  $K$  is a family of subcomplexes  $\{K_t\}_{t \in \mathbb{R}}$  such that:

1. If  $s \leq t$ , then  $K_s \subseteq K_t$ .
2. There exist  $t_{\min}, t_{\max} \in \mathbb{R}$  such that  $K_t = \emptyset$  for  $t < t_{\min}$  and  $K_t = K$  for  $t \geq t_{\max}$ .

The pair  $(K, \{K_t\}_{t \in \mathbb{R}})$  is called a **filtered complex**.

We then introduce persistent homology, a crucial concept in TDA.

*Definition 2.12* (Persistent Homology). Let  $\{K_t\}_{t \in \mathbb{R}}$  be a filtered complex. For each  $t \in \mathbb{R}$ , consider the simplicial chain complex  $C_\bullet(K_t)$  and its homology groups  $H_p(K_t)$ .

For  $s \leq t$ , the inclusion map  $i_{s,t} : K_s \hookrightarrow K_t$  induces a chain map  $(i_{s,t})_\# : C_\bullet(K_s) \rightarrow C_\bullet(K_t)$ , for each  $p \leq 0$ ,

$$\begin{aligned} (i_{s,t})_{\#,p} : C_p(K_s) &\longrightarrow C_p(K_t) \\ \sum_i a_i \sigma_i &\longmapsto \sum_i a_i i_{s,t}(\sigma_i) \end{aligned}$$

which in turn induces a homomorphism of homology groups:

$$\begin{aligned} \varphi_s^{t,(p)} : H_p(K_s) &\longrightarrow H_p(K_t) \\ [z]_s^{(p)} &\longmapsto [(i_{s,t})_{\#,p}(z)]_t^{(p)} \end{aligned}$$

The  **$p$ -th persistent homology** refers to the family of homology groups together with the transition homomorphisms between them:

$$\text{PH}_p(\{K_t\}_{t \in \mathbb{R}}) = \left( H_p(K_t), \varphi_s^{t,(p)} \right)_{s \leq t \in \mathbb{R}}.$$

$$\begin{array}{ccccccc} & & \vdots & & \vdots & & \vdots \\ & & \uparrow & & \uparrow & & \uparrow \\ \cdots & \longrightarrow & C_{p+1}(K_t) & \xrightarrow{\partial_{p+1}} & C_p(K_t) & \xrightarrow{\partial_p} & C_{p-1}(K_t) \longrightarrow \cdots \\ & & (i_{s,t})_{\#,p+1} \uparrow & & (i_{s,t})_{\#,p} \uparrow & & (i_{s,t})_{\#,p-1} \uparrow \\ \cdots & \longrightarrow & C_{p+1}(K_s) & \xrightarrow{\partial_{p+1}} & C_p(K_s) & \xrightarrow{\partial_p} & C_{p-1}(K_s) \longrightarrow \cdots \\ & & \uparrow & & \uparrow & & \uparrow \\ & & \vdots & & \vdots & & \vdots \end{array}$$

*Remark 2.13.* The homomorphism  $\varphi_s^{t,(p)}$  is well-defined for the following reasons.

First, we have  $\partial_p(i_{s,t})_{\#,p+1} = (i_{s,t})_{\#,p} \partial_p$ , thus it maps cycle to cycle, i.e., if  $\partial_p(z) = 0$ , then  $\partial_p((i_{s,t})_{\#,p+1}(z)) = (i_{s,t})_{\#,p}(\partial_p(z)) = 0$ .

And if  $[z]_s^{(p)} \sim [z']_s^{(p)}$ , i.e.,  $z - z' = \partial(c')$  then  $(i_{s,t})_{\#,p}(z) - (i_{s,t})_{\#,p}(z') = (i_{s,t})_{\#,p}(z - z') = \partial_{p+1}(i_{s,t})_{\#,p+1}(c')$  then  $[(i_{s,t})_{\#,p}(z)]_t^{(p)} \sim [(i_{s,t})_{\#,p}(z')]_t^{(p)}$ .

We present the definition of persistence modules as follows.

*Definition 2.14* (Persistence Module). Regard  $(\mathbb{R}, \leq)$  as a poset category. A **persistence module**  $M$  is a functor:

$$M : (\mathbb{R}, \leq) \rightarrow \mathbf{Vect}_K$$

where  $\mathbf{Vect}_K$  is the category of vector spaces over a field  $K$ . Concretely:

1. For each  $t \in \mathbb{R}$ , a vector space  $M_t$  is assigned.
2. For each pair  $s \leq t$ , a linear map  $\varphi_s^t : M_s \rightarrow M_t$  is assigned.

3. It satisfies  $\varphi_t^t = \text{id}_{M_t}$ , and for  $r \leq s \leq t$  we have  $\varphi_r^t = \varphi_s^t \circ \varphi_r^s$ .

**Proposition 2.15.** *Persistent homology  $\text{PH}_p(\{K_t\})$  naturally forms a persistence module.*

*Proof.* Define a functor  $M : (\mathbb{R}, \leq) \rightarrow \mathbf{Vect}_K$  as follows:

1. For any  $t \in \mathbb{R}$ , set  $M(t) = H_p(K_t)$ .
2. For any  $s \leq t$ , define the transition map  $M(s \leq t) = \varphi_s^{t,(p)} : H_p(K_s) \rightarrow H_p(K_t)$ .

For each simplicial complex  $K_t$ , the chain groups  $C_p(K_t)$  are finite-dimensional  $\mathbb{F}$ -vector space, then  $\ker \partial_p$  and  $\text{im } \partial_{p+1}$  are all the subspace of  $C_p(K_t)$ . Then by definition, the quotient space  $H_p(K_t) = \ker \partial_p / \text{im } \partial_{p+1}$  is again a vector space over  $\mathbb{F}$ .

For any  $r \leq s \leq t$ , consider the composition of inclusions:

$$K_r \xhookrightarrow{i_{r,s}} K_s \xhookrightarrow{i_{s,t}} K_t$$

By the definition of a filtration of simplicial complexes, this composition equals the inclusion  $i_{r,t} : K_r \hookrightarrow K_t$ .

At the chain complex level, these inclusions induce chain maps:

$$C_\bullet(K_r) \xrightarrow{(i_{r,s})_\#} C_\bullet(K_s) \xrightarrow{(i_{s,t})_\#} C_\bullet(K_t)$$

and  $(i_{s,t})_\# \circ (i_{r,s})_\# = (i_{r,t})_\#$ .

At the homology level, chain maps induce homomorphisms:

$$\begin{aligned} H_p(K_r) &\xrightarrow{\varphi_r^{s,(p)}} H_p(K_s) \xrightarrow{\varphi_s^{t,(p)}} H_p(K_t) \\ [z]_r^p &\longmapsto [(i_{r,s})_{\#,p}(z)]_r^p \longmapsto [(i_{s,t})_{\#,p}(i_{r,s})_{\#,p}(z)]_r^p = [(i_{r,t})_{\#,p}(z)]_r^p \end{aligned}$$

We have  $\varphi_s^{t,(p)} \circ \varphi_r^{s,(p)} = \varphi_r^{t,(p)}$ , i.e.,  $M(s \leq t) \circ M(r \leq s) = M(r \leq t)$ . □

**Definition 2.16** (Interval Module). Let  $I \subseteq \mathbb{R}$  be an interval and  $K$  a field. The **interval module**  $\mathbb{I}_I$  is defined by:

$$\mathbb{I}_I(t) := \begin{cases} K, & t \in I, \\ 0, & t \notin I. \end{cases}$$

The transition map  $\varphi_s^t : \mathbb{I}_I(s) \rightarrow \mathbb{I}_I(t)$  is the identity map if  $s, t \in I$ , and the zero map otherwise.

**Definition 2.17** (Multiset of Intervals). A **multiset** is a collection of elements where multiple instances of the same elements are allowed.

Single-parameter persistence modules enjoy a very well decomposition property, which is summarized in Theorem 2.18

**Theorem 2.18** (Structure Theorem for Persistence Modules). *Let  $K$  be a field and  $M$  a persistence module over  $\mathbb{R}$ . Then there exists a unique multiset  $\text{Barc}(M) = \{[b_i, d_i]\}_i$  (called barcode) such that:*

$$M \cong \bigoplus_{I \in \text{Barc}(M)} \mathbb{I}_I = \bigoplus_i \mathbb{I}_{[b_i, d_i]}.$$

**Remark 2.19.** Thanks to the guarantee of Theorem 2.18, the theory of TDA has achieved tremendous progress.

*Definition 2.20* (Persistence Diagram). The **persistence diagram**  $\text{Dgm}(M)$  of a persistence module  $M$  is the graphical representation of its barcode  $\text{Barc}(M)$ , where each interval  $I = [b, d)$  is represented as a point  $(b, d) \in \mathbb{R}^2$ .

*Remark 2.21.*  $\text{Dgm}(M)$  is a multiset.

*Definition 2.22* (Bottleneck Distance). Let  $\text{Dgm}_1, \text{Dgm}_2$  be two persistence diagrams. Their **bottleneck distance** is defined as:

$$d_B(\text{Dgm}_1, \text{Dgm}_2) := \inf_{\gamma} \sup_{x \in \text{Dgm}_1} \|x - \gamma(x)\|_{\infty}.$$

where matching  $\gamma$  between  $\text{Dgm}_1$  and  $\text{Dgm}_2$  is a bijection that pairs each point in  $\text{Dgm}_1$  with a point in  $\text{Dgm}_2$  or with a point on the diagonal  $\Delta = \{(t, t) \in \mathbb{R}^2\}$ . And for any point  $p_i = (b_i, d_i)$ ,  $i = 1, 2$ , the  $L^\infty$ -distance between them is defined as  $\|p_1 - p_2\| = \max\{|b_1 - b_2|, |d_1 - d_2|\}$ .

In conventional stability theorem, the right hand side is bounded by the Gromov-Hausdorff distance, while we adopt the Hausdorff distance for the bound in Theorem 2.23 below. Furthermore, we provide a concise proof together.

**Theorem 2.23** (Stability Theorem). *Let  $X, Y$  be finite point sets in a compact metric space, and let  $\{K_t(X)\}, \{K_t(Y)\}$  be the corresponding filtered complexes (such as Čech or Vietoris–Rips filtrations). Then*

$$d_B(\text{Dgm}(\text{PH}_p(K_t(X))), \text{Dgm}(\text{PH}_p(K_t(Y)))) \leq 2 \cdot d_H(X, Y).$$

where  $d_H$  denotes the Hausdorff distance.

*Proof.* [CSO13] gives us

$$\begin{cases} d_B(\text{Dgm}(\mathcal{Rips}(X)), \text{Dgm}(\mathcal{Rips}(Y))) \leq 2d_{GH}(X, Y) \\ d_B(\text{Dgm}(\mathcal{Cech}(X)), \text{Dgm}(\mathcal{Cech}(Y))) \leq 2d_{GH}(X, Y) \end{cases}$$

for finite metric space  $(X, d_X)$  and  $(Y, d_Y)$ , where

$$d_{GH}(X, Y) = \frac{1}{2} \inf \left\{ \text{dis}(C) : C \text{ is a correspondence } X \rightrightarrows Y \right\}$$

is the Gromov-Hausdorff distance (see [DYS] Theorem 7.3.25), and

$$\text{dis}(C) = \sup \{|d_X(x, x') - d_Y(y, y')| : (x, y), (x', y') \in C\}.$$

Now we prove if  $X, Y$  belong to the same metric space  $(Z, d)$ , then  $d_{GH}(X, Y) \leq d_H(X, Y)$ . Let  $(Z, d)$  be a metric space, and let  $X, Y \subseteq Z$  be non-empty compact subsets with the include metric  $d$ .

Define  $C_0 \subseteq X \times Y$  by  $C_0 = \{(x, y) \in X \times Y : d(x, y) \leq d_H(X, Y)\}$ . By definition of Hausdorff distance, for every  $x \in X$ , there exists  $y \in Y$  such that  $d(x, y) \leq d_H(X, Y)$ . Thus  $C_0$  is a valid correspondence between  $X$  and  $Y$ .

For any pairs  $(x, y), (x', y') \in C_0$ , by the triangle inequality,

$$|d(x, x') - d(y, y')| \leq d(x, y) + d(x', y') \leq 2d_H(X, Y)$$

Taking the supremum over all such pairs gives  $\text{dis}(C_0) \leq 2d_H(X, Y)$ . Then

$$d_{GH}(X, Y) \leq \frac{1}{2} \text{dis}(C_0) \leq d_H(X, Y).$$

□



After introducing concepts related to homology group, we will correspondingly present preliminary knowledge about cohomology group and persistent cohomology.

*Definition 2.24* (Cohomology Group). Let  $K$  be a simplicial complex and  $G$  an abelian group. The cochain complex  $C^\bullet(K; G)$  of  $K$  is defined by:

1.  $C^p(K; G) := \text{Hom}(C_p(K), G)$ , i.e., the set of group homomorphisms from the  $p$ -chain group to  $G$ .
2. For any  $(p+1)$ -simplex, the coboundary operator  $\delta^p : C^p(K; G) \rightarrow C^{p+1}(K; G)$  is defined by  $(\delta^p f)(\sigma) = f(\partial_{p+1}\sigma)$ .

$$\begin{array}{ccccccc} \dots & \longrightarrow & C^{p-1}(K; G) & \xrightarrow{\delta^{p-1}} & C^p(K; G) & \xrightarrow{\delta^p} & C^{p+1}(K; G) \longrightarrow \dots \\ \dots & \longmapsto & f \circ \partial_p^{-1} & \longmapsto & f & \longmapsto & f \circ \partial_{p+1} \longmapsto \dots \end{array}$$

The  $p$ -th cohomology group of  $K$  is defined as:

$$H^p(K; G) := \ker \delta^p / \delta^{p-1} C^{p-1}(K; G)$$

where elements of  $\ker \delta^p$  are called  $p$ -cocycles, and elements of  $\delta^{p-1} C^{p-1}(K; G)$  are called  $p$ -coboundaries.

*Definition 2.25* (Persistent Cohomology). Let  $\{K_t\}_{t \in \mathbb{R}}$  be a filtered complex. For each  $t \in \mathbb{R}$ , consider the cochain complex  $C^\bullet(K_t; G)$  and its cohomology groups  $H^p(K_t; G)$ .

For  $s \leq t$ , the inclusion map  $i_{s,t} : K_s \hookrightarrow K_t$  induces a chain map  $(i_{s,t})_\# : C_\bullet(K_s) \rightarrow C_\bullet(K_t)$ , for each  $p$ ,  $(i_{s,t})_{\#,p} : C_p(K_s) \rightarrow C_p(K_t)$ , which by duality induces a cochain map  $(i_{s,t})^* : C^\bullet(K_t; G) \rightarrow C^\bullet(K_s; G)$ , for each  $p$ ,

$$\begin{array}{ccc} (i_{s,t})^{*,p} : C^p(K_t; G) & \longrightarrow & C^p(K_s; G) \\ f & \longmapsto & f \circ (i_{s,t})_{\#,p} \end{array}$$

and finally a homomorphism of cohomology groups:

$$\begin{array}{ccc} \psi_s^{t,(p)} : H^p(K_t; G) & \longrightarrow & H^p(K_s; G) \\ [f]^{t,(p)} & \longmapsto & [(i_{s,t})^{*,p} f]^{s,(p)} \end{array}$$

The  $p$ -th persistent cohomology refers to the family of cohomology groups together with the transition homomorphisms between them:

$$\text{PH}^p(\{K_t\}_{t \in \mathbb{R}}; G) := \left( H^p(K_t; G), \psi_s^{t,(p)} \right)_{s \leq t \in \mathbb{R}}$$

where  $\psi_s^{t,(p)} : H^p(K_t) \rightarrow H^p(K_s)$  is the pullback homomorphism on cohomology induced by the inclusion  $K_s \hookrightarrow K_t$ .

In addition, Defitions 2.26,2.27,2.28, Lemma 2.29,2.30 are supplied as preliminary preparation.

*Definition 2.26* (Nerve of an Open Cover). Let  $X$  be a topological space and  $\mathcal{U} = \{U_i\}_{i \in I}$  an open cover of  $X$ . The **nerve** of the cover  $\mathcal{U}$  is the abstract simplicial complex  $\mathcal{N}(\mathcal{U})$  defined by:

1. Vertex set:  $V = \{v_i : i \in I\}$ , where each vertex  $v_i$  corresponds to the open set  $U_i$ .

2. **Simplices:** A finite subset  $\{v_{i_0}, v_{i_1}, \dots, v_{i_k}\} \subseteq V$  forms a  $k$ -simplex if and only if the intersection of the corresponding open sets is nonempty:

$$U_{i_0} \cap U_{i_1} \cap \dots \cap U_{i_k} \neq \emptyset.$$

**Definition 2.27** (Contractible Space). A topological space  $X$  is called **contractible space** or **good space** if identity map  $\text{id}_X: X \rightarrow X$  is homotopic to a constant map

**Definition 2.28** (Geometric Realization). Let  $K$  be an (abstract) simplicial complex. The **geometric realization**  $|K|$  of  $K$  is the topological space obtained by taking a standard geometric  $n$ -simplex  $\Delta_\sigma$  for each  $n$ -simplex  $\sigma \in \Sigma$ , and for each inclusion  $\tau \subseteq \sigma$ , identifying  $\Delta_\tau$  with a face of  $\Delta_\sigma$  via an affine homeomorphism. The union of these simplices is then given the weak topology

**Lemma 2.29** (Nerve Lemma). *Let  $\mathcal{U} = \{U_i\}_{i \in I}$  be an open cover of a topological space  $X$ . If every nonempty finite intersection  $U_{i_0} \cap U_{i_1} \cap \dots \cap U_{i_k}$  is good (i.e. contractible), then the geometric realization of  $\mathcal{N}(\mathcal{U})$  is homotopy equivalent to  $X$ :*

$$|\mathcal{N}(\mathcal{U})| \simeq X.$$

*Proof.* See [Hat02] Appendix Corollary 4G.3. □

**Lemma 2.30** (Čech Cover Approximation Lemma). *Let  $K$  be a compact subset of  $\mathbb{R}^D$ , let  $\mathcal{U} = \{B(z, \varepsilon) \mid z \in K\}$  be the  $\varepsilon$ -ball cover of  $K$ , and let  $\mathcal{V} = \{B(z_n, \varepsilon) \mid z_n \in Z\}$  be a “sample subcover” of  $\mathcal{U}$  with  $Z \subset K$ . If  $Z$  is  $\rho$ -dense in  $K$ , i.e.,  $\forall z \in K, \exists z_n \in Z$  such that  $d(z, z_n) < \rho$ , and  $\rho < \varepsilon/2$ , then  $\mathcal{V}$  is an open cover of  $K$  and  $|\mathcal{N}(\mathcal{U})| \simeq |\mathcal{N}(\mathcal{V})|$ .*

*Proof.* Let  $x \in K$ . By  $\rho$ -density, there exists  $z_n \in Z$  such that  $d(x, z_n) < \rho < \varepsilon/2$ . Then  $x \in B(z_n, \varepsilon)$ , so  $K \subseteq \bigcup_{z_n \in Z} B(z_n, \varepsilon) = \bigcup \mathcal{V}$ . Note that every nonempty finite intersection of balls in  $\mathcal{U}$  or  $\mathcal{V}$  is a convex set, hence contractible.

By Lemma 2.29, we have  $|\mathcal{N}(\mathcal{U})| \simeq K$ ,  $|\mathcal{N}(\mathcal{V})| \simeq K$ , then  $|\mathcal{N}(\mathcal{U})| \simeq |\mathcal{N}(\mathcal{V})|$  □

## 2.2 ČECH COHOMOLOGY AND SHEAVES

In this subsection, we present some fundamental knowledge of geometry and sheaf theory, which helps readers better comprehend the subsequent chapters.

**Definition 2.31** (Presheaf and Sheaf). Let  $X$  be a topological space. A **presheaf**  $\mathcal{F}$  on  $X$  is an assignment that to every nonempty open set<sup>1</sup>  $U$  on  $X$ , associates a set  $\mathcal{F}(U)$  and to every inclusion  $V \subseteq U$ , associates a homomorphism, called the restriction homomorphism  $r_V^U: \mathcal{F}(U) \rightarrow \mathcal{F}(V)$ , such that

$$1. \ r_U^U = \text{id}.$$

$$2. \ \text{For any } W \subseteq V \subseteq U, \text{ we have } r_W^U = r_W^V \circ r_V^U.$$

Elements of  $\mathcal{F}(U)$  are called sections of  $\mathcal{F}$  over  $U$ .

Let  $\mathcal{F}, \mathcal{G}$  be presheaves on  $X$ . A morphism  $h$  from  $\mathcal{F}$  to  $\mathcal{G}$  assigns to each nonempty open set  $U$  a homomorphism  $h_U: \mathcal{F}(U) \rightarrow \mathcal{G}(U)$  such that for any  $V \subseteq U$ , the following diagram commutes:

---

<sup>1</sup>More generally, an abelian group, a ring, etc.

$$\begin{array}{ccc}
\mathcal{F}(U) & \xrightarrow{h_U} & \mathcal{G}(U) \\
\downarrow r_V^U & & \downarrow \rho_V^U \\
\mathcal{F}(V) & \xrightarrow{h_V} & \mathcal{G}(V)
\end{array}$$

Let  $X$  be a topological space. A **sheaf**  $\mathcal{F}$  on  $X$  is a presheaf on  $X$  satisfying

1. If  $U$  is an open set in  $X$  and  $U = \bigcup_{i \in I} U_i$  is an open cover of  $U$ , then for any two sections  $s, t \in \mathcal{F}(U)$

$$r_{U_i}^U(s) = r_{U_i}^U(t), \forall i \in I \implies s = t$$

2. If  $U$  is an open set in  $X$  and  $U = \bigcup_{i \in I} U_i$  is an open cover of  $U$ , then for any family of sections  $s_i \in \mathcal{F}(U_i)$

$$r_{U_i \cap U_j}^{U_i}(s_i) = r_{U_i \cap U_j}^{U_j}(s_j), \forall i, j \in I \text{ with } U_i \cap U_j \neq \emptyset$$

then exists a global section  $s \in \mathcal{F}(U)$  such that  $r_{U_i}^U(s) = s_i, \forall i \in I$

*Definition 2.32* (Sheafification). For any  $p \in X$ , the stalk  $\mathcal{F}_p$  of a presheaf  $\mathcal{F}$  on  $X$  at  $p$  is defined by

$$\mathcal{F}_p := \varinjlim_{U \ni p} \mathcal{F}(U) := \coprod_{U \ni p} \mathcal{F}(U) / \sim$$

where  $s \in \mathcal{F}(U) \sim t \in \mathcal{F}(V)$  if and only if there exists an open set  $W \subseteq U \cap V$  such that  $r_W^U(s) = r_W^V(t)$ .

Elements of  $\mathcal{F}_p$  are called germs.

For any  $U \ni p$ , define the map  $\mathcal{F}(U) \rightarrow \mathcal{F}_p$ ,  $f \mapsto f_p := [f]$ .

The **sheafification** of a presheaf  $\mathcal{F}$  is defined by

$$\mathcal{F}^\#(U) := \left\{ s: U \rightarrow \prod_{p \in U} \mathcal{F}_p \left| \begin{array}{l} \forall q \in U, \exists \text{ an open neighborhood } V \subseteq U \text{ and a section} \\ f \in \mathcal{F}(V) \text{ such that } s(\xi) = f_\xi \text{ for all } \xi \in V \end{array} \right. \right\}$$

with restriction homomorphisms  $r_V^U := s|_V$ . Then  $\mathcal{F}^\#$  is a sheaf.

Naturally, one defines a morphism  $\theta: \mathcal{F} \rightarrow \mathcal{F}^\#$  from the presheaf to its sheafification, such that for any open set  $U$ ,

$$\theta_U: \mathcal{F}(U) \rightarrow \mathcal{F}^\#(U), \quad s \mapsto \left( \begin{array}{ccc} \tilde{s}: U & \longrightarrow & \prod_{p \in U} \mathcal{F}_p \\ p & \longmapsto & s_p \end{array} \right)$$

*Remark 2.33.* 1. If  $\mathcal{F}$  is a sheaf, then  $\theta$  is an isomorphism.

2. Let  $\mathcal{G}$  be a sheaf. For any morphism  $h: \mathcal{F} \rightarrow \mathcal{G}$ , there exists a unique  $\tilde{h}: \mathcal{F}^\# \rightarrow \mathcal{G}$  such that the following diagram commutes.

$$\begin{array}{ccc}
\mathcal{F} & \xrightarrow{h} & \mathcal{G} \\
\searrow \theta & & \nearrow \tilde{h} \\
& \mathcal{F}^\# &
\end{array}$$

*Definition 2.34* (Čech Cohomology of a Cover). Let  $\mathcal{F}$  be a sheaf on  $X$ , and let  $\mathcal{U} = \{U_i\}_{i \in I}$  be an open cover of  $X$ . For a multi-index  $(i_0, \dots, i_p) \in I^{p+1}$ , write  $U_{i_0, \dots, i_p} := U_{i_0} \cap \dots \cap U_{i_p}$ . Define

$$C^p(\mathcal{U}, \mathcal{F}) := \prod_{(i_0, \dots, i_p) \in I^{p+1}} \mathcal{F}(U_{i_0, \dots, i_p}),$$

concretely, for  $\sigma \in C^p(\mathcal{U}, \mathcal{F})$ , we have  $\sigma = \{\sigma_{i_0, \dots, i_p} : \sigma_{i_0, \dots, i_p} \in \mathcal{F}(U_{i_0, \dots, i_p})\}$ .

Define the coboundary operator  $\delta : C^p(\mathcal{U}, \mathcal{F}) \rightarrow C^{p+1}(\mathcal{U}, \mathcal{F})$ ,  $\sigma \mapsto \delta\sigma$ , where

$$(\delta\sigma)_{i_0, \dots, i_{p+1}} := \sum_{j=0}^{p+1} (-1)^j r_{U_{i_0, \dots, i_{p+1}}}^{U_{i_0, \dots, \hat{i}_j, \dots, i_{p+1}}} (\sigma_{i_0, \dots, \hat{i}_j, \dots, i_{p+1}}) =: \sum_{j=0}^{p+1} (-1)^j \sigma_{i_0, \dots, \hat{i}_j, \dots, i_{p+1}}|_{U_{i_0, \dots, i_{p+1}}}.$$

Set  $Z^p(\mathcal{U}, \mathcal{F}) := \ker \delta \subseteq C^p(\mathcal{U}, \mathcal{F})$ ,  $B^p(\mathcal{U}, \mathcal{F}) := \delta C^{p-1}(\mathcal{U}, \mathcal{F})$ . Then the **Čech cohomology groups** of the cover  $\mathcal{U}$  are defined by

$$\check{H}^p(\mathcal{U}; \mathcal{F}) := Z^p(\mathcal{U}, \mathcal{F}) / B^p(\mathcal{U}, \mathcal{F}).$$

If  $\mathcal{V} = \{V_\alpha\}_{\alpha \in \Gamma}$  is a refinement of  $\mathcal{U} = \{U_i\}_{i \in I}$ , choose a map  $\tau : \Gamma \rightarrow I$  such that  $V_\alpha \subseteq U_{\tau(\alpha)}$ . Then one defines a homomorphism to the refinement

$$\Phi_{\mathcal{V}}^{\mathcal{U}} : \check{H}^p(\mathcal{U}; \mathcal{F}) \rightarrow \check{H}^p(\mathcal{V}; \mathcal{F}), \quad [\sigma] = [\{\sigma_{i_0, \dots, i_p}\}] \mapsto [\eta] = [\{\eta_{\alpha_0, \dots, \alpha_p}\}]$$

where  $\eta_{\alpha_0, \dots, \alpha_p} \in \mathcal{F}(V_{\alpha_0, \dots, \alpha_p}) \subseteq \mathcal{F}(U_{\tau(\alpha_0), \dots, \tau(\alpha_p)})$ .

*Definition 2.35* (Čech Cohomology of a Space). Furthermore, the **Čech cohomology groups** of a space  $X$  are defined by

$$\check{H}^p(X; \mathcal{F}) := \varinjlim_{\mathcal{U}} \check{H}^p(\mathcal{U}; \mathcal{F}) := \prod_{\mathcal{U}} \check{H}^p(\mathcal{U}; \mathcal{F}) \Big/ \sim$$

where the equivalence relation is given by:

$$[\sigma] \in \check{H}^p(\mathcal{U}; \mathcal{F}) \sim [\eta] \in \check{H}^p(\mathcal{V}; \mathcal{F}) \iff \exists \text{ a common refinement } \mathcal{W} \text{ such that } \Phi_{\mathcal{W}}^{\mathcal{U}}([\sigma]) = \Phi_{\mathcal{W}}^{\mathcal{V}}([\eta]).$$

A sequence of sheaves is called exact if the induced sequence on stalks is exact at every point.

*Definition 2.36* (Resolution). Let  $\mathcal{F}$  be a sheaf on a topological space  $X$ . A **resolution** of  $\mathcal{F}$  is an exact sequence

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{S}^0 \rightarrow \mathcal{S}^1 \rightarrow \mathcal{S}^2 \rightarrow \dots$$

where each  $\mathcal{S}^p$  is a sheaf on  $X$ .

*Definition 2.37* (Acyclic Resolution). Let  $\mathcal{F}$  be a sheaf on a topological space  $X$ . A resolution  $0 \rightarrow \mathcal{F} \rightarrow \mathcal{S}^\bullet$  is called **acyclic** if each  $\mathcal{S}^p$  is acyclic, i.e.,  $\check{H}^i(X; \mathcal{S}^p) = 0$  for all  $i > 0$ .

*Definition 2.38* (Fine Sheaf). Let  $\mathcal{F}$  be a sheaf on a topological space  $X$ .  $\mathcal{F}$  is called a **fine sheaf** if for any locally finite open cover  $\{U_i\}$  of  $X$ , there exists a partition of unity  $\{e_i\}$  subordinate to the cover, i.e.,

1.  $\text{supp}(e_i) \subseteq U_i$ .
2.  $\sum_i e_i = 1$ .

**Theorem 2.39** (Abstract de Rham Theorem). *Let  $\mathcal{F}$  be a sheaf on a topological space  $X$ , and let  $0 \rightarrow \mathcal{F} \rightarrow \mathcal{S}^\bullet$  be an acyclic resolution of  $\mathcal{F}$  such that each  $\mathcal{S}^p$  is a fine sheaf. Then the Čech cohomology  $\check{H}^p(X, \mathcal{F})$  of  $\mathcal{F}$  is isomorphic to the  $p$ -th cohomology of the complex  $\Gamma(X, \mathcal{S}^\bullet)$ :*

$$\check{H}^p(X, \mathcal{F}) \cong \ker(\Gamma(X, \mathcal{S}^p) \rightarrow \Gamma(X, \mathcal{S}^{p+1})) / \operatorname{im}(\Gamma(X, \mathcal{S}^{p-1}) \rightarrow \Gamma(X, \mathcal{S}^p)).$$

*In particular, if  $\mathcal{U}$  is an open cover of  $X$  such that for every finite nonempty intersection  $U_{i_0, \dots, i_k}$  of sets in  $\mathcal{U}$ , we have  $\check{H}^j(U_{i_0, \dots, i_k}; \mathcal{S}^p) = 0$  for all  $j > 0$ , then there is a natural isomorphism*

$$\check{H}^p(\mathcal{U}; \mathcal{F}) \cong \check{H}^p(X, \mathcal{F}).$$

*Proof.* See [Han25]. □

### 3 DYNAMIC ČECH COMPLEXES AND DYNAMIC ČECH SHEAVES

#### 3.1 THEORY OF DYNAMIC ČECH COMPLEXES

When dealing with dynamical systems, traditional TDA typically uses a discrete approach: select a discrete sequence  $\{t_1, \dots, t_m\}$  on the time axis, construct a static Čech complex at each time  $t_i$ , then compare topological features among these discrete complexes. However, this method is essentially a discrete approximation and cannot describe the continuous evolution of topological structures over the parameter space.

In this subsection, we propose a fundamental viewpoint: model the dynamical process as a continuous map from the parameter space  $T$  to the space of complexes  $\mathcal{S}(M)$ , thereby establishing a truly continuous dynamical theory.<sup>2</sup>

*Definition 3.1* (Continuous Complex Flow). Let the parameter space  $T$  be a connected topological space, and let  $(M, g)$  be a compact  $d$ -dimensional Riemannian manifold. A **continuous complex flow** is a continuous map:

$$\mathcal{C}: T \rightarrow \mathcal{S}(M)$$

where  $\mathcal{S}(M)$  denotes the topological space of all abstract simplicial complexes on  $M$ , endowed with the Whitney topology. That is, for any open set  $U \subseteq M$  and integer  $k \geq 0$ , a subbase of open sets is given by

$$\mathcal{U}(U, k) := \{K \in \mathcal{S}(M) \mid K \cap U \text{ contains at least one } k\text{-simplex}\}.$$

The core idea is to model the dynamical process as a “path” in the parameter space, whose continuous movement in  $\mathcal{S}(M)$  describes the continuous evolution of topological structure.

*Definition 3.2* (Parametrized Topological Space). Let the parameter space  $T$  be a connected topological space, and let  $M$  be a compact  $d$ -dimensional manifold. A **parametrized topological space** is a trivial bundle  $\pi: E = M \times T \rightarrow T$  satisfying fiber isomorphism  $\pi^{-1}(t) \cong M$ .

*Definition 3.3* (Dynamic Family of Embeddings). A **dynamic family of embeddings** is a smooth map  $\Phi: E = M \times T \rightarrow \mathbb{R}^D$  such that

1.  $\Phi_t := \Phi(\cdot, t): M \rightarrow \mathbb{R}^D$  is an embedding;
2. There exists  $c > 0$  such that, for every  $t \in T$ , the norm of the Jacobian matrix of  $\Phi_t$  satisfies  $\|\mathbf{J}\Phi_t\| \leq c$  for all  $x \in M$ ;
3. The map  $t \mapsto \Phi_t$  is  $C^\infty$ .

Within the abstract framework of continuous complex flows, we require concrete geometric realizations. The dynamic Čech continuous flow serves as precisely such a bridge.

*Definition 3.4* (Dynamic Čech Continuous Flow). Given a dynamic family of embeddings  $\Phi: M \times T \rightarrow \mathbb{R}^D$  and a radius parameter  $\varepsilon > 0$ , the **dynamic Čech continuous flow** is defined by

$$\begin{aligned} \mathcal{C}_{\Phi, \varepsilon}: T &\longrightarrow \mathcal{S}(M) \\ t &\longmapsto \left\{ \sigma \subseteq M : \bigcap_{x \in \sigma} B(\Phi_t(x), \varepsilon) \neq \emptyset \right\} \end{aligned}$$

---

<sup>2</sup>Although real-world measured data is discrete, it can first be extended to continuous form using numerical methods, and then analyzed within the framework of continuous dynamic theory. This is a better method compared with traditional discrete approximation.

**Theorem 3.5** (Continuity Guarantee). *If the dynamic family of embeddings  $\Phi$  is continuous, then the dynamic Čech continuous flow  $\mathcal{C}_{\Phi,\varepsilon}$  is continuous with respect to the Whitney topology.*

*Proof.* Note that the Whitney topology on  $\mathcal{S}(M)$  is generated by the sets  $\mathcal{U}(U, k)$ . Thus to prove continuity of  $\mathcal{C}_{\Phi,\varepsilon}$ , it suffices to show that for every  $\mathcal{U}(U, k)$ , its preimage

$$\mathcal{C}_{\Phi,\varepsilon}^{-1}(\mathcal{U}(U, k)) := \{t \in T \mid \mathcal{C}_{\Phi,\varepsilon}(t) \in \mathcal{U}(U, k)\}$$

is open in  $T$ .

Let  $t_0 \in \mathcal{C}_{\Phi,\varepsilon}^{-1}(\mathcal{U}(U, k))$ . Then there exists a  $k$ -simplex  $\sigma = \{x_1, \dots, x_{k+1}\} \subseteq U$  in  $\mathcal{C}_{\Phi,\varepsilon}(t_0)$ , so there exists  $y \in \mathbb{R}^D$  such that  $y \in \bigcap_{i=1}^{k+1} B(\Phi_{t_0}(x_i), \varepsilon)$ , i.e.,  $d(\Phi_{t_0}(x_i), y) < \varepsilon$  for all  $i$ .

By continuity of  $\Phi$ , for each  $x_i$ , the map  $t \mapsto \Phi_t(x_i)$  is continuous at  $t_0$ . Hence there exists  $\delta_i > 0$  such that whenever  $d_T(t, t_0) < \delta_i$ ,

$$d_{\mathbb{R}^D}(\Phi_t(x_i), \Phi_{t_0}(x_i)) < \eta, \quad \eta = \frac{1}{2}(\varepsilon - \max_i d(\Phi_{t_0}(x_i), y)) > 0.$$

Set  $\delta = \min\{\delta_i \mid 1 \leq i \leq k+1\}$ . Then for  $d_T(t, t_0) < \delta$ , for every  $i$ ,

$$d(\Phi_t(x_i), y) \leq d(\Phi_t(x_i), \Phi_{t_0}(x_i)) + d(\Phi_{t_0}(x_i), y) < \eta + d(\Phi_{t_0}(x_i), y) < \varepsilon.$$

Thus  $y \in \bigcap_{i=1}^{k+1} B(\Phi_t(x_i), \varepsilon)$ , so  $\sigma \in \mathcal{C}_{\Phi,\varepsilon}(t)$ . Since  $\sigma \subseteq U$ , we have  $\mathcal{C}_{\Phi,\varepsilon}(t) \in \mathcal{U}(U, k)$ , so  $B_{d_T}(t_0, \delta) \subseteq \mathcal{C}_{\Phi,\varepsilon}^{-1}(\mathcal{U}(U, k))$ . This shows that the preimage is open.  $\square$

This result shows that continuity of the parameter space naturally descends to continuity of the complex structure. In what follows, we will gradually see that this preservation of continuity has profound implications.

To understand dynamical topological processes at a more abstract level, we lift the concept of continuous complex flow to a categorical setting. This lifting is not merely a formal nicety, but reveals the inherent algebraic structure of dynamical topological processes, providing a natural framework for subsequent sheaf-theoretic constructions.

In traditional static TDA, a Čech complex is viewed as an object in the category  $\mathbf{Simp}$ . While concise, this viewpoint cannot capture the dynamical behavior of complexes varying with parameters. To describe such dynamics, we elevate our perspective from “objects” to “functors”. Thus we uniformly define a dynamic Čech complex as a functor from the parameter space to the category of simplicial complexes, a construction that simultaneously encodes the continuous evolution of topology and its algebraic structure.

**Definition 3.6** (Dynamic Čech Complex Functor). Let  $(M, g)$  be a compact  $d$ -dimensional Riemannian manifold, and let the parameter space  $T$  be a connected topological space. Given a dynamic family of embeddings  $\Phi: M \times T \rightarrow \mathbb{R}^D$  and a radius parameter  $\varepsilon > 0$ , the **dynamic Čech complex functor** is defined by:

$$\mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}} : T \rightarrow \mathbf{Simp}$$

where:

1. For any  $t \in T$ ,  $\mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}}(t) := \left\{ \sigma \subseteq M : \bigcap_{x \in \sigma} B(\Phi_t(x), \varepsilon) \neq \emptyset \right\}$ .
2. For any  $s, t \in T$  and path  $\gamma : s \rightarrow t$ , i.e.,  $\gamma : [0, 1] \rightarrow T$  with  $\gamma(0) = s$ ,  $\gamma(1) = t$ ,  $\mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}}(\gamma) : \mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}}(s) \rightarrow \mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}}(t)$  is the induced simplicial map.

This unified construction has a dual significance:

- From the continuous flow viewpoint, regarding  $\mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}}$  as a map from  $T$  to the complex space  $\mathcal{S}(M)$ , it describes the continuous evolution of topological structure, capturing the infinitesimal changes of topological features in the dynamical process (see later sections).
- From the functorial viewpoint, regarding  $\mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}}$  as a functor from the parameter category  $T$  to the simplicial complex category **Simp**, it reveals the algebraic structure of the dynamical process, providing a natural framework for subsequent sheaf and cohomology theories.

In practical applications, we typically only have access to a finite sample set of the manifold  $M$ . This motivates the sampled version of the definition:

*Definition 3.7* (Sampled Dynamic Čech Complex Functor). Let  $X_n \subset M$  be a finite sample set. The **sampled dynamic Čech complex functor** is defined by:

$$\mathcal{C}_{\Phi,\varepsilon}^{\text{sm}} : T \rightarrow \text{Simp}$$

where for any  $t \in T$ ,

$$\mathcal{C}_{\Phi,\varepsilon}^{\text{sm}}(t) := \left\{ \sigma \subseteq X_n \mid \bigcap_{x \in \sigma} B(\Phi_t(x), \varepsilon) \neq \emptyset \right\},$$

and for any path  $\gamma : s \rightarrow t$ ,  $\mathcal{C}_{\Phi,\varepsilon}^{\text{sm}}(\gamma)$  is the corresponding simplicial map.

**Proposition 3.8** (Restriction Relation). *The sampled dynamic Čech complex functor is the restriction of the full dynamic Čech complex functor to the sample set  $X_n$ :*

$$\mathcal{C}_{\Phi,\varepsilon}^{\text{sm}} = \mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}} \Big|_{X_n}.$$

That is, for any  $t \in T$ ,  $\mathcal{C}_{\Phi,\varepsilon}^{\text{sm}}(t)$  is the subcomplex of  $\mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}}(t)$  generated by the sample set  $X_n$ .

*Proof.* This follows directly from the definition: the simplices of  $\mathcal{C}_{\Phi,\varepsilon}^{\text{sm}}(t)$  are exactly those subsets of simplices in  $\mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}}(t)$  whose vertex set is contained in  $X_n$ .  $\square$

**Theorem 3.9** (Approximation Property). *If the sample set  $X_n$  is  $\delta$ -dense in  $M$ , i.e., for every  $x \in M$ , there exists  $x_i \in X_n$  such that  $d_M(x, x_i) < \delta$ , then as  $\delta \rightarrow 0$ , the sampled dynamic Čech complex functor converges to the full dynamic Čech complex functor in an appropriate topological sense:*

$$\mathcal{C}_{\Phi,\varepsilon}^{\text{sm}} \rightarrow \mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}} \quad (\delta \rightarrow 0).$$

In particular, for any  $t \in T$ , there is a homotopy equivalence

$$|\mathcal{C}_{\Phi,\varepsilon}^{\text{sm}}(t)| \simeq |\mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}}(t)|, \quad \text{for sufficiently small } \delta.$$

*Proof.* Fix  $t \in T$  and consider  $\Phi_t : M \rightarrow \mathbb{R}^D$ . For any  $x, y \in M$ , since  $M$  is a compact Riemannian manifold, there exists a smooth curve  $\gamma : [0, 1] \rightarrow M$  connecting  $x$  and  $y$  with  $\gamma(0) = x$ ,  $\gamma(1) = y$ . Then

$$\Phi_t(y) - \Phi_t(x) = \int_0^1 \mathbf{J}\Phi_t(\gamma(s)) \cdot \gamma'(s) \, ds.$$



Taking the Euclidean norm,

$$d_{\mathbb{R}^D}(\Phi_t(x), \Phi_t(y)) \leq \int_0^1 \|\mathbf{J}\Phi_t(\gamma(s))\| \cdot \|\gamma'(s)\| ds \leq c \int_0^1 \|\gamma'(s)\| ds = c \cdot d_M(x, y).$$

Hence  $\Phi_t$  is Lipschitz continuous with Lipschitz constant  $c$ .

If  $X_n \subset M$  is  $\delta$ -dense, then for any  $x \in M$  and  $t \in T$ ,

$$d_{\mathbb{R}^D}(\Phi_t(x), \Phi_t(x_n)) \leq c \cdot d_M(x, x_n) < c\delta.$$

That is,  $\Phi_t(X_n)$  is  $c\delta$ -dense in  $\Phi_t(M)$ .

Fix  $t \in T$  and consider the following two open covers:

1. The full cover  $\mathcal{U}_t := \{B(\Phi_t(x), \varepsilon) \mid x \in M\}$ , whose nerve is  $\mathcal{N}(\mathcal{U}_t) = \mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(t)$ .
2. The sampled cover  $\mathcal{U}_t^{\text{smp}} := \{B(\Phi_t(x), \varepsilon) \mid x \in X_n\}$ , whose nerve is  $\mathcal{N}(\mathcal{U}_t^{\text{smp}}) = \mathcal{C}_{\Phi, \varepsilon}^{\text{smp}}(t)$ .

We show that if  $\delta < \varepsilon/(2c)$ , then  $|\mathcal{N}(\mathcal{U}_t)| \simeq |\mathcal{N}(\mathcal{U}_t^{\text{smp}})|$ .

Define a map  $f : |\mathcal{N}(\mathcal{U}_t)| \rightarrow |\mathcal{N}(\mathcal{U}_t^{\text{smp}})|$  as follows. For any point  $y \in |\mathcal{N}(\mathcal{U}_t)|$ , write its barycentric coordinate representation as  $y = \sum_{x \in \sigma} \lambda_x \cdot v_x$ , where  $\sigma = \{x_0, \dots, x_k\} \in \mathcal{N}(\mathcal{U}_t)$ , so

$\bigcap_{x \in \sigma} B(\Phi_t(x), \varepsilon) \neq \emptyset$ ,  $\lambda_x \geq 0$ ,  $\sum \lambda_x = 1$ , and  $v_x$  is the vertex corresponding to  $x$ .

By  $\delta$ -density of  $X_n$ , for each  $x \in \sigma$ , choose  $\pi(x) \in X_n$  such that  $d_M(x, \pi(x)) < \delta$ . Define  $f(y) = \sum_{x \in \sigma} \lambda_x \cdot v_{\pi(x)}$ , where  $v_{\pi(x)}$  is the vertex in  $\mathcal{N}(\mathcal{U}_t^{\text{smp}})$  corresponding to  $\pi(x)$ .

We verify that  $f$  is well-defined, i.e., that  $\{\pi(x) \mid x \in \sigma\}$  forms a simplex in  $\mathcal{N}(\mathcal{U}_t^{\text{smp}})$ . Since  $\sigma \in \mathcal{N}(\mathcal{U}_t)$ , there exists  $z \in \bigcap_{x \in \sigma} B(\Phi_t(x), \varepsilon)$ , so  $d_{\mathbb{R}^D}(\Phi_t(x), z) < \varepsilon$ . For any  $x \in \sigma$ ,

$$d_{\mathbb{R}^D}(\Phi_t(\pi(x)), z) \leq d_{\mathbb{R}^D}(\Phi_t(\pi(x)), \Phi_t(x)) + d_{\mathbb{R}^D}(\Phi_t(x), z) < c\delta + \varepsilon.$$

Since  $\delta < \varepsilon/(2c)$ , we have  $d_{\mathbb{R}^D}(\Phi_t(\pi(x)), z) < 3\varepsilon/2$ . Thus  $\bigcap_{\pi(x) \in \pi(\sigma)} B(\Phi_t(\pi(x)), 3\varepsilon/2) \neq \emptyset$ .

Combined with classical Čech approximation theory, for sufficiently small  $\delta$ ,  $\pi(\sigma)$  is a simplex in  $\mathcal{N}(\mathcal{U}_t^{\text{smp}})$ .

Define a map  $g : |\mathcal{N}(\mathcal{U}_t^{\text{smp}})| \rightarrow |\mathcal{N}(\mathcal{U}_t)|$  as follows. For each vertex  $v_{x_n}$  in the sampled nerve ( $x_n \in X_n$ ), set  $g(v_{x_n}) = v_{x_n}$  and extend linearly over simplices.

We obtain a homotopy equivalence:

1.  $g \circ f \sim \text{id}_{|\mathcal{N}(\mathcal{U}_t)|}$ : construct a homotopy  $H(s, y) = \sum \lambda_x \cdot (sv_x + (1-s)v_{\pi(x)})$ . Since  $x$  and  $\pi(x)$  are sufficiently close, the simplex persists throughout the deformation.
2.  $f \circ g \sim \text{id}_{|\mathcal{N}(\mathcal{U}_t^{\text{smp}})|}$ : similar homotopy, using density of the sample to ensure continuity.

Hence  $|\mathcal{N}(\mathcal{U}_t)| \simeq |\mathcal{N}(\mathcal{U}_t^{\text{smp}})|$ .

Therefore, as  $\delta \rightarrow 0$ ,  $\mathcal{C}_{\Phi, \varepsilon}^{\text{smp}} \rightarrow \mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}$ , and for any  $t \in T$ ,  $|\mathcal{C}_{\Phi, \varepsilon}^{\text{smp}}(t)| \simeq |\mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(t)|$ .  $\square$

**Theorem 3.10** (Continuity Guarantee). *If the dynamic family of embeddings  $\Phi$  is continuous, then the dynamic Čech complex functor  $\mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}$  and the sampled dynamic Čech complex functor  $\mathcal{C}_{\Phi, \varepsilon}^{\text{smp}}$  are continuous in the following sense:*

1. As maps from  $T$  to  $\mathcal{S}(M)$ , they are continuous in the Whitney topology.
2. As functors from  $T$  to  $\text{Simp}$ , they preserve the path structure of the parameter space.

*Proof.* The proof of topological continuity is analogous to earlier arguments. We prove functorial continuity (i.e., as functors from  $T$  to  $\mathbf{Simp}$ , they preserve the path structure of the parameter space); the sampled version is similar.

Preservation of identities is clear: for any  $t \in T$ , the constant path  $\text{id}_t : t \rightarrow t$  induces the identity simplicial map:

$$\mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(\text{id}_t) = \text{id}_{\mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(t)}.$$

Let  $\gamma : s \rightarrow t$  and  $\delta : t \rightarrow u$  be paths in  $T$  (assumed at least  $C^1$ ). That is,  $\gamma : [0, 1] \rightarrow T$  with  $\gamma(0) = s$ ,  $\gamma(1) = t$ , and  $\delta : [0, 1] \rightarrow T$  with  $\delta(0) = t$ ,  $\delta(1) = u$ . Define the composite path  $\delta \circ \gamma : [0, 1] \rightarrow T$  by

$$(\delta \circ \gamma)(\tau) = \begin{cases} \gamma(2\tau), & \tau \in [0, 1/2], \\ \delta(2\tau - 1), & \tau \in (1/2, 1]. \end{cases}$$

We show composition is preserved:

$$\mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(\delta \circ \gamma) = \mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(\delta) \circ \mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(\gamma).$$

For the path  $\gamma : s \rightarrow t$ , consider the simplicial map  $\mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(\gamma) : \mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(s) \rightarrow \mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(t)$ . For each vertex  $x \in M$ , since  $\Phi$  is  $C^\infty$ ,

$$\frac{d}{d\tau} \Phi_{\gamma(\tau)}(x) = \frac{\partial \Phi}{\partial t}(x, \gamma(\tau)) \cdot \gamma'(\tau).$$

With initial condition  $\Phi_{\gamma(0)}(x) = \Phi_s(x)$ , there is a unique solution  $\Phi_{\gamma(\tau)}(x)$ , depending continuously on  $\tau$ .

Thus for each vertex  $x \in \mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(s)$ , we define its image to be  $x$  itself (the vertex set is unchanged), giving the identity map on vertices.

For a simplex  $\sigma = \{x_1, \dots, x_k\} \in \mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(s)$ , we determine its image. Since the vertex map is the identity, the vertex set of  $\sigma$  maps to  $\{x_1, \dots, x_k\}$ . However,  $\{x_1, \dots, x_k\}$  may no longer be a simplex at time  $t$ .

We define: the image of  $\sigma$  under the simplicial map is  $\{x_1, \dots, x_k\}$  if and only if  $\sigma$  remains a simplex along the entire path  $\gamma$ , i.e.,

$$\bigcap_{i=1}^k B(\Phi_{\gamma(\tau)}(x_i), \varepsilon) \neq \emptyset, \quad \forall \tau \in [0, 1].$$

If this holds, set  $\mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(\gamma)(\sigma) = \sigma$ ; otherwise, take the maximal subsimplex  $\tau \subset \sigma$  that remains a simplex along  $\gamma$ , and set  $\mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(\gamma)(\sigma) = \tau$ .

We now prove composition is preserved. For any simplex  $\sigma \in \mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(s)$ ,

$$\begin{aligned} \mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(\delta \circ \gamma)(\sigma) = \sigma &\iff \bigcap_{i=1}^k B(\Phi_{(\delta \circ \gamma)(\tau)}(x_i), \varepsilon) \neq \emptyset, \quad \forall \tau \in [0, 1], \\ \mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(\delta) \circ \mathcal{C}_{\Phi, \varepsilon}^{\text{dyn}}(\gamma)(\sigma) = \sigma &\iff \begin{cases} \bigcap_{i=1}^k B(\Phi_{\gamma(\tau)}(x_i), \varepsilon) \neq \emptyset, & \forall \tau \in [0, 1], \\ \bigcap_{i=1}^k B(\Phi_{\delta(\tau)}(x_i), \varepsilon) \neq \emptyset, & \forall \tau \in [0, 1]. \end{cases} \end{aligned}$$

By definition of the composite path, these two conditions are equivalent. Therefore,

$$\mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}}(\delta \circ \gamma)(\sigma) = \sigma \iff \mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}}(\delta) \circ \mathcal{C}_{\Phi,\varepsilon}^{\text{dyn}}(\gamma)(\sigma) = \sigma,$$

so the induced simplicial maps coincide, preserving composition. The case of degenerate sub-simplices is analogous and immediate.  $\square$

By uniformly defining the dynamic Čech complex as a functor and clearly distinguishing between the full and sampled versions, we have established a rigorous and practical theoretical framework, laying a solid mathematical foundation for dynamic TDA.

To gain deeper insight into the evolution mechanism of topological features, we go beyond continuity and study the infinitesimal behavior of evolution. Infinitesimal generation theory provides a differential description of topological evolution, allowing us to precisely characterize topological changes in the language of differential equations.

*Definition 3.11* (Tangent Space of the Complex Flow). Let  $\mathcal{S}(M)$  be the space of all simplicial complexes on a smooth manifold  $M$ , endowed with the Whitney topology. For any  $K \in \mathcal{S}(M)$ , let  $\mathcal{U}_K$  denotes the set of all open neighborhoods of  $K$  in  $\mathcal{S}(M)$ , which forms a directed set under the reverse inclusion order  $U_1 \leq U_2 \iff U_2 \subseteq U_1$ .

For each  $U \in \mathcal{U}_K$ , let  $K_U \in U$  be a simplicial complex. The **tangent space** of the space of complexes at  $K$  is defined as the inverse limit of the sets  $\text{Hom}(K, K_U)$  over the directed set  $\mathcal{U}_K$ :

$$T_K \mathcal{S}(M) := \varprojlim_{U \in \mathcal{U}_K} \text{Hom}(K, K_U)$$

An element  $v \in T_K \mathcal{S}(M)$  is called an infinitesimal deformation of  $K$ , which encodes the infinitesimal behavior of topological evolution of  $K$ ,

The notion of tangent space allows us to define derivatives and differentials on the space of complexes, providing tools for studying the instantaneous behavior of topological evolution. The next theorem shows that the “instantaneous topological change” (tangent vector) of a dynamic Čech complex flow is well-defined and is uniquely determined by the instantaneous velocity of the embedding map.

**Theorem 3.12** (Infinitesimal Description of Dynamic Complex Flows). *The dynamic Čech continuous flow  $\mathcal{C}_{\Phi,\varepsilon}$  is differentiable at  $t$ , and its tangent vector satisfies*

$$\frac{d}{dt} \mathcal{C}_{\Phi,\varepsilon}(t) \in T_{\mathcal{C}_{\Phi,\varepsilon}(t)} \mathcal{S}(M).$$

*This tangent vector is uniquely determined by the embedding velocity field  $v_t = \frac{\partial \Phi}{\partial t}$ , with the explicit expression:*

$$\frac{d}{dt} \mathcal{C}_{\Phi,\varepsilon}(t) = \mathcal{P} \left( \bigcup_{x \in M} d\Phi_t(x)(v_t(x)) \right),$$

where

$$\mathcal{P}: \mathcal{D}(\mathcal{P}) = \left\{ \bigcup_{x \in M} w_t(x) \middle| w_t(x) \in T_{\Phi_t(x)} \mathbb{R}^D \right\} \rightarrow T_{\mathcal{C}_{\Phi,\varepsilon}(t)} \mathcal{S}(M)$$

denotes the projection operator onto the tangent space.

*Proof.* Let  $K_t = \mathcal{C}_{\Phi, \varepsilon}(t) \in \mathcal{S}(M)$ . We expand the tangent space:

$$T_{K_t}\mathcal{S}(M) = \prod_{U \in \mathcal{U}_{K_t}} \text{Hom}(K_t, K_U) \Big/ \sim,$$

where  $\mathcal{U}_{K_t} = \{U \subset \mathcal{S}(M) \mid U \text{ is a Whitney open neighborhood of } K_t\}$ ,  $K_U \in U$  is any complex in  $U$ , and the equivalence relation  $\sim$  is

$$(f_U) \sim (g_U) \iff \exists V \in \mathcal{U}_{K_t}, \forall U \subseteq V, f_U = g_U.$$

The equivalence class is denoted  $[(f_U)] = \{(g_U) \in \prod \text{Hom}(K_t, K_U) \mid (g_U) \sim (f_U)\}$ .

Linearized velocity:  $w_t(x) = d\Phi_t(x)(v_t(x)) \in T_{\Phi_t(x)}\mathbb{R}^D \cong \mathbb{R}^D$ . We prove that  $\mathcal{P}(\bigcup w_t(x)) = [(f_U)]$  for some compatible family of homomorphisms  $(f_U)$ .

First, the complex flow is differentiable at  $t$  if and only if there exists a unique  $\dot{K}_t \in T_{K_t}\mathcal{S}(M)$  such that

$$\lim_{h \rightarrow 0} \frac{[(f_U^h)] - [(f_U^0)]}{h} = \dot{K}_t,$$

where  $f_U^h \in \text{Hom}(K_t, K_{t+h})$ ,  $f_U^0 = \text{id}_{K_t}$ , and the limit is taken in the quotient topology of  $T_{K_t}\mathcal{S}(M)$ .

We construct the induced family  $(f_U^h)$ . For any  $U \in \mathcal{U}_{K_t}$ , by  $\Phi \in C^\infty(M \times T, \mathbb{R}^D)$ , there exists  $\delta_U > 0$  such that for  $|h| < \delta_U$ ,  $\Phi_{t+h}(x) = \Phi_t(x) + h \cdot w_t(x) + o(h)$  and  $K_{t+h} \in U$ .

For any simplex  $\sigma = \{x_1, \dots, x_k\} \in K_t$ , there exists  $z \in \bigcap_{i=1}^k B(\Phi_t(x_i), \varepsilon)$ , so  $d(\Phi_t(x_i), z) < \varepsilon$ .

Then

$$d(\Phi_{t+h}(x_i), z) \leq d(\Phi_t(x_i), z) + h \cdot \|w_t(x_i)\| + \|o(h)\| < \varepsilon + hC + \|o(h)\|,$$

where  $C = \max_{x \in M} \|w_t(x)\| < \infty$  (since  $M$  is compact and  $w_t$  continuous). Set  $\delta = \min \left\{ \delta_U, \frac{\varepsilon - d(\Phi_t(x_i), z)}{2C} \right\}$ .

For  $|h| < \delta$ ,  $d(\Phi_{t+h}(x_i), z) < \varepsilon$ , so  $\sigma \in K_{t+h}$ .

Define  $f_U^h : K_t \rightarrow K_{t+h}$  to be the simplicial homomorphism with vertex map  $f_U^h(x) = x$  and simplex extension  $f_U^h(\sigma) = \sigma$  for all  $\sigma \in K_t$ . Clearly  $f_U^h \in \text{Hom}(K_t, K_{t+h})$ , so  $(f_U^h)_{U \in \mathcal{U}_{K_t}} \in \prod \text{Hom}(K_t, K_{t+h})$  when  $|h| < \delta$ .

We now establish the existence of the limiting homomorphism family  $(f_U)$ . Define  $(f_U) \in \prod \text{Hom}(K_t, K_U)$  such that, for any  $U \in \mathcal{U}_{K_t}$  and  $\sigma \in K_t$ ,

$$f_U(\sigma) = \lim_{h \rightarrow 0} \frac{f_U^h(\sigma) - \sigma}{h}.$$

We clarify the geometric realizations:

1. For any  $k$ -simplex  $\sigma = \{x_0, x_1, \dots, x_k\} \in K_t$ , its geometric realization is the convex set in  $\mathbb{R}^D$ :

$$|\sigma| = \left\{ \sum_{i=0}^k \lambda_i \Phi_t(x_i) \mid \lambda_i \geq 0, \sum_{i=0}^k \lambda_i = 1 \right\}.$$

2. Since  $f_U^h(\sigma) = \sigma$  (identity on vertices), its geometric realization is:

$$|f_U^h(\sigma)| = \left\{ \sum_{i=0}^k \lambda_i \Phi_{t+h}(x_i) \mid \lambda_i \geq 0, \sum_{i=0}^k \lambda_i = 1 \right\}.$$

3. The geometric meaning of the difference quotient  $\frac{f_U^h(\sigma) - \sigma}{h}$  is the “instantaneous rate of change” of the convex set:

$$\left| \frac{f_U^h(\sigma) - \sigma}{h} \right| = \left\{ \sum_{i=0}^k \lambda_i \cdot \frac{\Phi_{t+h}(x_i) - \Phi_t(x_i)}{h} \middle| \lambda_i \geq 0, \sum_{i=0}^k \lambda_i = 1 \right\}.$$

We use the Hausdorff distance to characterize convergence of convex sets. Recall that for two compact convex sets  $A, B \subset \mathbb{R}^D$ , the Hausdorff distance is

$$d_H(A, B) = \max \left\{ \sup_{a \in A} \inf_{b \in B} \|a - b\|, \sup_{b \in B} \inf_{a \in A} \|a - b\| \right\}.$$

A sequence of simplices  $\{\sigma_h\}$  converges to  $\sigma$  if and only if their geometric realizations satisfy  $d_H(|\sigma_h|, |\sigma|) \rightarrow 0$  as  $h \rightarrow 0$ .

First, convergence of the vertex difference quotients holds trivially. For any vertex  $x_i$  of  $\sigma$ , since  $\Phi \in C^\infty(M \times T, \mathbb{R}^D)$ , the Taylor expansion gives

$$\Phi_{t+h}(x_i) = \Phi_t(x_i) + h \cdot w_t(x_i) + o(h) \implies \exists \lim_{h \rightarrow 0} \frac{\Phi_{t+h}(x_i) - \Phi_t(x_i)}{h} = w_t(x_i).$$

This limit exists in the Euclidean topology of  $\mathbb{R}^D$ .

Next, convergence of the simplex difference quotients in the Hausdorff distance. We show  $\left| \frac{f_U^h(\sigma) - \sigma}{h} \right| \xrightarrow{d_H} |f_U(\sigma)|$ , where

$$|f_U(\sigma)| = \left\{ \sum_{i=0}^k \lambda_i w_t(x_i) \middle| \lambda_i \geq 0, \sum_{i=0}^k \lambda_i = 1 \right\}.$$

1. Pointwise convergence. For any  $y = \sum_{i=0}^k \lambda_i \Phi_t(x_i) \in |\sigma|$ , let  $y_h = \sum_{i=0}^k \lambda_i \Phi_{t+h}(x_i) \in |f_U^h(\sigma)|$ .

Then

$$\frac{y_h - y}{h} = \sum_{i=0}^k \lambda_i \cdot \frac{\Phi_{t+h}(x_i) - \Phi_t(x_i)}{h} \rightarrow \sum_{i=0}^k \lambda_i w_t(x_i) \in |f_U(\sigma)|.$$

2. Uniform convergence. Since  $|\sigma|$  is compact (convex hull of a compact set in  $\mathbb{R}^D$ ) and  $w_t : M \rightarrow \mathbb{R}^D$  is continuous, by compactness of  $M$ ,  $w_t$  is uniformly continuous. Hence the vertex difference quotients  $q_h(x_i) := \frac{\Phi_{t+h}(x_i) - \Phi_t(x_i)}{h}$  converge uniformly to  $w_t(x_i)$  for  $x_i \in \sigma$ .

Let

$$Q_h = \left\{ \sum_{i=0}^k \lambda_i q_h(x_i) \middle| \lambda_i \geq 0, \sum \lambda_i = 1 \right\},$$

$$Q = \left\{ \sum_{i=0}^k \lambda_i w_t(x_i) \middle| \lambda_i \geq 0, \sum \lambda_i = 1 \right\}.$$

(a) Take any  $a \in Q_h$ . By convexity, there exist  $\lambda_i \geq 0$ ,  $\sum \lambda_i = 1$  such that

$$a = \sum_{i=0}^k \lambda_i q_h(x_i).$$

Set  $b = \sum_{i=0}^k \lambda_i w_t(x_i) \in Q$ . By the triangle inequality and linearity of convex combinations,

$$\|a - b\| = \left\| \sum_{i=0}^k \lambda_i q_h(x_i) - \sum_{i=0}^k \lambda_i w_t(x_i) \right\| = \left\| \sum_{i=0}^k \lambda_i (q_h(x_i) - w_t(x_i)) \right\|.$$

By subadditivity of the norm,

$$\|a - b\| \leq \sum_{i=0}^k \lambda_i \cdot \|q_h(x_i) - w_t(x_i)\|.$$

For  $|h| < \delta$ ,  $\|q_h(x_i) - w_t(x_i)\| < \varepsilon$  for all  $i$ , and  $\sum \lambda_i = 1$ , so

$$\|a - b\| < \sum_{i=0}^k \lambda_i \cdot \varepsilon = \varepsilon.$$

Thus for every  $a \in Q_h$ , there exists  $b \in Q$  with  $\|a - b\| < \varepsilon$ , so  $\inf_{b \in Q} \|a - b\| < \varepsilon$ . Taking the supremum yields

$$\sup_{a \in Q_h} \inf_{b \in Q} \|a - b\| \leq \varepsilon. \quad (3.1)$$

(b) Similarly, for every  $b \in Q$ , there exists  $a \in Q_h$  with  $\|b - a\| < \varepsilon$ , so  $\inf_{a \in Q_h} \|b - a\| < \varepsilon$ . Taking the supremum yields

$$\sup_{b \in Q} \inf_{a \in Q_h} \|b - a\| \leq \varepsilon. \quad (3.2)$$

Combining (3.1), (3.2) and the definition of Hausdorff distance:

$$d_H(Q_h, Q) = \max \left\{ \sup_{a \in Q_h} \inf_{b \in Q} \|a - b\|, \sup_{b \in Q} \inf_{a \in Q_h} \|b - a\| \right\} < \varepsilon.$$

Since  $\varepsilon > 0$  is arbitrary, as  $h \rightarrow 0$ ,

$$d_H(Q_h, Q) \rightarrow 0.$$

That is, the difference quotient set  $Q_h$  converges to the target set  $Q = |f_U(\sigma)|$  in the Hausdorff distance.

In summary,  $f_U(\sigma) = \lim_{h \rightarrow 0} \frac{f_U^h(\sigma) - \sigma}{h}$  exists, with geometric realization  $|f_U(\sigma)|$ .

One also verifies that  $f_U$  is a simplicial homomorphism, i.e., it preserves simplex structure and inclusions. This verification is straightforward. Hence  $f_U \in \text{Hom}(K_t, K_U)$ .

It remains to check compatibility of the homomorphism family. For any  $U_2 \subset U_1$ , define the restriction morphism  $\rho_{U_2, U_1} : \text{Hom}(K_t, K_{U_1}) \rightarrow \text{Hom}(K_t, K_{U_2})$  by

$$\rho_{U_2, U_1}(f_{U_1})(\sigma) = f_{U_1}(\sigma) \cap K_{U_2}.$$

Since  $U_2 \subset U_1$ , the simplices of  $K_{U_2}$  are a subset of those of  $K_{U_1}$ , and  $|f_{U_2}(\sigma)| = |f_{U_1}(\sigma)| \cap |K_{U_2}|$ , so  $f_{U_2}(\sigma) = \rho_{U_2, U_1}(f_{U_1})(\sigma)$ , i.e.,  $f_{U_2} = \rho_{U_2, U_1}(f_{U_1})$ .

Thus  $(f_U)_{U \in \mathcal{U}_{K_t}} \in \prod_{U \in \mathcal{U}_{K_t}} \text{Hom}(K_t, K_U)$  is a compatible family, as required.

We now confirm differentiability. By the equivalence relation  $\sim$ ,  $[(f_U^h)] = [(f_U^0) + h \cdot (f_U) + (o(h))]$ , where  $(o(h))$  denotes a family of homomorphisms each component of which is  $o(h)$ . By definition of convergence in the quotient topology,

$$\lim_{h \rightarrow 0} \frac{[(f_U^h)] - [(f_U^0)]}{h} = [(f_U)] \in T_{K_t} \mathcal{S}(M).$$

Hence  $\mathcal{C}_{\Phi, \varepsilon}$  is differentiable at  $t$ , and we set  $\frac{d}{dt} \mathcal{C}_{\Phi, \varepsilon}(t) = [(f_U)]$ .

Finally, we establish membership and uniqueness of the tangent vector. By construction,  $[(f_U)]$  is an equivalence class in the inverse limit, so  $\frac{d}{dt} \mathcal{C}_{\Phi, \varepsilon}(t) \in T_{K_t} \mathcal{S}(M)$ .

For uniqueness, suppose  $\dot{K}_t^{(1)} = [(f_U^{(1)})]$  and  $\dot{K}_t^{(2)} = [(f_U^{(2)})]$  both satisfy the definition of differentiability. Then

$$\lim_{h \rightarrow 0} \frac{[(f_U^h)] - [(f_U^0)]}{h} = [(f_U^{(1)})] = [(f_U^{(2)})].$$

By  $\sim$ , there exists  $V \in \mathcal{U}_{K_t}$  such that for all  $U \subseteq V$ ,  $f_U^{(1)} = f_U^{(2)}$ . Since both  $f_U^{(1)}$  and  $f_U^{(2)}$  are induced by  $w_t(x) = d\Phi_t(x)(v_t(x))$ , and  $v_t(x) = \frac{\partial \Phi}{\partial t}(x, t)$  is unique,  $w_t(x)$  is unique, hence  $f_U^h$  is unique, so  $f_U^{(1)} = f_U^{(2)}$  for all  $U \in \mathcal{U}_{K_t}$ , i.e.,  $[(f_U^{(1)})] = [(f_U^{(2)})]$ . Thus the tangent vector is uniquely determined by  $v_t$ .

We now examine the projection operator  $\mathcal{P}$ . For any input  $W = \bigcup_{x \in M} w_t(x) \in \mathcal{D}(\mathcal{P})$ , define  $\mathcal{P}(W) = [(f_U^W)]$ , where  $(f_U^W)$  is the compatible family induced by  $W$ , such that for any  $U \in \mathcal{U}_{K_t}$ ,

$$f_U^W(\sigma) = \lim_{h \rightarrow 0} \frac{f_U^{h, W}(\sigma) - \sigma}{h},$$

where  $f_U^{h, W}$  is the “approximate homomorphism” generated using velocities  $w_t(x)$  from  $W$  (similarly to  $f_U^h$ , only replacing the velocity source with  $W$ ), and  $[(f_U^W)]$  is the equivalence class of  $(f_U^W)$  in the tangent space.

When the input is  $W = \bigcup_{x \in M} d\Phi_t(x)(v_t(x))$ , the induced family  $(f_U^W)$  coincides exactly with the family  $(f_U)$  constructed in the differentiability proof, i.e.,  $f_U^W = f_U$ . Since  $(f_U)$  is compatible, its equivalence class  $[(f_U)] \in T_{K_t} \mathcal{S}(M)$ , so  $\mathcal{P}(W) = [(f_U)]$  is well-defined.

We have shown that the tangent vector of the complex flow is  $\frac{d}{dt} \mathcal{C}_{\Phi, \varepsilon}(t) = [(f_U)]$ , and the projection gives  $\mathcal{P}(W) = [(f_U)]$ . Therefore,

$$\frac{d}{dt} \mathcal{C}_{\Phi, \varepsilon}(t) = \mathcal{P} \left( \bigcup_{x \in M} d\Phi_t(x)(v_t(x)) \right).$$

In conclusion, the dynamic Čech continuous flow  $\mathcal{C}_{\Phi, \varepsilon}$  is differentiable at  $t$ , its tangent vector lies in  $T_{K_t} \mathcal{S}(M)$ , is uniquely determined by the embedding velocity field  $v_t$ , and satisfies the projection formula.  $\square$

In this subsection, we have developed a rigorous mathematical framework for dynamic Čech complexes. Starting from the concept of continuous complex flows, via functorial constructions and tangent space theory, we have provided new tools for understanding the continuous evolution of topological structures. In subsequent sections, we build sheaf-theoretic frameworks and cohomology theories upon these foundations.

### 3.2 CONSTRUCTION OF THE DYNAMIC ČECH SHEAF

In dynamic Topological Data Analysis (TDA), the traditional sheaf-theoretic framework faces fundamental challenges: how to describe the local-global relationships evolving continuously with parameters? This subsection will construct the theory of dynamic Čech sheaves, which provides a new mathematical language for understanding the continuous changes of topological features by endogenously encoding parametric evolution into the sheaf structure.

The core idea of constructing the dynamic Čech presheaf is to incorporate parametric dependence into the classical Čech construction, such that local sections can reflect the continuous evolution of topological structures.

*Definition 3.13* (Dynamic Čech Presheaf). Let  $(M, g)$  be a  $d$ -dimensional compact Riemannian manifold, the parameter space  $T$  be a connected topological space, and  $\Phi : M \times T \rightarrow \mathbb{R}^D$  be a dynamic embedding family, given a radius  $\varepsilon > 0$ . The **dynamic Čech presheaf**  $\mathcal{P}_{\Phi, \varepsilon}$  is defined as follows:

For any open set  $U \subseteq M \times T$ ,

$$\mathcal{P}_{\Phi, \varepsilon}(U) := \left\{ s : U \rightarrow \mathbb{R} \left| \begin{array}{l} \forall (x_0, t_0) \in U, \exists \text{ open neighborhood } V \times W \subseteq U, \\ \text{such that } s|_{V \times W} \in C^\infty(V \times W), \text{ and } \forall t \in W, \\ s(\cdot, t) \text{ is locally constant on each simplex of } \mathcal{C}_{\Phi, \varepsilon}(t) \end{array} \right. \right\}$$

The restriction morphisms are the natural restrictions of functions.

This definition requires sections to maintain the local constancy determined by the Čech complex in the spatial direction and smoothness in the parametric direction, thereby capturing the continuous essence of topological evolution.

Dynamic Čech presheaves usually do not satisfy the gluing axiom of sheaves, so we need to obtain a genuine sheaf structure through sheafification.

*Definition 3.14* (Dynamic Čech Sheaf). The sheafification  $\mathcal{D}_{\Phi, \varepsilon} := (\mathcal{P}_{\Phi, \varepsilon})^\#$  of the dynamic Čech presheaf  $\mathcal{P}_{\Phi, \varepsilon}$  forms a sheaf on  $M \times T$ , called the **dynamic Čech sheaf**. Specifically,

$$\mathcal{D}_{\Phi, \varepsilon}(U) = \left\{ s : U \rightarrow \coprod_{(x, t) \in U} (\mathcal{P}_{\Phi, \varepsilon})_{(x, t)} \left| \begin{array}{l} \forall (x, t) \in U, \exists \text{ open neighborhood } V \ni (x, t) \text{ and a section} \\ f \in \mathcal{P}_{\Phi, \varepsilon}(V), \text{ such that for all } \xi \in V, s(\xi) = f_\xi. \end{array} \right. \right\}$$

The universal property of the dynamic Čech sheaf establishes its central position in the category of sheaves satisfying the dynamic local constancy condition.

**Theorem 3.15** (Universal Property). *Let  $\mathcal{F}$  be a sheaf on  $M \times T$ . If for any open set  $U \subseteq M \times T$  and any section  $s \in \mathcal{F}(U)$ , there exists an open cover  $\{U_i\}$  and sections  $f_i \in \mathcal{P}_{\Phi, \varepsilon}(U_i)$  such that  $s|_{U_i} = f_i$  in the sheaf  $\mathcal{F}$ , then there exists a unique sheaf morphism  $\varphi : \mathcal{F} \rightarrow \mathcal{D}_{\Phi, \varepsilon}$ .*

*Proof.* We prove this by constructing the sheaf morphism  $\varphi : \mathcal{F} \rightarrow \mathcal{D}_{\Phi, \varepsilon}$ .

For any open set  $U \subseteq M \times T$  and  $s \in \mathcal{F}(U)$ , by assumption, there exists an open cover  $\{U_i\}$  and  $f_i \in \mathcal{P}_{\Phi, \varepsilon}(U_i)$  such that  $s|_{U_i} = f_i$  in  $\mathcal{F}$ . Consider the sheafification morphism  $\theta : \mathcal{P}_{\Phi, \varepsilon} \rightarrow \mathcal{D}_{\Phi, \varepsilon}$ , and define  $\tilde{f}_i = \theta_{U_i}(f_i) \in \mathcal{D}_{\Phi, \varepsilon}(U_i)$ . By the definition of sheafification,  $\tilde{f}_i$  is the germ section induced by  $f_i$ , i.e., for any  $(x, t) \in U_i$ ,  $\tilde{f}_i(x, t) = (f_i)_{(x, t)}$ .



On the intersection  $U_i \cap U_j$ , we have  $s|_{U_i \cap U_j} = f_i|_{U_i \cap U_j} = f_j|_{U_i \cap U_j}$  in  $\mathcal{F}$ . Since  $\mathcal{F}$  is a sheaf,  $f_i$  and  $f_j$  define the same section on  $U_i \cap U_j$ .

By the definition of the dynamic Čech presheaf,  $f_i|_{U_i \cap U_j}$  and  $f_j|_{U_i \cap U_j}$  are equal in  $\mathcal{P}_{\Phi, \varepsilon}(U_i \cap U_j)$ , so they induce the same germ section  $\theta_{U_i \cap U_j}(f_i|_{U_i \cap U_j}) = \theta_{U_i \cap U_j}(f_j|_{U_i \cap U_j})$ . Thus  $\tilde{f}_i|_{U_i \cap U_j} = \tilde{f}_j|_{U_i \cap U_j}$ . By the sheaf axioms of  $\mathcal{D}_{\Phi, \varepsilon}$ , there exists a unique  $\tilde{s} \in \mathcal{D}_{\Phi, \varepsilon}(U)$  such that  $\tilde{s}|_{U_i} = \tilde{f}_i$ .

Define  $\varphi_U : \mathcal{F}(U) \rightarrow \mathcal{D}(U)$ ,  $s \mapsto \tilde{s}$ .

It is clear that the  $\varphi$  defined in this way is indeed a sheaf morphism. Next, we verify the uniqueness.

Let  $\psi : \mathcal{F} \rightarrow \mathcal{D}_{\Phi, \varepsilon}$  be another sheaf morphism satisfying the conditions. For any  $s \in \mathcal{F}(U)$ , consider the open cover  $\{U_i\}$  and  $f_i \in \mathcal{P}_{\Phi, \varepsilon}(U_i)$  such that  $s|_{U_i} = f_i$ . On  $U_i$ ,  $\varphi(s|_{U_i}) = \theta_{U_i}(f_i) = \psi(s|_{U_i})$ . By locality,  $\varphi(s) = \psi(s)$ .  $\square$

**Corollary 3.16.** *If there exists another sheaf  $\mathcal{G}$  satisfying the same universal property, then  $\mathcal{G} \cong \mathcal{D}_{\Phi, \varepsilon}$ .*

*Proof.* By the universal property of  $\mathcal{G}$ , there exists a unique sheaf morphism  $\varphi : \mathcal{D}_{\Phi, \varepsilon} \rightarrow \mathcal{G}$ . By the universal property of  $\mathcal{D}_{\Phi, \varepsilon}$ , there exists a unique sheaf morphism  $\psi : \mathcal{G} \rightarrow \mathcal{D}_{\Phi, \varepsilon}$ .

For any  $s \in \mathcal{D}_{\Phi, \varepsilon}(U)$ , by the local dynamic local constancy condition, there exists an open cover  $\{U_i\}$  and  $f_i \in \mathcal{P}_{\Phi, \varepsilon}(U_i)$  such that  $s|_{U_i} = \theta_{U_i}(f_i)$ . Then  $(\psi \circ \varphi)(s|_{U_i}) = \psi(\varphi(\theta_{U_i}(f_i))) = \psi(\theta_{U_i}(f_i)) = \theta_{U_i}(f_i) = s|_{U_i}$ , so  $\psi \circ \varphi = \text{id}$ .

Similarly,  $\varphi \circ \psi = \text{id}_{\mathcal{G}}$ . Therefore  $\mathcal{G} \cong \mathcal{D}_{\Phi, \varepsilon}$ .  $\square$

The following concept of dynamic star neighborhood generalizes the classical star neighborhood to the parametric case, providing a geometric foundation for understanding local consistency.

**Definition 3.17** (Dynamic Star Neighborhood). The **dynamic star neighborhood** of a point  $(x, t) \in M \times T$  is defined as:

$$\text{St}_{\Phi, \varepsilon}(x, t) := \{(y, \tau) \in M \times T \mid \{x, y\} \in \mathcal{C}_{\Phi, \varepsilon}(\tau)\}$$

i.e., all pairs of points that are covered by the same  $\varepsilon$ -ball as  $x$  at parameter  $\tau$ .

**Proposition 3.18** (Local Consistency Condition). *On the dynamic star neighborhood  $\text{St}_{\Phi, \varepsilon}(x, t)$ , the sections of the dynamic Čech sheaf satisfy local consistency:*

$$s(y, \tau) = s(x, t) + O(|\tau - t| + d_M(x, y))$$

i.e., the variation of section values is jointly controlled by parametric variation and spatial distance.

*Proof.* Let  $s \in \mathcal{D}_{\Phi, \varepsilon}(U)$  be a section of the dynamic Čech sheaf on an open set  $U \subseteq M \times T$ . According to the definition of the dynamic Čech sheaf, for any  $(x, t) \in U$ , there exists an open neighborhood  $V \ni (x, t)$  and a presheaf section  $f \in \mathcal{P}_{\Phi, \varepsilon}(V)$  such that for any  $\xi \in V$ ,  $s(\xi) = f_\xi$ , where  $f_\xi$  is the germ of  $f$  at  $\xi$ .

By the definition of the dynamic Čech presheaf,  $f$  satisfies: for any  $(y, \tau) \in V$ ,  $f(y, \tau)$  is locally constant on each simplex of  $\mathcal{C}_{\Phi, \varepsilon}(\tau)$ , and  $f$  is smooth on  $V$ .

Consider  $(y, \tau) \in \text{St}_{\Phi, \varepsilon}(x, t) \cap V$ . By the definition of the dynamic star neighborhood,  $\{x, y\} \in \mathcal{C}_{\Phi, \varepsilon}(\tau)$ , i.e., there exists  $z \in \mathbb{R}^D$  such that:

$$z \in B(\Phi_\tau(x), \varepsilon) \cap B(\Phi_\tau(y), \varepsilon)$$

Since  $f(\cdot, \tau)$  is locally constant on each simplex of  $\mathcal{C}_{\Phi, \varepsilon}(\tau)$ , and  $\{x, y\}$  is a 1-simplex in  $\mathcal{C}_{\Phi, \varepsilon}(\tau)$ , we have:

$$f(x, \tau) = f(y, \tau)$$

Since  $f \in C^\infty(V)$ , at the point  $(x, t)$ , for  $(y, \tau) \in \text{St}_{\Phi, \varepsilon}(x, t) \cap V$ , we have:

$$f(y, \tau) = f(x, t) + \frac{\partial f}{\partial t}(x, t)(\tau - t) + \nabla_x f(x, t) \cdot (y - x) + R(y, \tau)$$

where the remainder  $R(y, \tau)$  satisfies  $|R(y, \tau)| \leq C(|\tau - t|^2 + d_M(x, y)^2)$ . In particular:

$$f(y, \tau) = f(x, \tau) = f(x, t) + \frac{\partial f}{\partial t}(x, t)(\tau - t) + R(x, \tau)$$

Since  $f$  is locally constant in the spatial direction,  $\nabla_x f(x, t) = 0$ , so:

$$f(y, \tau) = f(x, t) + \frac{\partial f}{\partial t}(x, t)(\tau - t) + O(|\tau - t|^2)$$

Since  $s(\xi) = f_\xi$  for all  $\xi \in V$ , in particular:

$$s(y, \tau) = f(y, \tau), \quad s(x, t) = f(x, t)$$

Therefore,

$$s(y, \tau) = s(x, t) + \frac{\partial f}{\partial t}(x, t)(\tau - t) + O(|\tau - t|^2)$$

Since  $\frac{\partial f}{\partial t}(x, t)$  is bounded (as  $f$  is smooth and  $M \times T$  is compact), we obtain:

$$s(y, \tau) = s(x, t) + O(|\tau - t|)$$

Considering that on the dynamic star neighborhood,  $d_M(x, y)$  may not be zero, but by the Lipschitz property of the embedding and the compactness of  $M$ ,

$$d_M(x, y) \leq L^{-1} d_{\mathbb{R}^D}(\Phi_\tau(x), \Phi_\tau(y)) < 2\varepsilon L^{-1} \implies O(|\tau - t|) = O(|\tau - t| + d_M(x, y)).$$

we finally get  $s(y, \tau) = s(x, t) + O(|\tau - t| + d_M(x, y))$ .  $\square$

A direct corollary of the local consistency condition is the continuity of dynamic Čech sheaf sections:

**Corollary 3.19** (Continuity). *The sections of the dynamic Čech sheaf are continuous on the dynamic star neighborhood. In particular, if  $(y_n, \tau_n) \rightarrow (x, t)$  in  $\text{St}_{\Phi, \varepsilon}(x, t)$ , then  $s(y_n, \tau_n) \rightarrow s(x, t)$ .*

*Proof.* By the local consistency condition, when  $(y_n, \tau_n) \rightarrow (x, t)$ , we have  $|\tau_n - t| \rightarrow 0$  and  $d_M(y_n, x) \rightarrow 0$ , so:

$$|s(y_n, \tau_n) - s(x, t)| \leq C(|\tau_n - t| + d_M(y_n, x)) \rightarrow 0$$

i.e.,  $s(y_n, \tau_n) \rightarrow s(x, t)$ .  $\square$

The Theorem of Limit Representation of Stalks reveals the profound algebraic properties of the local structure of dynamic Čech sheaves, establishing a bridge between dynamic sheaf theory and classical smooth function theory.

**Theorem 3.20** (Limit Representation of Stalks). *The stalk of the dynamic Čech sheaf at  $(x, t)$  is isomorphic to:*

$$(\mathcal{D}_{\Phi, \varepsilon})_{(x, t)} \cong \varinjlim_{U \ni (x, t)} \left( C^\infty(U) / \sim_{\Phi, \varepsilon} \right)$$

where the equivalence relation  $\sim_{\Phi, \varepsilon}$  is defined as:  $f \sim g$  if and only if there exists a neighborhood  $V$  of  $(x, t)$  such that for any  $(\xi, \tau) \in V$ ,

$$f(\xi, \tau) = g(\xi, \tau) \quad \text{on } \text{St}_{\Phi, \varepsilon}(\xi, \tau)$$

*Proof.* First, consider the left-hand side:

$$(\mathcal{D}_{\Phi, \varepsilon})_{(x, t)} = \varinjlim_{U \ni (x, t)} \mathcal{D}_{\Phi, \varepsilon}(U)$$

Restriction homomorphism: For  $U_2 \subseteq U_1$ , there exists  $\rho_{U_2, U_1} : \mathcal{D}_{\Phi, \varepsilon}(U_1) \rightarrow \mathcal{D}_{\Phi, \varepsilon}(U_2)$ ,  $s \mapsto s|_{U_2}$ .

Equivalence relation:  $s_1 \in \mathcal{D}_{\Phi, \varepsilon}(U_1) \sim s_2 \in \mathcal{D}_{\Phi, \varepsilon}(U_2)$  if and only if there exists  $V \ni (x, t)$ ,  $V \subseteq U_1 \cap U_2$ , such that  $s_1|_V = s_2|_V$ .

Next, consider the right-hand side:

$$\varinjlim_{U \ni (x, t)} C^\infty(U) / \sim_{\Phi, \varepsilon}$$

Restriction morphism: For  $U_2 \subseteq U_1$ , there exists  $\bar{\rho}_{U_2, U_1} : C^\infty(U_1) / \sim_{\Phi, \varepsilon} \rightarrow C^\infty(U_2) / \sim_{\Phi, \varepsilon}$ ,  $[f] \mapsto [f|_{U_2}]$ .

Equivalence relation:  $[f_1] \in C^\infty(U_1) / \sim_{\Phi, \varepsilon} \sim [f_2] \in C^\infty(U_2) / \sim_{\Phi, \varepsilon}$  if and only if there exists  $V \ni (x, t)$ ,  $V \subseteq U_1 \cap U_2$ , such that  $[f_1|_V] = [f_2|_V]$ .

For each open set  $U$  containing  $(x, t)$ , define the map:

$$\eta_U : \mathcal{D}_{\Phi, \varepsilon}(U) \rightarrow C^\infty(U) / \sim_{\Phi, \varepsilon}, \quad s \mapsto [f_s]$$

where  $f_s \in C^\infty(U)$  is the presheaf section inducing  $s$ , specifically:

By the definition of the dynamic Čech sheaf, for any  $s \in \mathcal{D}_{\Phi, \varepsilon}(U)$ , there exists an open cover  $\{U_i\}$  and  $f_i \in \mathcal{P}_{\Phi, \varepsilon}(U_i)$  such that  $s|_{U_i} = \theta_{U_i}(f_i)$ , where  $\theta : \mathcal{P}_{\Phi, \varepsilon} \rightarrow \mathcal{D}_{\Phi, \varepsilon}$  is the sheafification morphism, and  $f_i$  satisfies  $f_i|_{U_i \cap U_j} = f_j|_{U_i \cap U_j}$  on the overlapping regions  $U_i \cap U_j$ . These  $f_i$  can be glued to form a global smooth function  $f_s \in C^\infty(U)$ .

$\eta_U$  is well-defined:

1. If  $s_1 = s_2$ , then there exists a neighborhood  $V$  of  $(x, t)$  such that  $s_1|_V = s_2|_V$ , thus  $f_{s_1} \sim_{\Phi, \varepsilon} f_{s_2}$  on  $V$ , so  $[f_{s_1}] = [f_{s_2}]$ .
2. The map is independent of the choice of representation: if  $s$  is induced by different  $f'_i$ , then  $f_s \sim_{\Phi, \varepsilon} f'_s$ .

$\eta_U$  is bijective:

1. Injectivity: If  $\eta_U(s) = \eta_U(s')$ , then  $[f_s] = [f_{s'}]$ , there exists  $V \ni (x, t)$  such that  $f_s \sim_{\Phi, \varepsilon} f_{s'}$  on  $V$ , so  $s|_V = s'|_V$ , thus  $s_{(x, t)} = s'_{(x, t)}$ .
2. Surjectivity: For any  $[f] \in C^\infty(U) / \sim_{\Phi, \varepsilon}$ , since  $f \in \mathcal{P}_{\Phi, \varepsilon}(U)$ , we obtain  $s_f = \theta_U(f) \in \mathcal{D}_{\Phi, \varepsilon}(U)$  by sheafification, and  $\eta_U(s_f) = [f]$ .

We further claim that for any  $U_2 \subseteq U_1$ , the following diagram commutes:

$$\begin{array}{ccc} \mathcal{D}_{\Phi,\varepsilon}(U_1) & \xrightarrow{\eta_{U_1}} & C^\infty(U_1)/\sim_{\Phi,\varepsilon} \\ \rho_{U_2,U_1} \downarrow & & \downarrow \bar{\rho}_{U_2,U_1} \\ \mathcal{D}_{\Phi,\varepsilon}(U_2) & \xrightarrow{\eta_{U_2}} & C^\infty(U_2)/\sim_{\Phi,\varepsilon} \end{array}$$

Specifically, for any  $s \in \mathcal{D}_{\Phi,\varepsilon}(U_1)$ , let  $\eta_{U_1}(s) = [f_s]$ , then:

$$\begin{aligned} \eta_{U_2}(\rho_{U_2,U_1}(s)) &= \eta_{U_2}(s|_{U_2}) = [f_s|_{U_2}] \\ \bar{\rho}_{U_2,U_1}(\eta_{U_1}(s)) &= \bar{\rho}_{U_2,U_1}([f_s]) = [f_s|_{U_2}] \end{aligned}$$

Thus the diagram commutes.

We have therefore established the isomorphism:

$$\eta : \{\mathcal{D}_{\Phi,\varepsilon}(U), \rho_{U',U}\} \rightarrow \{C^\infty(U)/\sim_{\Phi,\varepsilon}, \bar{\rho}_{U',U}\}$$

By the universal property of direct limits, if two systems are naturally isomorphic, their direct limits are isomorphic. Therefore:

$$(\mathcal{D}_{\Phi,\varepsilon})_{(x,t)} \cong \varinjlim_{U \ni (x,t)} C^\infty(U)/\sim_{\Phi,\varepsilon}$$

□

As a direct application of this theorem, we can obtain the dimension formula for the stalks of dynamic sheaves:

**Corollary 3.21** (Dimension Formula for Stalks). *At a non-critical point  $(x, t)$ , the dimension of the stalk of the dynamic Čech sheaf is:*

$$\dim_{\mathbb{R}}(\mathcal{D}_{\Phi,\varepsilon})_{(x,t)} = \dim \text{St}_{\Phi,\varepsilon}(x, t) + 1$$

where  $\dim \text{St}_{\Phi,\varepsilon}(x, t)$  is the dimension of the dynamic star neighborhood.

*Proof.* Notice  $\dim \left( \varinjlim_{U \ni (x,t)} C^\infty(U)/\sim_{\Phi,\varepsilon} \right) = \dim \text{St}_{\Phi,\varepsilon}(x, t) + \dim T = \dim \text{St}_{\Phi,\varepsilon}(x, t) + 1$ . □

We note that the dynamic Čech sheaf is not only a sheaf, but more importantly, it is functorial, which reflects its natural compatibility with parametric evolution.

**Definition 3.22** (Dynamic Sheaf Functor). The construction of the dynamic Čech sheaf defines a functor from the parameter category to the sheaf category, called the **dynamic sheaf functor**:

$$\mathcal{D}_{\Phi,\varepsilon} : T \rightarrow \text{Sh}(M \times T)$$

Specifically:

1. For any  $t \in T$ ,  $\mathcal{D}_{\Phi,\varepsilon}(t) := \mathcal{D}_{\Phi,\varepsilon}|_{M \times \{t\}}$  is a sheaf on  $M$ .
2. For any path  $\gamma : s \rightarrow t$ , it induces a sheaf homomorphism  $\mathcal{D}_{\Phi,\varepsilon}(\gamma) : \mathcal{D}_{\Phi,\varepsilon}(t) \rightarrow \mathcal{D}_{\Phi,\varepsilon}(s)$ .

The dynamic Čech sheaf is well compatible with standard sheaf operations, which ensures the consistency of the theoretical framework.

**Proposition 3.23** (Compatibility with Pullback Operations). *Let  $f : T' \rightarrow T$  be a continuous map, then there exists a natural isomorphism:*

$$f^* \mathcal{D}_{\Phi, \varepsilon} \cong \mathcal{D}_{\Phi \circ (\text{id}_M \times f), \varepsilon}$$

where:

1.  $f^*$  is the pullback sheaf, i.e., for a sheaf  $\mathcal{F}$  on  $M \times T$ ,  $f^* \mathcal{F}$  is a sheaf on  $M \times T'$ .
2.  $\Phi \circ (\text{id}_M \times f) : M \times T' \rightarrow \mathbb{R}^D$  is the composition map, defined as  $(\Phi \circ (\text{id}_M \times f))(x, t') = \Phi(x, f(t'))$ .

*Proof.* We prove the global isomorphism by verifying the isomorphism of the two sheaves at each stalk.

Let  $F = \text{id}_M \times f : M \times T' \rightarrow M \times T$ , and  $\Phi' = \Phi \circ F$ .

For any  $(x, t') \in M \times T'$ , consider the stalk isomorphism:

$$(f^* \mathcal{D}_{\Phi, \varepsilon})_{(x, t')} \cong (\mathcal{D}_{\Phi, \varepsilon})_{F(x, t')} = (\mathcal{D}_{\Phi, \varepsilon})_{(x, f(t'))}$$

By the Limit Representation of Stalks ([Theorem 3.20](#)):

$$\begin{aligned} (\mathcal{D}_{\Phi, \varepsilon})_{(x, f(t'))} &\cong \varinjlim_{U \ni (x, f(t'))} C^\infty(U) / \sim_{\Phi, \varepsilon} \\ (\mathcal{D}_{\Phi', \varepsilon})_{(x, t')} &\cong \varinjlim_{U' \ni (x, t')} C^\infty(U') / \sim_{\Phi', \varepsilon} \end{aligned}$$

Consider the two direct systems:

- $\mathcal{A} = \{C^\infty(U) / \sim_{\Phi, \varepsilon}\}_{U \ni (x, f(t'))}$ , where  $U \subseteq M \times T$  is open.
- $\mathcal{B} = \{C^\infty(U') / \sim_{\Phi', \varepsilon}\}_{U' \ni (x, t')}$ , where  $U' \subseteq M \times T'$  is open.

Define the functor  $\varphi : \mathcal{A} \rightarrow \mathcal{B}$  as follows:

For any  $U \ni (x, f(t'))$ , let  $U' = F^{-1}(U)$ , and define:

$$\varphi_U : C^\infty(U) / \sim_{\Phi, \varepsilon} \rightarrow C^\infty(U') / \sim_{\Phi', \varepsilon}, \quad [g] \mapsto [g \circ F]$$

It is straightforward to claim that the  $\varphi_U$  defined above is a well-defined linear isomorphism.

For any inclusion relation  $U_1 \subseteq U_2$ , consider the diagram:

$$\begin{array}{ccc} C^\infty(U_2) / \sim_{\Phi, \varepsilon} & \xrightarrow{\varphi_{U_2}} & C^\infty(F^{-1}(U_2)) / \sim_{\Phi', \varepsilon} \\ \rho_{U_1, U_2} \downarrow & & \downarrow \rho_{F^{-1}(U_1), F^{-1}(U_2)} \\ C^\infty(U_1) / \sim_{\Phi, \varepsilon} & \xrightarrow{\varphi_{U_1}} & C^\infty(F^{-1}(U_1)) / \sim_{\Phi', \varepsilon} \end{array}$$

Commutativity follows directly from the definition of restriction morphisms.

Since  $\varphi$  is a natural equivalence of direct systems, by the universal property of direct limits, it induces an isomorphism of stalks:

$$(f^* \mathcal{D}_{\Phi, \varepsilon})_{(x, t')} \cong (\mathcal{D}_{\Phi, \varepsilon})_{(x, f(t'))} \cong (\mathcal{D}_{\Phi', \varepsilon})_{(x, t')}$$

Take a basis of local coordinate neighborhoods of  $M \times T'$ . On each coordinate neighborhood, the stalk isomorphisms are consistent on overlapping regions by stalk isomorphism, so they can be glued to form a global sheaf isomorphism.

Finally, we verify the naturality of the obtained isomorphism with respect to pullback operations. For any continuous map  $g : T'' \rightarrow T'$ , consider the composition  $f \circ g : T'' \rightarrow T$ , and we need to verify the commutativity of the diagram:

$$\begin{array}{ccc}
(f \circ g)^* \mathcal{D}_{\Phi, \varepsilon} & \xrightarrow{\cong} & g^* f^* \mathcal{D}_{\Phi, \varepsilon} \\
\cong \downarrow & & \downarrow \cong \\
\mathcal{D}_{\Phi \circ (\text{id}_M \times (f \circ g)), \varepsilon} & \xrightarrow{\cong} & g^* \mathcal{D}_{\Phi \circ (\text{id}_M \times f), \varepsilon}
\end{array}$$

Commutativity is obvious from the functoriality of inductive limits and the definition of dynamic sheaves.  $\square$

This result shows that the transformation of the parameter space is compatible with the construction of dynamic sheaves.

**Corollary 3.24.** *Let  $f : T' \rightarrow T$  be a homeomorphism, then the dynamic topological features are preserved under parametric transformation, i.e.:*

$$\mathcal{D}_{\Phi, \varepsilon} \cong f_* \mathcal{D}_{\Phi', \varepsilon}$$

where  $f_*$  is the pushforward sheaf.

Comparisons between different dynamic Čech sheaves are realized through dynamic sheaf morphisms defined as follows.

**Definition 3.25** (Dynamic Sheaf Morphism). Let  $\mathcal{D}_{\Phi, \varepsilon}$  and  $\mathcal{D}_{\Phi', \varepsilon'}$  be two dynamic Čech sheaves. A **dynamic sheaf morphism**  $f : \mathcal{D}_{\Phi, \varepsilon} \rightarrow \mathcal{D}_{\Phi', \varepsilon'}$  is a family of sheaf homomorphisms  $\{f_t : \mathcal{D}_{\Phi, \varepsilon}(t) \rightarrow \mathcal{D}_{\Phi', \varepsilon'}(t)\}_{t \in T}$  such that for any path  $\gamma : s \rightarrow t$ , the following diagram commutes:

$$\begin{array}{ccc}
\mathcal{D}_{\Phi, \varepsilon}(t) & \xrightarrow{f_t} & \mathcal{D}_{\Phi', \varepsilon'}(t) \\
\mathcal{D}_{\Phi, \varepsilon}(\gamma) \downarrow & & \downarrow \mathcal{D}_{\Phi', \varepsilon'}(\gamma) \\
\mathcal{D}_{\Phi, \varepsilon}(s) & \xrightarrow{f_s} & \mathcal{D}_{\Phi', \varepsilon'}(s)
\end{array}$$

Dynamic sheaf morphisms enable us to compare the topological evolution behaviors under different dynamic processes (such as different embedding families or different scale parameters).

Through the development of this subsection, we have established a complete theoretical framework for dynamic Čech sheaves, from presheaf construction and sheafification process to the geometric description of stalk structures and functorial properties, providing a powerful mathematical tool for understanding the local-global relationships of dynamic topological processes. This framework lays a solid foundation for the persistent dynamic cohomology theory we will discuss next.

## 4 PERSISTENT DYNAMIC COHOMOLOGY THEORY

In dynamic topological data analysis, the classical single-parameter persistent homology theory faces a fundamental challenge: how to simultaneously describe the co-evolution of topological features along both the scale parameter  $\varepsilon$  and the time parameter  $t$ ? This section establishes a rigorous mathematical framework for persistent dynamic cohomology, resolving the difficulties of the two-parameter setting.

Recent research has shown that multi-parameter persistence modules are generically indecomposable [BS25], which brings challenges to our theory while also offering new insight. Unlike the classical single-parameter case, two-parameter persistence modules require more refined algebraic structures. This discovery motivates us to develop new theoretical tools to handle the complex phenomena in dynamic topological evolution.

### 4.1 FUNDAMENTAL THEORY OF PERSISTENT DYNAMIC COHOMOLOGY

Classical persistent homology is based on filtered complexes with a single scale parameter, while the dynamic setting requires the simultaneous interaction of scale and time. Inspired by the study of persistent homology on multi-grid posets [Día25], we establish the following categorical framework:

*Definition 4.1* (Two-parameter Persistence Module). Let  $\mathcal{P} = \mathbb{R}_{\geq 0} \times T$  be the scale-time parameter space equipped with the partial order:

$$(\varepsilon_1, t_1) \preceq (\varepsilon_2, t_2) \iff \varepsilon_1 \leq \varepsilon_2, t_1 \leq t_2.$$

A **two-parameter persistence module** is a functor:

$$M : (\mathcal{P}, \preceq) \rightarrow \mathbf{Vect}_K$$

such that for each pair  $(\varepsilon, t)$ , it assigns a vector space  $M(\varepsilon, t)$ , and for any  $(\varepsilon_1, t_1) \preceq (\varepsilon_2, t_2)$ , it assigns a transition homomorphism  $\varphi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)}$ , satisfying the functorial axioms.

*Definition 4.2* (Persistent Dynamic Cohomology). Let a dynamic embedding family  $\Phi : M \times T \rightarrow \mathbb{R}^D$  satisfy: for any  $t_1 \leq t_2 \in T$  and any fixed scale  $\varepsilon > 0$ , the earlier dynamic Čech complex is a subcomplex of the later one, i.e.,  $\mathcal{C}_{\Phi, \varepsilon}(t_1) \hookrightarrow \mathcal{C}_{\Phi, \varepsilon}(t_2)$ .

Under this condition, its  **$p$ -th persistent dynamic cohomology** is defined as the two-parameter persistence module:

$$\begin{aligned} \text{PDHP}(\Phi)(\varepsilon, t) &= H^p(\mathcal{C}_{\Phi, \varepsilon}(t); K), \\ \text{PDHP}(\Phi) &:= \left( \text{PDHP}(\Phi)(\varepsilon, t), \psi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)} \right)_{(\varepsilon_1, t_1) \preceq (\varepsilon_2, t_2)} \end{aligned}$$

where the transition homomorphism  $\psi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)} : \text{PDHP}(\Phi)(\varepsilon_2, t_2) \rightarrow \text{PDHP}(\Phi)(\varepsilon_1, t_1)$  is the pullback cohomology homomorphism induced by the inclusion  $\mathcal{C}_{\Phi, \varepsilon_1}(t_1) \hookrightarrow \mathcal{C}_{\Phi, \varepsilon_2}(t_2)$ .

This inclusion condition guarantees the causal structure of the dynamic process: as time evolves, the topological structure does not collapse drastically but evolves smoothly. Technically, it ensures that the partial order  $\preceq$  on the parameter space is compatible with the inclusion of complexes, so that the transition homomorphisms are well-defined.

**Theorem 4.3** (Functoriality). *The  $p$ -th persistent dynamic cohomology construction defines a functor from the category of dynamic embeddings to the category of two-parameter persistence modules:*

$$\text{PDHP} : \text{DynEmb}(M, \mathbb{R}^D) \rightarrow \text{PersMod}_2(K).$$

This functor preserves homotopy relations of dynamic embeddings: if  $\Phi \simeq \Psi$ , then  $\mathbb{PDH}^p(\Phi) \cong \mathbb{PDH}^p(\Psi)$ .

*Proof.* Let  $f : \Phi \rightarrow \Psi$  be a morphism of dynamic embeddings. For each parameter  $(\varepsilon, t)$ ,  $f$  induces a simplicial map  $f_{\varepsilon, t} : \mathcal{C}_{\Phi, \varepsilon}(t) \rightarrow \mathcal{C}_{\Psi, \varepsilon}(t)$ , which further induces a cohomology pullback:

$$f_{\varepsilon, t}^* : H^p(\mathcal{C}_{\Psi, \varepsilon}(t); K) \rightarrow H^p(\mathcal{C}_{\Phi, \varepsilon}(t); K).$$

Define  $\mathbb{PDH}^p(f)$  to be the natural transformation whose component at  $(\varepsilon, t)$  is  $f_{\varepsilon, t}^*$ .

It remains to verify naturality: for any  $(\varepsilon_1, t_1) \preceq (\varepsilon_2, t_2)$ , the following diagram commutes:

$$\begin{array}{ccc} H^p(\mathcal{C}_{\Psi, \varepsilon_2}(t_2); K) & \xrightarrow{\psi_{\Psi}} & H^p(\mathcal{C}_{\Psi, \varepsilon_1}(t_1); K) \\ f_{\varepsilon_2, t_2}^* \downarrow & & \downarrow f_{\varepsilon_1, t_1}^* \\ H^p(\mathcal{C}_{\Phi, \varepsilon_2}(t_2); K) & \xrightarrow{\psi_{\Phi}} & H^p(\mathcal{C}_{\Phi, \varepsilon_1}(t_1); K) \end{array}$$

This is guaranteed by the naturality of cohomology pullbacks.

The preservation of homotopy relations follows from a fundamental theorem in algebraic topology: simplicial homotopies induce isomorphisms in cohomology. Concretely, if  $H : \Phi \simeq \Psi$  is a homotopy, then for each fixed  $(\varepsilon, t)$ ,  $H$  induces a chain homotopy, which yields a cohomology isomorphism. ([Hat02], p.201, Paragraph begin with ‘‘Homotopy Invariance ...’’.)  $\square$

Persistent dynamic cohomology not only captures topological features but also preserves important algebraic structures that reflect the essential properties of dynamic processes.

**Theorem 4.4** (Monotonicity). *Under the defining conditions of persistent dynamic cohomology, the  $p$ -th persistent dynamic cohomology module satisfies monotonicity: for any chain of parameters  $(\varepsilon_1, t_1) \preceq (\varepsilon_2, t_2) \preceq (\varepsilon_3, t_3)$ , the transition homomorphisms satisfy*

$$\psi_{(\varepsilon_1, t_1)}^{(\varepsilon_3, t_3)} = \psi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)} \circ \psi_{(\varepsilon_2, t_2)}^{(\varepsilon_3, t_3)},$$

*and all transition homomorphisms are injective.*

*Proof.* By the inclusion condition in the definition, for  $\varepsilon_1 \leq \varepsilon_2$  and  $t_1 \leq t_2$ , we have  $\mathcal{C}_{\Phi, \varepsilon_1}(t_1) \hookrightarrow \mathcal{C}_{\Phi, \varepsilon_2}(t_2)$ . Thus for  $(\varepsilon_1, t_1) \preceq (\varepsilon_2, t_2) \preceq (\varepsilon_3, t_3)$ ,

$$\mathcal{C}_{\Phi, \varepsilon_1}(t_1) \hookrightarrow \mathcal{C}_{\Phi, \varepsilon_2}(t_2) \hookrightarrow \mathcal{C}_{\Phi, \varepsilon_3}(t_3).$$

The transition homomorphisms are included by pullback homomorphisms

$$\psi_{(\varepsilon_k, t_k)}^{(\varepsilon_j, t_j)}[f]_j = [i_{kj}^{*, p}(f)]_k, \quad i_{kj} : \mathcal{C}_k \hookrightarrow \mathcal{C}_j, \quad \forall 1 \leq k < j \leq 3.$$

By contravariance of the cohomology functor, the pullback of a composition equals the composition of pullbacks in reverse order:  $i_{13}^* = i_{12}^* \circ i_{23}^*$ . Substituting gives  $\psi_{(\varepsilon_1, t_1)}^{(\varepsilon_3, t_3)} = \psi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)} \circ \psi_{(\varepsilon_2, t_2)}^{(\varepsilon_3, t_3)}$ .

Injectivity follows from the fact that the pullback cohomology homomorphism induced by an inclusion map is injective in the same dimension.  $\square$

**Theorem 4.5** (Sheaf Property). *Under the defining conditions of persistent dynamic cohomology, the  $p$ -th persistent dynamic cohomology module admits an intrinsic sheaf structure: for any open cover  $\{U_i\}$  of the parameter space  $\mathcal{P}$ , there is an exact sequence:*

$$0 \rightarrow \mathbb{PDH}^p(\Phi)(U) \rightarrow \prod_i \mathbb{PDH}^p(\Phi)(U_i) \rightarrow \prod_{i, j} \mathbb{PDH}^p(\Phi)(U_i \cap U_j),$$

where  $\mathbb{PDH}^p(\Phi)(U) = \varprojlim_{(\varepsilon, t) \in U} \mathbb{PDH}^p(\Phi)(\varepsilon, t)$ , and  $U \subseteq \mathcal{P}$  is an open subset of the parameter space.



*Proof.* For any inclusion of open sets  $V \subseteq U \subseteq \mathcal{P}$ , define the restriction map:

$$r_V^U : \mathbb{PDHP}(\Phi)(U) \rightarrow \mathbb{PDHP}(\Phi)(V).$$

If  $s \in \mathbb{PDHP}(\Phi)(U)$  is a global section over  $U$ , then  $s$  is a compatible family of cohomology classes:

$$s = \{s((\varepsilon, t)) \in \mathbb{PDHP}(\Phi)(\varepsilon, t) \mid (\varepsilon, t) \in U\},$$

such that for any  $(\varepsilon_1, t_1) \preceq (\varepsilon_2, t_2) \in U$ , we have  $\psi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)}(s((\varepsilon_2, t_2))) = s((\varepsilon_1, t_1))$ . The restriction  $r_V^U(s)$ , written  $s|_V$ , is defined by  $s|_V = \{s((\varepsilon, t)) \mid (\varepsilon, t) \in V\}$ , which clearly satisfies compatibility over  $V$ .

By the universal property of inverse limits, the restriction maps satisfy the presheaf axioms:

1. Identity:  $r_U^U(s) = s|_U = s$ , i.e.,  $r_U^U = \text{id}$ ;
2. Transitivity: for  $W \subseteq V \subseteq U$ ,  $r_W^U(s) = (s|_V)|_W = r_W^V \circ r_V^U(s)$ , i.e.,  $r_W^U = r_W^V \circ r_V^U$ .

Therefore,  $\mathbb{PDHP}(\Phi)$  forms a presheaf over  $\mathcal{P}$  via the above restriction maps.

Define the left-hand map  $\rho : \mathbb{PDHP}(\Phi)(U) \rightarrow \prod_i \mathbb{PDHP}(\Phi)(U_i)$ ,  $s \mapsto (s|_{U_i})_i$ , which restricts a global section  $s$  to each open set  $U_i$  in the cover, yielding a family of local sections  $(s|_{U_i})_i$ . We now show that  $\rho$  is injective.

Let  $s, t \in \mathbb{PDHP}(\Phi)(U)$  satisfy  $\rho(s) = \rho(t)$ , meaning  $s|_{U_i} = t|_{U_i}$  for all  $i$ . For any  $(\varepsilon, t) \in U$ , since  $\{U_i\}$  covers  $U$ , there exists some  $U_i$  such that  $(\varepsilon, t) \in U_i$ , so:

$$s((\varepsilon, t)) = (s|_{U_i})((\varepsilon, t)) = (t|_{U_i})((\varepsilon, t)) = t((\varepsilon, t)).$$

Since  $(\varepsilon, t)$  is arbitrary,  $s = t$ , so  $\rho$  is injective and the sequence is exact at  $\mathbb{PDHP}(\Phi)(U)$ .

Define the right-hand map  $\delta : \prod_i \mathbb{PDHP}(\Phi)(U_i) \rightarrow \prod_{i,j} \mathbb{PDHP}(\Phi)(U_i \cap U_j)$  by:

$$\delta((s_i)_i) = ((s_i|_{U_i \cap U_j}) - (s_j|_{U_i \cap U_j}))_{i,j},$$

where  $-$  denotes pointwise subtraction.

We prove  $\text{im}(\rho) = \ker(\delta)$ , i.e., both inclusions  $\text{im}(\rho) \subseteq \ker(\delta)$  and  $\ker(\delta) \subseteq \text{im}(\rho)$ .

1.  $\text{im}(\rho) \subseteq \ker(\delta)$ .

For any  $s \in \mathbb{PDHP}(\Phi)(U)$ ,  $\rho(s) = (s|_{U_i})_i$ . By transitivity of restrictions,  $s|_{U_i}|_{U_i \cap U_j} = s|_{U_i \cap U_j}$  and similarly  $s|_{U_j}|_{U_i \cap U_j} = s|_{U_i \cap U_j}$ . Thus for any  $(\varepsilon, t) \in U_i \cap U_j$ :

$$(s|_{U_i}|_{U_i \cap U_j})((\varepsilon, t)) - (s|_{U_j}|_{U_i \cap U_j})((\varepsilon, t)) = s((\varepsilon, t)) - s((\varepsilon, t)) = 0,$$

so  $\delta(\rho(s)) = 0$  and hence  $\text{im}(\rho) \subseteq \ker(\delta)$ .

2.  $\ker(\delta) \subseteq \text{im}(\rho)$ .

Let  $(s_i)_i \in \ker(\delta)$ . Then for all  $i, j$  and  $(\varepsilon, t) \in U_i \cap U_j$ ,

$$(s_i|_{U_i \cap U_j})((\varepsilon, t)) = (s_j|_{U_i \cap U_j})((\varepsilon, t)).$$

Construct a global section  $s \in \mathbb{PDHP}(\Phi)(U)$ : for any  $(\varepsilon, t) \in U$ , choose an open set  $U_i$  containing  $(\varepsilon, t)$  and set  $s((\varepsilon, t)) = s_i((\varepsilon, t))$ .

Two things must be checked:

(a) Well-definedness.

If  $(\varepsilon, t) \in U_i \cap U_j$ , then by the definition of  $\ker(\delta)$ ,  $(s_i|_{U_i \cap U_j})(\varepsilon, t) = (s_j|_{U_i \cap U_j})(\varepsilon, t)$ . Since  $s_i((\varepsilon, t)) = (s_i|_{U_i \cap U_j})(\varepsilon, t)$  and  $s_j((\varepsilon, t)) = (s_j|_{U_i \cap U_j})(\varepsilon, t)$ , we have  $s_i((\varepsilon, t)) = s_j((\varepsilon, t))$ , independent of the choice of  $U_i$ .

(b) Compatibility.

For any  $(\varepsilon_1, t_1) \preceq (\varepsilon_2, t_2) \in U$ , by the monotonicity condition for persistent dynamic cohomology ( $\mathcal{C}_{\Phi, \varepsilon_1}(t_1) \hookrightarrow \mathcal{C}_{\Phi, \varepsilon_2}(t_2)$ ), the transition map  $\psi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)}$  is well-defined. Choose  $U_k$  containing both  $(\varepsilon_1, t_1)$  and  $(\varepsilon_2, t_2)$  (such  $U_k$  exists because  $U$  is open and the cover is exhaustive). Since  $s_k$  is a section over  $U_k$  (hence compatible), we have:

$$\psi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)}(s((\varepsilon_2, t_2))) = \psi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)}(s_k((\varepsilon_2, t_2))) = s_k((\varepsilon_1, t_1)) = s((\varepsilon_1, t_1)).$$

Therefore,  $s$  is a well-defined global section over  $U$ , and for any  $i$ ,  $s|_{U_i} = s_i$ , meaning  $(s_i)_i = \rho(s) \in \text{im}(\rho)$ . Thus  $\ker(\delta) \subseteq \text{im}(\rho)$ .

Finally, by construction of inverse limits,  $\mathbb{PDH}^p(\Phi)(U)$  satisfies:

1. For every open set  $U$ ,  $\mathbb{PDH}^p(\Phi)(U)$  is a  $K$ -vector space;
2. The transition homomorphisms  $\psi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)}$  are linear and satisfy transitivity;
3. The restriction maps  $r_V^U$  are compatible with transition homomorphisms, ensuring globality after gluing local sections.

In summary, the sequence

$$0 \rightarrow \mathbb{PDH}^p(\Phi)(U) \xrightarrow{\rho} \prod_i \mathbb{PDH}^p(\Phi)(U_i) \xrightarrow{\delta} \prod_{i,j} \mathbb{PDH}^p(\Phi)(U_i \cap U_j)$$

is exact, so  $\mathbb{PDH}^p(\Phi)$  forms a sheaf over  $\mathcal{P}$ . □

The sheaf property reflects the local-global principle in dynamic topological analysis and provides a theoretical foundation for distributed computation of dynamic topological features.

## 4.2 CONVERGENCE, STABILITY, AND DYNAMIC TOPOLOGICAL FEATURE MEASURES

The core of classical persistent homology is barcode theory, but in the two-parameter setting we face a fundamental challenge: recent work by Bauer and Scoccola [BS25] shows that multi-parameter persistence modules are generically indecomposable, meaning the traditional barcode decomposition no longer applies.

**Theorem 4.6** (Indecomposability). *Let  $\Phi$  be a generic dynamic embedding family. Then its  $p$ -th persistent dynamic cohomology module  $\mathbb{PDH}^p(\Phi)$  is **indecomposable**, i.e., it admits no non-trivial direct sum decomposition into interval modules.*

*Proof.* The proof relies on the generic indecomposability of multi-parameter persistence modules [BS25]. We verify that persistent dynamic cohomology modules satisfy the conditions of that theory:

1.  $\mathbb{PDH}^p(\Phi)$  is a finitely presented multi-parameter persistence module.

Since  $M$  is a compact manifold and the dynamic embedding  $\Phi$  is continuous, for any fixed  $(\varepsilon, t) \in \mathcal{P}$ , the Čech complex  $\mathcal{C}_{\Phi, \varepsilon}(t)$  generated by an  $\varepsilon$ -neighborhood cover is a finite

simplicial complex, so its  $p$ -th cohomology group  $H^p(\mathcal{C}_{\Phi, \varepsilon}(t); K)$  is a finite-dimensional  $K$ -vector space. Furthermore, by monotonicity of the dynamic embedding  $(\mathcal{C}_{\Phi, \varepsilon_1}(t_1) \hookrightarrow \mathcal{C}_{\Phi, \varepsilon_2}(t_2))$  whenever  $\varepsilon_1 \leq \varepsilon_2, t_1 \leq t_2$ , there exists a finite scale-time grid  $\mathcal{P} \subseteq \mathcal{P}$  such that  $\mathbb{PDH}^p(\Phi)$  is isomorphic to the left Kan extension of its restriction to  $\mathcal{P}$  (i.e.,  $\mathbb{PDH}^p(\Phi) \cong \text{Lan}(\mathbb{PDH}^p(\Phi)|_{\mathcal{P}})$ ). By [BS25] Lemma 2.1 (finitely presented modules are equivalent to finite-grid restriction–extension modules),  $\mathbb{PDH}^p(\Phi)$  is finitely presented. It can therefore be expressed as the cokernel of a morphism  $\bigoplus_j P_j \rightarrow \bigoplus_i P_i$  of basic modules  $P_i$ , where  $\mathcal{P} = \mathbb{R}_{\geq 0} \times T$  is a two-parameter space ( $n = 2$ ).

2. For generic  $\Phi$ , the module  $\mathbb{PDH}^p(\Phi)$  is generic.

A “generic dynamic embedding family” is defined as one in the space of  $C^1$  dynamic embeddings that does not belong to any lower-dimensional algebraic subset (i.e., the generic case with no instantaneous collapse of topological features and no redundant parameter dependence). By [BS25] Theorem A (indecomposable modules are dense in the space of finitely presented modules) and Theorem B ( $\varepsilon$ -indecomposable modules form an open set), the module induced by such an embedding belongs to a dense open set of indecomposable modules, hence is generic.

3. No non-trivial direct sum decomposition into interval modules.

Suppose  $\mathbb{PDH}^p(\Phi) \cong A \oplus B$  with  $A, B$  non-zero interval modules. Then  $A, B$  are finitely presented interval modules. However, [BS25] Corollary 1.4 implies that decomposable modules relying on finite-dimensional indecomposable interval modules are nowhere dense, contradicting that  $\mathbb{PDH}^p(\Phi)$  is generic (lies in a dense open set). Therefore  $\mathbb{PDH}^p(\Phi)$  admits no non-trivial direct sum decomposition into interval modules, i.e., it is indecomposable.

□

The profound implication of this result is that we can no longer rely on traditional barcodes to represent multi-parameter topological features and must develop new descriptive methods.

Inspired by [WLY25], we give the following definition, which uses measure-theoretic tools to describe the distribution of topological features and bypasses the indecomposability problem.

*Definition 4.7* (Dynamic  $p$ -dimensional Topological Feature Measure). Under the defining conditions of persistent dynamic cohomology, the **dynamic topological feature measure of  $p$ -th persistent dynamic cohomology** is defined as:

$$\mu_{\Phi}^p : \mathcal{B}(\mathcal{P}) \rightarrow \mathbb{R}_{\geq 0},$$

where  $\mathcal{B}(\mathcal{P})$  is the Borel  $\sigma$ -algebra of the parameter space  $\mathcal{P} = \mathbb{R}_{\geq 0} \times T$ , and for any measurable set  $A \subseteq \mathcal{P}$ , the measure is given by the normalized integral:

$$\mu_{\Phi}^p(A) = \begin{cases} \frac{1}{\mathcal{R}_{\Phi}^p} \cdot \int_T \int_{\mathbb{R}_{\geq 0}} \dim(\mathbb{PDH}^p(\Phi)(\varepsilon, t)) \cdot \chi_A(\varepsilon, t) \, d\varepsilon dt, & \text{if } \mathcal{R}_{\Phi}^p \neq 0, \\ 0, & \text{if } \mathcal{R}_{\Phi}^p = 0, \end{cases}$$

where  $\mathcal{R}_{\Phi}^p := \int_T \int_{\mathbb{R}_{\geq 0}} \dim(\mathbb{PDH}^p(\Phi)(\varepsilon, t)) \, d\varepsilon dt$  is the total integral average of the dimension of the  $p$ -th cohomology group (the normalization factor).

Due to the indecomposability of multi-parameter persistence modules, we cannot decompose  $\mathbb{PDH}^p(\Phi)$  into a direct sum of interval modules as in the single-parameter case. The dynamic

topological feature measure provides an alternative description, capturing the “density distribution” of topological features over the parameter space rather than discrete “lifetimes”.

Stability is the key guarantee for the practical utility of persistent homology. In the two-parameter setting, we introduce a new metric tool.

**Definition 4.8** (Dynamic Interpolation Distance). Let  $M, N : \mathcal{P} \rightarrow \mathbf{Vec}_K$  be two two-parameter persistence modules. Their **dynamic interpolation distance** is defined as:

$$d_{DI}(M, N) = \inf_{\gamma} \sup_{(\varepsilon, t) \in \mathcal{P}} \|\gamma_{(\varepsilon, t)}\|_{\text{op}},$$

where  $\|\cdot\|_{\text{op}}$  denotes the operator norm of linear maps between finite-dimensional vector spaces, and  $\gamma$  ranges over all contravariant natural transformations from  $M$  to  $N$  satisfying:

1. For any  $(\varepsilon, t) \in \mathcal{P}$ ,  $\gamma_{(\varepsilon, t)} : M(\varepsilon, t) \rightarrow N(\varepsilon, t)$  is a  $K$ -linear map;
2. For any pair  $(\varepsilon', t') \preceq (\varepsilon, t)$ , the transition maps of  $M$  and  $N$  go from  $(\varepsilon, t)$  to  $(\varepsilon', t')$  and are compatible with  $\gamma$ :

$$\begin{array}{ccc} M(\varepsilon, t) & \xrightarrow{\gamma_{(\varepsilon, t)}} & N(\varepsilon, t) \\ \downarrow r_{(\varepsilon', t')}^{(\varepsilon, t)} & & \downarrow \rho_{(\varepsilon', t')}^{(\varepsilon, t)} \\ M(\varepsilon', t') & \xrightarrow{\gamma_{(\varepsilon', t')}} & N(\varepsilon', t') \end{array}$$

Here  $r_{(\varepsilon', t')}^{(\varepsilon, t)}$  and  $\rho_{(\varepsilon', t')}^{(\varepsilon, t)}$  are the transition morphisms of the contravariant persistence modules.

When  $M = N$ , the identity natural transformation  $\gamma = \text{id}$  satisfies  $\|\gamma_{(\varepsilon, t)}\|_{\text{op}} = 1$ , so  $d_{DI}(M, M) = 0$ .

**Theorem 4.9** (Stability Theorem). *Under the defining conditions of persistent dynamic cohomology, the  $p$ -th persistent dynamic cohomology is Lipschitz stable with respect to the dynamic interpolation distance: for any two dynamic embedding families  $\Phi$  and  $\Psi$ ,*

$$d_{DI}(\text{PDHP}(\Phi), \text{PDHP}(\Psi)) \leq C \cdot d_{C^1}(\Phi, \Psi),$$

where

$$d_{C^1}(\Phi, \Psi) = \sup_{t \in T, x \in M} (\|\Phi(t, x) - \Psi(t, x)\|_{\mathbb{R}^D} + \|\mathbf{J}\Phi(t, x) - \mathbf{J}\Psi(t, x)\|_{\text{op}})$$

is the  $C^1$  dynamic distance, and  $C$  is a constant depending only on the manifold  $M$  and the parameter space  $T$ .

The stability theorem ensures that our theory is robust to noise and small perturbations in the data.

**Theorem 4.10** (Convergence Theorem). *Under the defining conditions of persistent dynamic cohomology, let  $\{\Phi_n\}$  be a sequence of dynamic embedding families converging to  $\Phi$  in the  $C^1$  topology. Then the corresponding  $p$ -th persistent dynamic cohomology modules converge:*

$$\lim_{n \rightarrow \infty} \text{PDHP}(\Phi_n) = \text{PDHP}(\Phi)$$

with convergence in the dynamic interpolation distance.

The convergence theorem provides a theoretical foundation for numerical approximation, ensuring that discrete approximations preserve topological features in the limit.

Although the indecomposability of multi-parameter persistence modules limits the application of traditional barcode methods, dynamic topological feature measures and stability theory still provide powerful tools for analyzing dynamic topological features. These tools do not depend on decomposition theorems but directly on the global structural properties of persistence modules.

### 4.3 STATISTICS AND ANALYSIS OF DYNAMIC TOPOLOGICAL FEATURES

Having established a rigorous mathematical framework for dynamic topological feature measures and proved their stability, we now explore the rich geometric and statistical information encoded in these measures.

Based on the dynamic topological feature measure  $\mu_\Phi^p$ , we can introduce various statistical moments to describe the distributional properties of topological features, providing powerful tools for quantifying topological evolution.

*Definition 4.11* (Moments of Topological Features). The  **$k$ -th moment** of the dynamic topological feature measure  $\mu_\Phi^p$  is defined as:

$$M_k^p(\Phi) = \int_{\mathcal{P}} d((\varepsilon, t), (\varepsilon_0, t_0))^k d\mu_\Phi^p(\varepsilon, t),$$

where  $d$  is a suitable metric on  $\mathcal{P}$  (e.g., Euclidean), and  $(\varepsilon_0, t_0)$  is a reference point (e.g., the origin or the barycenter of the distribution).

In fact, it is clear that:

- The first moment  $M_1^p$  describes the “center of mass” of the topological features;
- The second moment  $M_2^p$  describes their dispersion.

Specifically, we may define directional moments by

$$M_{k,\varepsilon}^p := \int_{\mathcal{P}} d((\varepsilon, t_0), (\varepsilon_0, t_0))^k d\mu_\Phi^p(\varepsilon, t), \quad M_{k,t}^p := \int_{\mathcal{P}} d((\varepsilon_0, t), (\varepsilon_0, t_0))^k d\mu_\Phi^p(\varepsilon, t).$$

The first moments then have clear geometric meaning:

1. A larger  $M_{1,\varepsilon}^p$  indicates that  $p$ -dimensional topological features appear mainly at large scales;
2.  $M_{1,t}^p$  reflects the average time of appearance along the time axis.

Similarly, the magnitude of  $M_2^p$  quantifies the concentration of topological features over the parameter space.

To compare topological features between different dynamic processes, we introduce the following Wasserstein distance to measure the discrepancy between  $p$ -dimensional topological feature distributions. For background on Wasserstein distances, we refer to the survey by Victor M. Panaretos and Yoav Zemel (2019) [PZ19], and in particular, the following definition is inspired by [PZ19] Eq. (2) in Section 1.2 with  $p = 1$ .

*Definition 4.12* (1st-order Wasserstein Topological Distance). Let dynamic embedding families  $\Phi, \Psi : T \times M \rightarrow \mathbb{R}^D$  have  $p$ -dimensional dynamic topological feature measures  $\mu_\Phi^p, \mu_\Psi^p : \mathcal{B}(\mathcal{P}) \rightarrow \mathbb{R}_{\geq 0}$ , both being probability measures<sup>3</sup>, i.e.,  $\mu_\Phi^p(\mathcal{P}) = \mu_\Psi^p(\mathcal{P}) = 1$ .

<sup>3</sup>This satisfies the prerequisite for Wasserstein distances, see [PZ19], Section 1.2.

The **1st-order Wasserstein topological distance** between  $\Phi$  and  $\Psi$  is defined by

$$d_W^{1,p}(\Phi, \Psi) = \inf_{\pi \in \Pi(\mu_\Phi^p, \mu_\Psi^p)} \int_{\mathcal{P} \times \mathcal{P}} d_{\mathcal{P}}((\varepsilon_1, t_1), (\varepsilon_2, t_2)) \, d\pi(\varepsilon_1, t_1; \varepsilon_2, t_2),$$

where the metric  $d_{\mathcal{P}}$  on the parameter space is the weighted product Euclidean metric<sup>4</sup>

$$d_{\mathcal{P}}((\varepsilon_1, t_1), (\varepsilon_2, t_2)) = \sqrt{\omega_\varepsilon \cdot (\varepsilon_1 - \varepsilon_2)^2 + \omega_t \cdot d_T(t_1, t_2)^2},$$

with  $\omega_\varepsilon > 0$ ,  $\omega_t > 0$  weight coefficients, and where  $d_T(t_1, t_2) = |t_1 - t_2|$  if  $T \subseteq \mathbb{R}$ .

The set of couplings  $\Pi(\mu_\Phi^p, \mu_\Psi^p)$  consists of all joint probability measures with marginals  $\mu_\Phi^p$  and  $\mu_\Psi^p$ , i.e., for all  $\pi \in \Pi(\mu_\Phi^p, \mu_\Psi^p)$ ,

$$\int_{\mathcal{P}} d\pi(\varepsilon_1, t_1; \varepsilon_2, t_2) = \mu_\Phi^p(d\varepsilon_1, dt_1), \quad \int_{\mathcal{P}} d\pi(\varepsilon_1, t_1; \varepsilon_2, t_2) = \mu_\Psi^p(d\varepsilon_2, dt_2).$$

**Theorem 4.13** (Stability of the Wasserstein Topological Distance). *Let  $\Phi, \Psi : T \times M \rightarrow \mathbb{R}^D$  be  $C^1$  dynamic embedding families, and let  $\mu_\Phi^p, \mu_\Psi^p$  be their  $p$ -dimensional dynamic topological feature probability measures. Then the 1st-order Wasserstein topological distance is Lipschitz stable with respect to the  $C^1$  dynamic distance  $d_{C^1}$ :*

$$d_W^{1,p}(\Phi, \Psi) \leq C \cdot d_{C^1}(\Phi, \Psi),$$

where  $C > 0$  is a constant depending only on  $M, T, D$ , and the weights  $\omega_\varepsilon, \omega_t$  in the parameter space metric  $d_{\mathcal{P}}$ , independent of  $\Phi$  and  $\Psi$ .

In dynamic topological data analysis, topological features at different scales often encode distinct geometric and topological information. Fine-scale (small  $\varepsilon$ ) features reflect local geometric structure, while coarse-scale (large  $\varepsilon$ ) features reveal global topological properties of the data. To systematically characterize the distribution of topological features across scales, we introduce the following scale spectrum.

**Definition 4.14** (Scale Spectrum of Topological Features). For a dynamic embedding family  $\Phi$ , the **scale spectrum of its  $p$ -dimensional topological features** is the function  $S_\Phi^p : \mathbb{R}_{\geq 0} \rightarrow [0, 1]$  given by

$$S_\Phi^p(\varepsilon_0) = \mu_\Phi^p(\{(\varepsilon, t) \in \mathcal{P} \mid \varepsilon \leq \varepsilon_0\}),$$

where  $\mu_\Phi^p$  is the dynamic topological feature probability measure (a Borel probability measure).

Since  $\{(\varepsilon, t) \in \mathcal{P} \mid \varepsilon \leq \varepsilon_0\} \in \mathcal{B}(\mathcal{P})$ , the measure is well-defined.

The scale spectrum  $S_\Phi^p(\varepsilon_0)$  has an intuitive meaning: it is the total proportion of  $p$ -dimensional topological features with scale at most  $\varepsilon_0$  over the full parameter space  $\mathcal{P}$ .

Since  $S_\Phi^p(\varepsilon_0)$  is essentially the marginal cumulative distribution function of topological features over the scale parameter  $\varepsilon$  (ignoring variation in time), it increases monotonically from 0 to 1 as  $\varepsilon_0$  goes from 0 to  $+\infty$ , giving clear geometric intuition:

1. If  $S_\Phi^p(\varepsilon_0)$  rapidly approaches 1 at small  $\varepsilon_0$ , then  $p$ -dimensional topological features are concentrated at fine scales;
2. If  $S_\Phi^p(\varepsilon_0)$  approaches 1 slowly at large  $\varepsilon_0$ , then topological features remain significant at coarse scales.

---

<sup>4</sup>Adapted to the product structure of  $\mathcal{P}$ , consistent with the general metric space formulation in the survey, see [PZ19], Section 1.2.

This visually encodes the scale preference of dynamic topological features.

**Proposition 4.15** (Properties of the Scale Spectrum). *The scale spectrum  $S_{\Phi}^p(\varepsilon_0)$  is a non-decreasing function of  $\varepsilon_0$ , and*

$$\lim_{\varepsilon_0 \rightarrow 0} S_{\Phi}^p(\varepsilon_0) = 0, \quad \lim_{\varepsilon_0 \rightarrow \infty} S_{\Phi}^p(\varepsilon_0) = 1.$$

## 5 CONCLUSIONS AND OUTLOOK

This paper systematically constructs a unified mathematical framework for dynamic Čech cohomology and sheaf theory on manifolds. The main contributions are as follows:

1. Proposing a rigorous mathematical definition of the dynamic Čech complex and establishing the concept of continuous complex flow. Extending the classical Čech cohomology theory from the static setting to the dynamic setting, enriching the application scope of algebraic topology in data analysis.
2. Constructing a theory of dynamic Čech sheaves. Through dynamic star neighborhoods and local consistency conditions, a local–global correspondence for the continuous evolution of topological features is established, providing a new mathematical language for dynamic topological analysis.
3. The persistent dynamic cohomology module  $\mathbb{PDH}^p(\Phi)$  is defined, and dynamic topological feature measures are introduced to characterize the distribution of topological features in the scale–time parameter space, effectively avoiding the indecomposability issue of multi-parameter persistent modules.

Based on the framework established in this paper, further research can be carried out in the following directions:

1. Further taking the dimension  $p$  as an additional parameter, regarding the persistent dynamic cohomology module as a three-parameter persistent module  $\mathbb{PDH}^\bullet(\Phi)$ . The correlation between different dimensions remains a challenging problem.
2. Developing efficient parallel computing algorithms to handle larger-scale dynamic data, such as video sequences and sensor network data.
3. Conducting applied research in biomedicine, fluid mechanics, cosmology, and other fields to explore the value of dynamic topological features in the analysis of complex systems.
4. Combining dynamic topological features with deep learning to develop topology-aware neural network models and enhance the understanding of dynamic data.
5. Investigating online dynamic topological analysis algorithms to realize real-time topological monitoring and anomaly detection for streaming data.

The unified framework of dynamic Čech cohomology and sheaf theory on manifolds established in this paper not only provides new mathematical tools for topological data analysis, but also opens up a new research direction for understanding the evolution of topological structures in dynamic systems. With the improvement of computing power and the expansion of application demands, this framework is expected to play an important role in more fields and promote the in-depth development of topological data analysis theory and applications.



## A PROOF OF STABILITY THEOREM 4.9 AND THEOREM 4.13

*Proof.* There exists a uniform Lipschitz constant  $c > 0$  such that  $\|\mathbf{J}\Phi_t(x)\| \leq c$  and  $\|\mathbf{J}\Psi_t(x)\| \leq c$  hold for all  $x \in M$  and  $t \in T$ .

Let  $\delta = d_{C^1}(\Phi, \Psi)$ . We will prove that there exists a constant  $C > 0$  such that

$$d_{DI}(\mathbb{PDHP}(\Phi), \mathbb{PDHP}(\Psi)) \leq C \cdot \delta.$$

We first establish uniform boundedness of the Hausdorff distance.

For any fixed  $t \in T$ , consider the Hausdorff distance between the embedded images  $\Phi_t(M) \subseteq \mathbb{R}^D$  and  $\Psi_t(M) \subseteq \mathbb{R}^D$ :

$$d_H(\Phi_t(M), \Psi_t(M)) = \max \left\{ \sup_{x \in M} \inf_{y \in M} \|\Phi_t(x) - \Psi_t(y)\|, \sup_{y \in M} \inf_{x \in M} \|\Phi_t(x) - \Psi_t(y)\| \right\}.$$

By  $C^1$ -continuity of the dynamic embeddings,  $\|\Phi_t(x) - \Psi_t(x)\| \leq \delta$  for all  $x \in M$ . By Lipschitz continuity of  $\Phi_t$ , for any  $y \in M$ :

$$\|\Phi_t(x) - \Psi_t(y)\| \leq \|\Phi_t(x) - \Phi_t(y)\| + \|\Phi_t(y) - \Psi_t(y)\| \leq c \cdot d_M(x, y) + \delta.$$

Taking  $y = x$  yields  $\inf_{y \in M} \|\Phi_t(x) - \Psi_t(y)\| \leq \delta$ . Similarly,  $\inf_{x \in M} \|\Phi_t(x) - \Psi_t(y)\| \leq \delta$ . Therefore:

$$d_H(\Phi_t(M), \Psi_t(M)) \leq \delta. \tag{A.1}$$

Now, for any  $(\varepsilon, t) \in \mathcal{P}$ , we have the property:

$$|\mathcal{C}_{\Phi, \varepsilon}(t)| \subseteq |\mathcal{C}_{\Psi, \varepsilon + \delta}(t)|, \quad |\mathcal{C}_{\Psi, \varepsilon}(t)| \subseteq |\mathcal{C}_{\Phi, \varepsilon + \delta}(t)|,$$

where  $|\cdot|$  denotes the geometric realization. For instance, the first inclusion can also be expressed as:

$$\forall \sigma \in \mathcal{C}_{\Phi, \varepsilon}(t), \exists \tau \in \mathcal{C}_{\Psi, \varepsilon + \delta}(t) \quad \text{s.t.} \quad \dim \sigma = \dim \tau \quad \text{and} \quad \sigma \simeq \tau.$$

A brief justification follows. Take any simplex  $\sigma \in \mathcal{C}_{\Phi, \varepsilon}(t)$ . By definition of the dynamic Čech complex, there exists  $z \in \mathbb{R}^D$  such that  $z \in \bigcap_{x \in \sigma} B(\Phi_t(x), \varepsilon)$ . For any  $x \in \sigma$ , by the Hausdorff bound equation (A.1), there exists  $y_x \in M$  such that  $\|\Phi_t(x) - \Psi_t(y_x)\| \leq \delta$ . By the triangle inequality:

$$\|\Psi_t(y_x) - z\| \leq \|\Psi_t(y_x) - \Phi_t(x)\| + \|\Phi_t(x) - z\| < \delta + \varepsilon.$$

Thus  $z \in B(\Psi_t(y_x), \varepsilon + \delta)$ . Let  $\tau = \{y_x \mid x \in \sigma\}$ . Then

$$z \in \bigcap_{y_x \in \tau} B(\Psi_t(y_x), \varepsilon + \delta) \neq \emptyset,$$

so  $\tau \in \mathcal{C}_{\Psi, \varepsilon + \delta}(t)$ . Moreover,  $\dim \tau = \dim \sigma$ , and by the Nerve Lemma 2.29, the geometric realizations of  $\sigma$  and  $\tau$  are homotopy equivalent  $\sigma \simeq \tau$ . Hence the topological structure of  $\sigma$  is fully preserved in  $\mathcal{C}_{\Psi, \varepsilon + \delta}(t)$ .

The second inclusion follows similarly.

Accordingly, there exists a simplicial morphism  $f_{\Phi \rightarrow \Psi} : \mathcal{C}_{\Phi, \varepsilon}(t) \rightarrow \mathcal{C}_{\Psi, \varepsilon + \delta}(t)$  satisfying  $f_{\Phi \rightarrow \Psi}(x) = y_x$  and  $f_{\Phi \rightarrow \Psi}(\sigma) = \tau$ .

By basic properties of algebraic topology, a simplicial morphism induces a cohomology homomorphism:

$$f_{\Phi \rightarrow \Psi}^* : H^p(\mathcal{C}_{\Psi, \varepsilon + \delta}(t); K) \rightarrow H^p(\mathcal{C}_{\Phi, \varepsilon}(t); K).$$

Since  $\sigma \simeq \tau$ , the geometric realizations of  $\mathcal{C}_{\Phi, \varepsilon}(t)$  and  $\mathcal{C}_{\Psi, \varepsilon + \delta}(t)$  are homotopy equivalent, so  $f_{\Phi \rightarrow \Psi}^*$  is an isomorphism.

On the other hand, by definition of persistent dynamic cohomology, the inclusion  $j_{\Psi} : \mathcal{C}_{\Psi, \varepsilon}(t) \hookrightarrow \mathcal{C}_{\Psi, \varepsilon + \delta}(t)$  induces a transition homomorphism:

$$\psi_{(\varepsilon, t)}^{(\varepsilon + \delta, t)} : H^p(\mathcal{C}_{\Psi, \varepsilon + \delta}(t); K) \rightarrow H^p(\mathcal{C}_{\Psi, \varepsilon}(t); K).$$

Define the linear map  $\gamma_{(\varepsilon, t)} : \mathbb{PDH}^p(\Phi)(\varepsilon, t) \rightarrow \mathbb{PDH}^p(\Psi)(\varepsilon, t)$  by:

$$\gamma_{(\varepsilon, t)} = \psi_{(\varepsilon, t)}^{(\varepsilon + \delta, t)} \circ (f_{\Phi \rightarrow \Psi}^*)^{-1}.$$

Since  $M$  is a compact Riemannian manifold, both  $\mathcal{C}_{\Phi, \varepsilon}(t)$  and  $\mathcal{C}_{\Psi, \varepsilon}(t)$  are finite simplicial complexes, and their cohomology groups are finite-dimensional vector spaces. Combining the following two facts:

1. The operator norm of the isomorphism  $\|(f_{\Phi \rightarrow \Psi}^*)^{-1}\|_{\text{op}} \leq C_1$ , where  $C_1$  is a constant depending only on the compactness of  $M$ , as the family of simplicial morphisms is uniformly bounded over the compact manifold;
2. The operator norm of the transition homomorphism  $\|\psi_{(\varepsilon, t)}^{(\varepsilon + \delta, t)}\|_{\text{op}} \leq C_2$ , where  $C_2$  is controlled by the combinatorial complexity of  $\mathcal{C}_{\Psi, \varepsilon}(t)$ , hence by the dimension  $d$  and compactness of  $M$ ,

by submultiplicativity of the operator norm:

$$\|\gamma_{(\varepsilon, t)}\|_{\text{op}} \leq \|\psi_{(\varepsilon, t)}^{(\varepsilon + \delta, t)}\|_{\text{op}} \cdot \|(f_{\Phi \rightarrow \Psi}^*)^{-1}\|_{\text{op}} \leq C_1 C_2. \quad (\text{A.2})$$

We verify that  $\gamma = \{\gamma_{(\varepsilon, t)}\}_{(\varepsilon, t) \in \mathcal{P}}$  satisfies functorial compatibility for the dynamic interpolation distance, i.e., for any  $(\varepsilon_1, t_1) \preceq (\varepsilon_2, t_2)$ , the following diagram commutes:

$$\begin{array}{ccc} \mathbb{PDH}^p(\Phi)(\varepsilon_2, t_2) & \xrightarrow{\gamma_{(\varepsilon_2, t_2)}} & \mathbb{PDH}^p(\Psi)(\varepsilon_2, t_2) \\ \varphi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)} \downarrow & & \downarrow \psi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)} \\ \mathbb{PDH}^p(\Phi)(\varepsilon_1, t_1) & \xrightarrow{\gamma_{(\varepsilon_1, t_1)}} & \mathbb{PDH}^p(\Psi)(\varepsilon_1, t_1) \end{array}$$

Substitute  $\gamma_{(\varepsilon, t)} = \psi_{(\varepsilon, t)}^{(\varepsilon + \delta, t)} \circ (f_{\Phi \rightarrow \Psi}^*)^{-1}$  and expand both compositions:

$$\begin{aligned} \text{Left-hand side : } & \gamma_{(\varepsilon_1, t_1)} \circ \varphi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)} = \psi_{(\varepsilon_1, t_1)}^{(\varepsilon_1 + \delta, t_1)} \circ (f_{\Phi \rightarrow \Psi}^*)^{-1} \circ \varphi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)}, \\ \text{Right-hand side : } & \psi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)} \circ \gamma_{(\varepsilon_2, t_2)} = \psi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)} \circ \psi_{(\varepsilon_2, t_2)}^{(\varepsilon_2 + \delta, t_2)} \circ (f_{\Phi \rightarrow \Psi}^*)^{-1}. \end{aligned}$$

By naturality of transition maps in persistent dynamic cohomology—meaning that  $f_{\Phi \rightarrow \Psi}$  commutes with transition maps—we have

$$(f_{\Phi \rightarrow \Psi}^*)^{-1} \circ \varphi_{(\varepsilon_1, t_1)}^{(\varepsilon_2, t_2)} = \psi_{(\varepsilon_1 + \delta, t_1)}^{(\varepsilon_2 + \delta, t_2)} \circ (f_{\Phi \rightarrow \Psi}^*)^{-1},$$

where  $\psi_{(\varepsilon_1+\delta, t_1)}^{(\varepsilon_2+\delta, t_2)} : \text{PDHP}(\Psi)(\varepsilon_1 + \delta, t_1) \rightarrow \text{PDHP}(\Psi)(\varepsilon_2 + \delta, t_2)$ . Substituting into the left-hand side:

$$\text{LHS} = \psi_{(\varepsilon_1, t_1)}^{(\varepsilon_1+\delta, t_1)} \circ \psi_{(\varepsilon_1+\delta, t_1)}^{(\varepsilon_2+\delta, t_2)} \circ (f_{\Phi \rightarrow \Psi}^*)^{-1}.$$

Therefore  $\text{LHS} = \text{RHS}$ , and functorial compatibility is proved.

From the above construction,  $\gamma = \{\gamma_{(\varepsilon, t)}\}$  is a valid family of morphisms satisfying functorial compatibility, with operator norm bounded by  $\|\gamma_{(\varepsilon, t)}\|_{\text{op}} \leq C_1 C_2$ .

We now relate this norm bound to the  $C^1$ -distance  $\delta = d_{C^1}(\Phi, \Psi)$ .

Since  $M$  is a compact Riemannian manifold and the dynamic embedding  $\Psi$  is uniformly Lipschitz with constant  $c$ , the cohomological perturbation induced by scaling from  $\varepsilon$  to  $\varepsilon + \delta$  is linearly controllable. Specifically, the operator norm of the inverse transition map satisfies

$$\left\| \psi_{(\varepsilon, t)}^{(\varepsilon+\delta, t)} \right\|_{\text{op}} \leq c' \cdot \delta,$$

where  $c' > 0$  is a universal constant depending only on  $M$  and the Lipschitz constant  $c$ , independent of  $\Phi$  and  $\Psi$ .

Note that  $C_1$  is also a constant depending only on the compactness of  $M$ . Set  $C = C_1 c'$ , which is still a constant depending only on the ambient manifold  $M$ . Then

$$C_1 C_2 \leq C_1 \cdot c' \cdot \delta = C \cdot \delta.$$

Thus the operator norm of the morphism family  $\gamma$  satisfies the uniform bound

$$\sup_{(\varepsilon, t) \in \mathcal{P}} \|\gamma_{(\varepsilon, t)}\|_{\text{op}} \leq C \cdot \delta = C \cdot d_{C^1}(\Phi, \Psi).$$

By definition of the dynamic interpolation distance:

$$d_{DI}(\text{PDHP}(\Phi), \text{PDHP}(\Psi)) = \inf_{\gamma} \sup_{(\varepsilon, t) \in \mathcal{P}} \|\gamma_{(\varepsilon, t)}\|_{\text{op}},$$

where the infimum is taken over all functorially compatible morphism families  $\gamma$ . Since we have shown that  $\gamma$  is an admissible morphism family, its supremum gives an upper bound for the dynamic interpolation distance, so

$$d_{DI}(\text{PDHP}(\Phi), \text{PDHP}(\Psi)) \leq C \cdot d_{C^1}(\Phi, \Psi).$$

The stability for the first-order Wasserstein distance  $d_W^{1,p}(\Phi, \Psi)$  between the dynamic topological feature measures  $\mu_{\Phi}^p$  and  $\mu_{\Psi}^p$  follows by analogous arguments.

In summary, there exist constants  $C, C' > 0$  depending only on the manifold  $M$  and the Lipschitz constant  $c$  of the dynamic embeddings such that:

$$d_{DI}(\text{PDHP}(\Phi), \text{PDHP}(\Psi)) \leq C \cdot d_{C^1}(\Phi, \Psi), \quad d_W^{1,p}(\Phi, \Psi) \leq C' \cdot d_{C^1}(\Phi, \Psi).$$

□

## B PROOF OF CONVERGENCE THEOREM 4.10 AND ITS COROLLARIES

In the main text, the convergence theorem 4.10 is stated as follows: if a sequence of dynamic embedding families  $\{\Phi_n\}$  converges to  $\Phi$  in the  $C^1$  topology, then the corresponding  $p$ -th persistent dynamic cohomology modules  $\mathbb{PDH}^p(\Phi_n)$  converge to  $\mathbb{PDH}^p(\Phi)$  with respect to the dynamic interpolation distance. It is natural to conjecture that the associated topological feature measures, Wasserstein distances, topological moments, and scale spectra also converge accordingly.

### B.1 CONVERGENCE IN THE DYNAMIC INTERPOLATION DISTANCE

Recall the definition of  $C^1$ -convergence: A sequence of dynamic embedding families  $\{\Phi_n\}$  converges to  $\Phi$  in the  $C^1$  topology if

$$\lim_{n \rightarrow \infty} d_{C^1}(\Phi_n, \Phi) = 0,$$

where

$$d_{C^1}(\Phi, \Psi) = \sup_{t \in T, x \in M} \left( \|\Phi(t, x) - \Psi(t, x)\|_{\mathbb{R}^N} + \|\mathbf{J}\Phi(t, x) - \mathbf{J}\Psi(t, x)\|_{\text{op}} \right).$$

For any  $\varepsilon > 0$ , there exists  $N \in \mathbb{N}$  such that for all  $n \geq N$ ,  $d_{C^1}(\Phi_n, \Phi) < \varepsilon$ .

By the stability theorem 4.9, there exists a universal constant  $C > 0$  depending only on the manifold  $M$ , the time interval  $T$ , and the Lipschitz constant, such that for any two dynamic embeddings  $\Psi_1, \Psi_2$ ,

$$d_{DI}(\mathbb{PDH}^p(\Psi_1), \mathbb{PDH}^p(\Psi_2)) \leq C \cdot d_{C^1}(\Psi_1, \Psi_2).$$

Substituting  $\Psi_1 = \Phi_n$  and  $\Psi_2 = \Phi$ , we obtain

$$d_{DI}(\mathbb{PDH}^p(\Phi_n), \mathbb{PDH}^p(\Phi)) \leq C \cdot d_{C^1}(\Phi_n, \Phi).$$

By the definition of  $C^1$ -convergence,  $\lim_{n \rightarrow \infty} d_{C^1}(\Phi_n, \Phi) = 0$ . Combining the above inequality and the squeeze theorem:

$$0 \leq \lim_{n \rightarrow \infty} d_{DI}(\mathbb{PDH}^p(\Phi_n), \mathbb{PDH}^p(\Phi)) \leq C \cdot \lim_{n \rightarrow \infty} d_{C^1}(\Phi_n, \Phi) = 0.$$

Therefore,

$$\lim_{n \rightarrow \infty} d_{DI}(\mathbb{PDH}^p(\Phi_n), \mathbb{PDH}^p(\Phi)) = 0,$$

i.e.,  $\mathbb{PDH}^p(\Phi_n)$  converges to  $\mathbb{PDH}^p(\Phi)$  in the dynamic interpolation distance.

### B.2 OTHER CONVERGENCE RESULTS

Convergence in the dynamic interpolation distance implies pointwise convergence of the Betti number function:

$$\lim_{n \rightarrow \infty} \beta_{\Phi_n}^p(\varepsilon, t) = \beta_{\Phi}^p(\varepsilon, t), \quad \forall (\varepsilon, t) \in \mathcal{P}.$$

*Proof.* By assumption,  $\lim_{n \rightarrow \infty} d_{DI}(\mathbb{PDH}^p(\Phi_n), \mathbb{PDH}^p(\Phi)) = 0$ , meaning that for any  $\epsilon > 0$ , there exists  $N \in \mathbb{N}$  such that for all  $n > N$ ,

$$\sup_{(\varepsilon, t) \in \mathcal{P}} \|\gamma_{n, (\varepsilon, t)}\|_{\text{op}} < \epsilon.$$

For any fixed  $(\varepsilon, t) \in \mathcal{P}$ , the Betti number  $\beta_{\Phi}^p(\varepsilon, t) = \dim(\mathbb{PDH}^p(\Phi)(\varepsilon, t))$  is a nonnegative integer characterizing the topological information of the cohomology group.

When  $n$  is sufficiently large, the difference in topological structure between  $\mathbb{PDH}^p(\Phi_n)(\varepsilon, t)$  and  $\mathbb{PDH}^p(\Phi)(\varepsilon, t)$  tends to zero, so their core topological invariant (the Betti number) must converge to the same value<sup>5</sup>, i.e.,  $\lim_{n \rightarrow \infty} \beta_{\Phi_n}^p(\varepsilon, t) = \beta_{\Phi}^p(\varepsilon, t)$ .  $\square$

Since both  $M$  and  $T$  are compact manifolds,  $\mathcal{P} = \mathbb{R}_{\geq 0} \times T$  is a locally compact, separable metric space. Moreover, there exists a constant  $B > 0$  depending only on the topology of  $M$  such that for any dynamic embedding  $\Psi$ ,

$$0 \leq \beta_{\Psi}^p(\varepsilon, t) \leq B, \quad \forall (\varepsilon, t) \in \mathcal{P}.$$

From the pointwise convergence  $\beta_{\Phi_n}^p \rightarrow \beta_{\Phi}^p$  and Lebesgue's dominated convergence theorem, for any Borel set  $A \subseteq \mathcal{P}$ ,

$$\lim_{n \rightarrow \infty} \int_A \beta_{\Phi_n}^p(\varepsilon, t) d\varepsilon dt = \int_A \beta_{\Phi}^p(\varepsilon, t) d\varepsilon dt.$$

For the normalization factor  $\mathcal{R}_{\Phi}^p = \int_{\mathcal{P}} \beta_{\Phi}^p(\varepsilon, t) d\varepsilon dt$ , taking  $A = \mathcal{P}$  gives  $\lim_{n \rightarrow \infty} \mathcal{R}_{\Phi_n}^p = \mathcal{R}_{\Phi}^p$ .

Recall that the dynamic topological feature measure is defined by

$$\mu_{\Phi_n}^p(A) = \frac{1}{\mathcal{R}_{\Phi_n}^p} \int_A \beta_{\Phi_n}^p(\varepsilon, t) d\varepsilon dt, \quad \mu_{\Phi}^p(A) = \frac{1}{\mathcal{R}_{\Phi}^p} \int_A \beta_{\Phi}^p(\varepsilon, t) d\varepsilon dt.$$

Combining  $\mathcal{R}_{\Phi_n}^p \rightarrow \mathcal{R}_{\Phi}^p$  and the convergence of integrals, we obtain

$$\lim_{n \rightarrow \infty} \mu_{\Phi_n}^p(A) = \frac{1}{\mathcal{R}_{\Phi}^p} \int_A \beta_{\Phi}^p(\varepsilon, t) d\varepsilon dt = \mu_{\Phi}^p(A), \quad \forall \text{ Borel } A \subseteq \mathcal{P},$$

i.e.,  $\mu_{\Phi_n}^p$  converges weakly to  $\mu_{\Phi}^p$ .

Since  $\mathcal{P}$  is a locally compact, separable metric space, all dynamic topological feature measures  $\mu_{\Phi_n}^p$  and  $\mu_{\Phi}^p$  are supported on a common compact subset of  $\mathcal{P}$ , and their density functions (the Betti number functions) are uniformly bounded. Together with the weak convergence of measures

$$\lim_{n \rightarrow \infty} \mu_{\Phi_n}^p(A) = \mu_{\Phi}^p(A), \quad \forall \text{ Borel set } A \subseteq \mathcal{P},$$

we obtain convergence of the first-order Wasserstein distance:

$$\lim_{n \rightarrow \infty} d_W^{1,p}(\Phi_n, \Phi) = 0.$$

Furthermore, for any  $k \geq 1$ , the topological moment is  $M_k^p(\Phi) = \int_{\mathcal{P}} \varepsilon^k d\mu_{\Phi}^p(\varepsilon, t)$ . By weak convergence of measures and continuity of integration ( $\varepsilon^k$  is continuous and bounded),

$$\lim_{n \rightarrow \infty} M_k^p(\Phi_n) = \int_{\mathcal{P}} \varepsilon^k d\mu_{\Phi}^p(\varepsilon, t) = M_k^p(\Phi).$$

---

<sup>5</sup>Since  $M$  is a compact manifold, the dimension of  $\mathbb{PDH}^p(\Phi)(\varepsilon, t)$  is bounded (denote the upper bound by  $B$ ), i.e.,  $0 \leq \beta_{\Phi}^p(\varepsilon, t) \leq B$ . The Betti number is a discrete nonnegative integer. If the sequence  $\{\beta_{\Phi_n}^p(\varepsilon, t)\}$  satisfies that the “topological difference tends to zero”, then the sequence must eventually converge to the unique integer  $\beta_{\Phi}^p(\varepsilon, t)$ .

Similarly, the scale spectrum is  $S_{\Phi}^p(\varepsilon) = \int_T \beta_{\Phi}^p(\varepsilon, t) dt$ . For a fixed  $\varepsilon$ , by the dominated convergence theorem,

$$\lim_{n \rightarrow \infty} S_{\Phi_n}^p(\varepsilon) = \int_T \beta_{\Phi}^p(\varepsilon, t) dt = S_{\Phi}^p(\varepsilon).$$

These results collectively form the convergence foundation of the persistent dynamic cohomology theory, providing solid theoretical support for numerical computation and practical applications.

## REFERENCES

- [BCM25] A. Balitskiy, B. Coskunuzer, and F. Mémoli. *Geometric Bounds for Persistence*. 2025. arXiv: [2403.13980](https://arxiv.org/abs/2403.13980) [math.AT]. URL: <https://arxiv.org/abs/2403.13980>.
- [BS25] U. Bauer and L. Scoccola. “Multi-parameter Persistence Modules are Generically Indecomposable”. In: *International Mathematics Research Notices* 2025.5 (2025), rna034. ISSN: 1073-7928. DOI: [10.1093/imrn/rnaf034](https://doi.org/10.1093/imrn/rnaf034). eprint: <https://academic.oup.com/imrn/article-pdf/2025/5/rnaf034/62170563/rnaf034.pdf>. URL: <https://doi.org/10.1093/imrn/rnaf034>.
- [CG09] Carlsson and Gunnar. “TOPOLOGY AND DATA.” In: *Bulletin of the American Mathematical Society* (2009).
- [Car14] G. Carlsson. “Topological pattern recognition for point cloud data”. In: *Acta Numerica* 23.23 (2014), pp. 289–368.
- [CSO13] F. Chazal, V. de Silva, and S. Oudot. *Persistence stability for geometric complexes*. 2013. arXiv: [1207.3885](https://arxiv.org/abs/1207.3885) [math.AT]. URL: <https://arxiv.org/abs/1207.3885>.
- [Día25] J. Díaz Boils. “Persistent Homology on a Lattice of Multigraphs”. In: *International Journal of Topology* 2.2 (2025). ISSN: 2813-9542. DOI: [10.3390/ijtop2020007](https://doi.org/10.3390/ijtop2020007). URL: <https://www.mdpi.com/2813-9542/2/2/7>.
- [DYS] B. Dmitri, B. Yuri, and I. Sergei. *A Course in Metric Geometry*. A course in metric geometry.
- [EH21] H. Edelsbrunner and J. L. Harer. *Computational Topology: An Introduction*. 2021.
- [ELZ02] Edelsbrunner, Letscher, and Zomorodian. “Topological Persistence and Simplification”. In: *Discrete & Computational Geometry* 28.4 (2002), pp. 511–533.
- [Fri25] S. Friedl. *Topology*. 2025.
- [GQ20] J. Gallier and J. Quaintance. *Algebra, Topology, Differential Calculus, and Optimization Theory For Computer Science and Machine Learning*. 2020.
- [Han25] J. Han. “lecture notes of PKU Summer School: Complex Geometry”. In: (2025).
- [Hat02] A. Hatcher. *Algebraic Topology*. 2002.
- [Jr08] R. O. W. Jr. *Differential Analysis on Complex Manifolds*. GTM 65. Springer Science+Business Media, LLC, 2008. ISBN: 978-0-387-90419-0.
- [KS14] M. Kashiwara and P. Schapira. *Sheaves on Manifolds*. 2014.
- [Ott+17] N. Otter, M. A. Porter, U. Tillmann, P. Grindrod, and H. A. Harrington. “A roadmap for the computation of persistent homology”. In: *Epj Data Science* 6.1 (2017), p. 17.
- [Oud15] S. Y. Oudot. *Persistence Theory: From Quiver Representations to Data Analysis*. 2015.
- [PZ19] V. M. Panaretos and Y. Zemel. “Statistical Aspects of Wasserstein Distances”. In: *Annual Review of Statistics and Its Application* 6.1 (2019), pp. 405–431. ISSN: 2326-831X. DOI: [10.1146/annurev-statistics-030718-104938](https://doi.org/10.1146/annurev-statistics-030718-104938). URL: <http://dx.doi.org/10.1146/annurev-statistics-030718-104938>.
- [PH13] J. Perea and J. Harer. “Sliding Windows and Persistence: An Application of Topological Methods to Signal Analysis”. In: *Foundations of Computational Mathematics* 15.3 (2013).
- [Rob14] M. Robinson. *Topological signal processing*. Springer, 2014.
- [SG07] V. D. Silva and R. Ghrist. “Homological sensor networks”. In: *Notices of the American Mathematical Society* 54.1 (2007).
- [Was16] L. Wasserman. “Topological Data Analysis”. In: *Annual Review of Statistics and Its Application* 5.1 (2016).
- [WLY25] H.-T. Wong, D. Lee, and H. Yan. *Towards Scalable Topological Regularizers*. 2025. arXiv: [2501.14641](https://arxiv.org/abs/2501.14641) [cs.LG]. URL: <https://arxiv.org/abs/2501.14641>.