

See discussions, stats, and author profiles for this publication at: <http://www.researchgate.net/publication/253763305>

A Convex Approach for Controlled Lotka Volterra Multi-Species Models

ARTICLE

DOWNLOADS

7

VIEWS

23

3 AUTHORS:



[Sonia Palomino Castro](#)

Federal University of Santa Catarina

9 PUBLICATIONS 2 CITATIONS

SEE PROFILE



[Daniel Coutinho](#)

Federal University of Santa Catarina

142 PUBLICATIONS 507 CITATIONS

SEE PROFILE



[K. A. Barbosa](#)

University of Santiago, Chile

41 PUBLICATIONS 418 CITATIONS

SEE PROFILE

A Convex Approach for Controlled Lotka Volterra Multi-Species Models

S. Palomino Bean

Mathematics Department, Universidade Federal de Santa Catarina - UFSC,
88040-900, Florianópolis, SC, Brazil. E-mail: palomino@mtm.ufsc.br

D. F. Coutinho

Group of Automation and Control Systems, Pontifícia Universidade Católica do Rio Grande do Sul - PUCRS,
Av. Ipiranga 6681, Porto Alegre, RS, 90619-900, Brazil. E-mail: dcoutinho@pucrs.br

K. A. Barbosa

Department of Systems and Control, Laboratório Nacional de Computação Científica - LNCC,
Av. Getúlio Vargas 333, Petrópolis, RJ, 25651-075, Brazil. E-mail: karinab@lncc.br

Keywords: Multi-species models, Lotka Volterra systems, control systems, coexistence, LMIs.

Recently, the coexistence of the multi-species, where the models consider switching or not, are analyzed in the references [1, 2]. The latter results employ a traditional approach and the coexistence of the species is taken into account by means of rational functions (switching terms) as in the references [3, 4, 5]. Since Lotka Volterra systems have more than one equilibrium point, an estimate of the region in which the species coexist is an important issue not yet fully addressed. One can provide an estimate of the coexistence region through the Linear Matrix Inequality (LMI [6]) framework and a Differential-Algebraic representation (DAR) of the system [7, 8]. It turns out that Lotka Volterra systems with no switching have no stable equilibrium points. In this work, a control term to stabilize the system is imposed extending the result obtained in [2]. An example illustrates the proposed methodology, where a system with three species (two prey and one predator in a two trophic level food chain) is considered.

References

- [1] S. Palomino Bean, A. C. S. Vilcarromero, O. Bonato, and J. Fernandes, Co-existência de espécies em sistemas presa-predador com switching [Coexistence of species in predator-prey systems with switching], *Tendências em Matemática Aplicada, TEMA/SBMAC*, vol. 7(2), 317–326, 2006 (in Portuguese).
- [2] S. Palomino Bean and L. M. Porto, Coexistence on predator-prey multi-species with control, 4th Workshop in Mathematical Methods and Modeling of Biophysical Phenomena, (IMPA, Angra dos Reis, RJ), 2006.
- [3] M. Tansky, Switching effect in prey-predator system, *J. Theoretical Biology*, vol. 70, 263–271, 1978.
- [4] E. Kuno, Principles of predator-prey interaction in theoretical, experimental and natural population systems, *Advances in Ecological Research*, vol. 16, 249–337, 1987.
- [5] M. Saleem, A. K. Tripathi, and A. H. Sadiyal, Coexistence of species in a defensive switching model, *Mathematical Biosciences*, vol. 181, 145–164, 2003.
- [6] S. Boyd, L. Ghaoui, E. Feron, and V. Balakrishnan, *Linear matrix inequalities in systems and control theory*. Philadelphia: SIAM Books, 1994.
- [7] S. Palomino Bean, D. F. Coutinho, and K. A. Barbosa, Coexistence of Lotka Volterra Multi-species: an LMI Approach, 6th International Congress on Industrial and Applied Mathematics, (submitted), 2007.
- [8] D. F. Coutinho, C. E. de Souza, and A. Trofino, Regional stability analysis of implicit polynomial systems, *Proc. 45th IEEE Conf. Decision and Control*, (San Diego, CA), 2006.