

# Classical Mechanics Tutorial I

## Engineering Physics

Indian Institute of Information Technology, Allahabad

# Calculus of Multivariable Functions

We would try to look at the introductory part of how one does the calculus of multiple variables sitting in a function.

# Calculus of Multivariable Functions

We would try to look at the introductory part of how one does the calculus of multiple variables sitting in a function.

- ▶ We denote the partial derivative of a function as  $\frac{\partial}{\partial x}$ .

# Calculus of Multivariable Functions

We would try to look at the introductory part of how one does the calculus of multiple variables sitting in a function.

- ▶ We denote the partial derivative of a function as  $\frac{\partial}{\partial x}$ .
- ▶ Let us consider a function  $f(x, y, t)$ ,

$$f(x, y, t) = 2x^3y + 4x^2t + xy \quad (1)$$

Write down the derivatives of  $f(x, y, t)$  with respect to  $x, y$  and  $t$ ?

# Calculus of Multivariable Functions

We would try to look at the introductory part of how one does the calculus of multiple variables sitting in a function.

- ▶ We denote the partial derivative of a function as  $\frac{\partial}{\partial x}$ .
- ▶ Let us consider a function  $f(x, y, t)$ ,

$$f(x, y, t) = 2x^3y + 4x^2t + xy \quad (1)$$

Write down the derivatives of  $f(x, y, t)$  with respect to  $x, y$  and  $t$ ?

$$\frac{\partial f}{\partial x} = f_x = \frac{\partial(2x^3y)}{\partial x} + \frac{\partial(4x^2t)}{\partial x} + \frac{\partial(xy)}{\partial x} = 6x^2y + 8xt + y \quad (2)$$

# Calculus of Multivariable Functions

We would try to look at the introductory part of how one does the calculus of multiple variables sitting in a function.

- ▶ We denote the partial derivative of a function as  $\frac{\partial}{\partial x}$ .
- ▶ Let us consider a function  $f(x, y, t)$ ,

$$f(x, y, t) = 2x^3y + 4x^2t + xy \quad (1)$$

Write down the derivatives of  $f(x, y, t)$  with respect to  $x, y$  and  $t$ ?

$$\frac{\partial f}{\partial x} = f_x = \frac{\partial(2x^3y)}{\partial x} + \frac{\partial(4x^2t)}{\partial x} + \frac{\partial(xy)}{\partial x} = 6x^2y + 8xt + y \quad (2)$$

$$\frac{\partial f}{\partial y} = f_y = \frac{\partial(2x^3y)}{\partial y} + \frac{\partial(4x^2t)}{\partial y} + \frac{\partial(xy)}{\partial y} = 2x^3 + x \quad (3)$$

# Calculus of Multivariable Functions

We would try to look at the introductory part of how one does the calculus of multiple variables sitting in a function.

- ▶ We denote the partial derivative of a function as  $\frac{\partial}{\partial x}$ .
- ▶ Let us consider a function  $f(x, y, t)$ ,

$$f(x, y, t) = 2x^3y + 4x^2t + xy \quad (1)$$

Write down the derivatives of  $f(x, y, t)$  with respect to  $x, y$  and  $t$ ?

$$\frac{\partial f}{\partial x} = f_x = \frac{\partial(2x^3y)}{\partial x} + \frac{\partial(4x^2t)}{\partial x} + \frac{\partial(xy)}{\partial x} = 6x^2y + 8xt + y \quad (2)$$

$$\frac{\partial f}{\partial y} = f_y = \frac{\partial(2x^3y)}{\partial y} + \frac{\partial(4x^2t)}{\partial y} + \frac{\partial(xy)}{\partial y} = 2x^3 + x \quad (3)$$

$$\frac{\partial f}{\partial t} = f_t = \frac{\partial(2x^3y)}{\partial t} + \frac{\partial(4x^2t)}{\partial t} + \frac{\partial(xy)}{\partial t} = 4x^2 \quad (4)$$

- Sometimes, we denote the partial derivative as a suffix to the function. E.g.  $F_x$  means derivative of function  $F$  w.r.t.  $x$ .



- ▶ Sometimes, we denote the partial derivative as a suffix to the function. E.g.  $F_x$  means derivative of function  $F$  w.r.t.  $x$ .
- ▶ When the derivative of a function  $F$  is expressed as  $\frac{dF}{dx}$ , we call it as total derivative. It is also known as the **Leibniz's notation**.

- ▶ Sometimes, we denote the partial derivative as a suffix to the function. E.g.  $F_x$  means derivative of function  $F$  w.r.t.  $x$ .
- ▶ When the derivative of a function  $F$  is expressed as  $\frac{dF}{dx}$ , we call it as total derivative. It is also known as the **Leibniz's notation**.
- ▶ In classical mechanics, we would see *dot* (e.g.  $\dot{x}$ ) for the (*total*) derivative w.r.t. to time parameter, and prime (e.g.  $\frac{dg}{d\omega} = g'(\omega)$ ) can be used for derivative w.r.t. spatial ( $x, y$  or  $z$ ) coordinates.

- ▶ Sometimes, we denote the partial derivative as a suffix to the function. E.g.  $F_x$  means derivative of function  $F$  w.r.t.  $x$ .
- ▶ When the derivative of a function  $F$  is expressed as  $\frac{dF}{dx}$ , we call it as total derivative. It is also known as the **Leibniz's notation**.
- ▶ In classical mechanics, we would see *dot* (e.g.  $\dot{x}$ ) for the (*total*) derivative w.r.t. to time parameter, and prime (e.g.  $\frac{dg}{d\omega} = g'(\omega)$ ) can be used for derivative w.r.t. spatial ( $x, y$  or  $z$ ) coordinates.

Now consider an example of total derivative. Let us consider a function  $f(x(t), y(t), t)$  as

$$f(x(t), y(t), t) = x^2yt + xy^2t + yt^2 \quad (5)$$

Here,  $x$  and  $y$  are also function of  $t$ . Write down  $\frac{df}{dt}$ ?

- ▶ Sometimes, we denote the partial derivative as a suffix to the function. E.g.  $F_x$  means derivative of function  $F$  w.r.t.  $x$ .
- ▶ When the derivative of a function  $F$  is expressed as  $\frac{dF}{dx}$ , we call it as total derivative. It is also known as the **Leibniz's notation**.
- ▶ In classical mechanics, we would see *dot* (e.g.  $\dot{x}$ ) for the (*total*) derivative w.r.t. to time parameter, and prime (e.g.  $\frac{dg}{d\omega} = g'(\omega)$ ) can be used for derivative w.r.t. spatial ( $x, y$  or  $z$ ) coordinates.

Now consider an example of total derivative. Let us consider a function  $f(x(t), y(t), t)$  as

$$f(x(t), y(t), t) = x^2yt + xy^2t + yt^2 \quad (5)$$

Here,  $x$  and  $y$  are also function of  $t$ . Write down  $\frac{df}{dt}$ ?

$$\frac{df}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} + \frac{\partial f}{\partial t} = f_x \dot{x} + f_y \dot{y} + f_t \quad (6)$$

now you can calculate partial derivatives of  $f$ , i.e.,  $f_x$ ,  $f_y$  and  $f_z$ .

$$f_x = 2xyt + y^2t \quad ; \quad f_y = x^2t + 2xyt + t^2 \quad ; \quad f_t = x^2y + xy^2 + 2yt \quad (7)$$

- Now we can replace  $f_x$ ,  $f_y$  and  $f_z$  in Eq.(6) to find the expression for  $\frac{df}{dt}$ .

now you can calculate partial derivatives of  $f$ , i.e.,  $f_x$ ,  $f_y$  and  $f_z$ .

$$f_x = 2xyt + y^2t \quad ; \quad f_y = x^2t + 2xyt + t^2 \quad ; \quad f_t = x^2y + xy^2 + 2yt \quad (7)$$

- ▶ Now we can replace  $f_x$ ,  $f_y$  and  $f_z$  in Eq.(6) to find the expression for  $\frac{df}{dt}$ .
- ▶ One can also compute  $\frac{d}{dt}f_x = \dot{f}_x = \frac{d}{dt}\left(\frac{\partial f}{\partial x}\right)$  as

$$\dot{f}_x = 2(\dot{x}yt + x\dot{y}t + xy) + 2y\dot{y}t + y^2 \quad (8)$$

now you can calculate partial derivatives of  $f$ , i.e.,  $f_x$ ,  $f_y$  and  $f_z$ .

$$f_x = 2xyt + y^2t \quad ; \quad f_y = x^2t + 2xyt + t^2 \quad ; \quad f_t = x^2y + xy^2 + 2yt \quad (7)$$

- ▶ Now we can replace  $f_x$ ,  $f_y$  and  $f_z$  in Eq.(6) to find the expression for  $\frac{df}{dt}$ .

- ▶ One can also compute  $\frac{d}{dt}f_x = \dot{f}_x = \frac{d}{dt}\left(\frac{\partial f}{\partial x}\right)$  as

$$\dot{f}_x = 2(\dot{x}yt + x\dot{y}t + xy) + 2y\dot{y}t + y^2 \quad (8)$$

- ▶ If we would like to find the differential element of the multi-variable function, we implement the *chain rule*. For function  $f(x(t), y(t), t)$ .

now you can calculate partial derivatives of  $f$ , i.e.,  $f_x$ ,  $f_y$  and  $f_z$ .

$$f_x = 2xyt + y^2t \quad ; \quad f_y = x^2t + 2xyt + t^2 \quad ; \quad f_t = x^2y + xy^2 + 2yt \quad (7)$$

- ▶ Now we can replace  $f_x$ ,  $f_y$  and  $f_z$  in Eq.(6) to find the expression for  $\frac{df}{dt}$ .

- ▶ One can also compute  $\frac{d}{dt}f_x = \dot{f}_x = \frac{d}{dt}\left(\frac{\partial f}{\partial x}\right)$  as

$$\dot{f}_x = 2(\dot{x}yt + x\dot{y}t + xy) + 2y\dot{y}t + y^2 \quad (8)$$

- ▶ If we would like to find the differential element of the multi-variable function, we implement the *chain rule*. For function  $f(x(t), y(t), t)$ .

$$df = \frac{\partial f}{\partial x}dx + \frac{\partial f}{\partial y}dy + \frac{\partial f}{\partial t}dt = f_x dx + f_y dy + f_t dt \quad (9)$$



now you can calculate partial derivatives of  $f$ , i.e.,  $f_x$ ,  $f_y$  and  $f_z$ .

$$f_x = 2xyt + y^2t \quad ; \quad f_y = x^2t + 2xyt + t^2 \quad ; \quad f_t = x^2y + xy^2 + 2yt \quad (7)$$

- ▶ Now we can replace  $f_x$ ,  $f_y$  and  $f_z$  in Eq.(6) to find the expression for  $\frac{df}{dt}$ .

- ▶ One can also compute  $\frac{d}{dt}f_x = \dot{f}_x = \frac{d}{dt}\left(\frac{\partial f}{\partial x}\right)$  as

$$\dot{f}_x = 2(\dot{x}yt + x\dot{y}t + xy) + 2y\dot{y}t + y^2 \quad (8)$$

- ▶ If we would like to find the differential element of the multi-variable function, we implement the *chain rule*. For function  $f(x(t), y(t), t)$ .

$$df = \frac{\partial f}{\partial x}dx + \frac{\partial f}{\partial y}dy + \frac{\partial f}{\partial t}dt = f_x dx + f_y dy + f_t dt \quad (9)$$

- ▶ As  $dx = \frac{dx}{dt}dt$ . Similarly for  $dy$ . Hence

$$df = f_x \dot{x}dt + f_y \dot{y}dt + f_t dt \quad (10)$$

now you can calculate partial derivatives of  $f$ , i.e.,  $f_x$ ,  $f_y$  and  $f_z$ .

$$f_x = 2xyt + y^2t \quad ; \quad f_y = x^2t + 2xyt + t^2 \quad ; \quad f_t = x^2y + xy^2 + 2yt \quad (7)$$

- ▶ Now we can replace  $f_x$ ,  $f_y$  and  $f_z$  in Eq.(6) to find the expression for  $\frac{df}{dt}$ .

- ▶ One can also compute  $\frac{d}{dt}f_x = \dot{f}_x = \frac{d}{dt}\left(\frac{\partial f}{\partial x}\right)$  as

$$\dot{f}_x = 2(\dot{x}yt + x\dot{y}t + xy) + 2y\dot{y}t + y^2 \quad (8)$$

- ▶ If we would like to find the differential element of the multi-variable function, we implement the *chain rule*. For function  $f(x(t), y(t), t)$ .

$$df = \frac{\partial f}{\partial x}dx + \frac{\partial f}{\partial y}dy + \frac{\partial f}{\partial t}dt = f_x dx + f_y dy + f_t dt \quad (9)$$

- ▶ As  $dx = \frac{dx}{dt}dt$ . Similarly for  $dy$ . Hence

$$df = f_x \dot{x}dt + f_y \dot{y}dt + f_t dt \quad (10)$$

If you divide by  $dt$ , you can recover the Eq.(6).

- Now let us look at an example of double partial derivative.

Consider a function  $f(x, y) = x^2y + xy^2$ . Calculate  $\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right)$ ,  $\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right)$  and  $\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y} \right)$ ?

► Now let us look at an example of double partial derivative.

Consider a function  $f(x, y) = x^2y + xy^2$ . Calculate  $\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right)$ ,  $\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right)$  and  $\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y} \right)$ ?

$$f_x = \frac{\partial f}{\partial x} = 2xy + y^2 \quad ; \quad f_y = \frac{\partial f}{\partial y} = x^2 + 2xy \quad (11)$$

► Now let us look at an example of double partial derivative.

Consider a function  $f(x, y) = x^2y + xy^2$ . Calculate  $\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right)$ ,  $\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right)$  and  $\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y} \right)$ ?

$$f_x = \frac{\partial f}{\partial x} = 2xy + y^2 \quad ; \quad f_y = \frac{\partial f}{\partial y} = x^2 + 2xy \quad (11)$$

Now another derivative is

$$f_{xx} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right) = 2y \quad (12)$$

- Now let us look at an example of double partial derivative.

Consider a function  $f(x, y) = x^2y + xy^2$ . Calculate  $\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right)$ ,  $\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right)$  and  $\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y} \right)$ ?

$$f_x = \frac{\partial f}{\partial x} = 2xy + y^2 \quad ; \quad f_y = \frac{\partial f}{\partial y} = x^2 + 2xy \quad (11)$$

Now another derivative is

$$f_{xx} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right) = 2y \quad (12)$$

Similarly, find the expression for  $\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right)$ .

► Now let us look at an example of double partial derivative.

Consider a function  $f(x, y) = x^2y + xy^2$ . Calculate  $\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right)$ ,  $\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right)$  and  $\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y} \right)$ ?

$$f_x = \frac{\partial f}{\partial x} = 2xy + y^2 \quad ; \quad f_y = \frac{\partial f}{\partial y} = x^2 + 2xy \quad (11)$$

Now another derivative is

$$f_{xx} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right) = 2y \quad (12)$$

Similarly, find the expression for  $\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right)$ .

$$f_{yy} = \frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right) = 2x \quad (13)$$

► Now let us look at an example of double partial derivative.

Consider a function  $f(x, y) = x^2y + xy^2$ . Calculate  $\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right)$ ,  $\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right)$  and  $\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y} \right)$ ?

$$f_x = \frac{\partial f}{\partial x} = 2xy + y^2 \quad ; \quad f_y = \frac{\partial f}{\partial y} = x^2 + 2xy \quad (11)$$

Now another derivative is

$$f_{xx} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right) = 2y \quad (12)$$

Similarly, find the expression for  $\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right)$ .

$$f_{yy} = \frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right) = 2x \quad (13)$$

Further,

$$f_{xy} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y} \right) = 2(x + y) \quad (14)$$



► Now let us look at an example of double partial derivative.

Consider a function  $f(x, y) = x^2y + xy^2$ . Calculate  $\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right)$ ,  $\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right)$  and  $\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y} \right)$ ?

$$f_x = \frac{\partial f}{\partial x} = 2xy + y^2 \quad ; \quad f_y = \frac{\partial f}{\partial y} = x^2 + 2xy \quad (11)$$

Now another derivative is

$$f_{xx} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right) = 2y \quad (12)$$

Similarly, find the expression for  $\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right)$ .

$$f_{yy} = \frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right) = 2x \quad (13)$$

Further,

$$f_{xy} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y} \right) = 2(x + y) \quad (14)$$

Calculate  $f_{yx} = \frac{\partial}{\partial y} \left( \frac{\partial f}{\partial x} \right)$  and check if they are equal?

# Plane Polar Coordinates

We can opt any coordinate system to solve a problem, but suitable and proper choice of coordinate system can vastly simplify the problem.

# Plane Polar Coordinates


We can opt any coordinate system to solve a problem, but suitable and proper choice of coordinate system can vastly simplify the problem.

- ▶ If a point on a plane is assigned  $(x, y)$  Cartesian coordinates and  $(r, \theta)$  polar coordinates, one can write down the coordinate transformations.

# Plane Polar Coordinates

We can opt any coordinate system to solve a problem, but suitable and proper choice of coordinate system can vastly simplify the problem.


- If a point on a plane is assigned  $(x, y)$  Cartesian coordinates and  $(r, \theta)$  polar coordinates, one can write down the coordinate transformations.



# Plane Polar Coordinates

We can opt any coordinate system to solve a problem, but suitable and proper choice of coordinate system can vastly simplify the problem.

- If a point on a plane is assigned  $(x, y)$  Cartesian coordinates and  $(r, \theta)$  polar coordinates, one can write down the coordinate transformations.




In  $\Delta OPQ$ ,  $\cos \theta = \frac{OP}{OQ} = \frac{x}{r} \Rightarrow x = r \cos \theta$

# Plane Polar Coordinates

We can opt any coordinate system to solve a problem, but suitable and proper choice of coordinate system can vastly simplify the problem.

- If a point on a plane is assigned  $(x, y)$  Cartesian coordinates and  $(r, \theta)$  polar coordinates, one can write down the coordinate transformations.




In  $\Delta OPQ$ ,  $\cos \theta = \frac{OP}{OQ} = \frac{x}{r} \Rightarrow x = r \cos \theta$

Similarly, from  $\Delta ORQ$ , we find  $y = r \sin \theta$ .

# Plane Polar Coordinates

We can opt any coordinate system to solve a problem, but suitable and proper choice of coordinate system can vastly simplify the problem.

- If a point on a plane is assigned  $(x, y)$  Cartesian coordinates and  $(r, \theta)$  polar coordinates, one can write down the coordinate transformations.



In  $\Delta OPQ$ ,  $\cos \theta = \frac{OP}{OQ} = \frac{x}{r} \Rightarrow x = r \cos \theta$

Similarly, from  $\Delta ORQ$ , we find  $y = r \sin \theta$ .


Hence, we have

$$x = r \cos \theta \quad ; \quad y = r \sin \theta \quad (15)$$

# Plane Polar Coordinates

We can opt any coordinate system to solve a problem, but suitable and proper choice of coordinate system can vastly simplify the problem.

- If a point on a plane is assigned  $(x, y)$  Cartesian coordinates and  $(r, \theta)$  polar coordinates, one can write down the coordinate transformations.



In  $\Delta OPQ$ ,  $\cos \theta = \frac{OP}{OQ} = \frac{x}{r} \Rightarrow x = r \cos \theta$

Similarly, from  $\Delta ORQ$ , we find  $y = r \sin \theta$ .

Hence, we have

$$x = r \cos \theta \quad ; \quad y = r \sin \theta \quad (15)$$

- The Position vector is:  $\vec{r} = x\hat{i} + y\hat{j} = r(\cos \theta \hat{i} + \sin \theta \hat{j})$




# Plane Polar Coordinates

## Velocity construction:

- From the figure, we can write down the unit vectors for  $\hat{r}$  and  $\hat{\theta}$  in the direction of increasing  $\theta$  and  $r$  respectively.

$$\hat{r} = \frac{\vec{r}}{|\vec{r}|} = \cos \theta \hat{i} + \sin \theta \hat{j} \quad (16)$$

$$\hat{\theta} = \cos \theta \hat{j} - \sin \theta \hat{i} \quad (17)$$




# Plane Polar Coordinates

## Velocity construction:

- From the figure, we can write down the unit vectors for  $\hat{r}$  and  $\hat{\theta}$  in the direction of increasing  $\theta$  and  $r$  respectively.

$$\hat{r} = \frac{\vec{r}}{|\vec{r}|} = \cos \theta \hat{i} + \sin \theta \hat{j} \quad (16)$$

$$\hat{\theta} = \cos \theta \hat{j} - \sin \theta \hat{i} \quad (17)$$



- As the unit vectors are not constant, they are changing with respect to time. Hence they can be finitely differentiated.


# Plane Polar Coordinates

## Velocity construction:

- From the figure, we can write down the unit vectors for  $\hat{r}$  and  $\hat{\theta}$  in the direction of increasing  $\theta$  and  $r$  respectively.

$$\hat{r} = \frac{\vec{r}}{|\vec{r}|} = \cos \theta \hat{i} + \sin \theta \hat{j} \quad (16)$$

$$\hat{\theta} = \cos \theta \hat{j} - \sin \theta \hat{i} \quad (17)$$



- As the unit vectors are not constant, they are changing with respect to time. Hence they can be finitely differentiated.

$$\frac{d\hat{r}}{dt} = \dot{\hat{r}} = \frac{d\hat{r}}{d\theta} \frac{d\theta}{dt} = \hat{\theta} \dot{\theta} \quad (18)$$

# Plane Polar Coordinates

- Similarly one can also calculate

$$\frac{d\hat{\theta}}{dt} = \dot{\hat{\theta}} = \frac{d\hat{\theta}}{d\theta} \frac{d\theta}{dt} = -\dot{\theta} \hat{r} \quad (19)$$

# Plane Polar Coordinates

- ▶ Similarly one can also calculate

$$\frac{d\hat{\theta}}{dt} = \dot{\hat{\theta}} = \frac{d\hat{\theta}}{d\theta} \frac{d\theta}{dt} = -\dot{\theta} \hat{r} \quad (19)$$

Now one can construct the velocity vector.

# Plane Polar Coordinates

- Similarly one can also calculate

$$\frac{d\hat{\theta}}{dt} = \dot{\hat{\theta}} = \frac{d\hat{\theta}}{d\theta} \frac{d\theta}{dt} = -\dot{\theta} \hat{r} \quad (19)$$

Now one can construct the velocity vector.

$$\vec{v} = \frac{d\vec{r}}{dt} = \frac{d}{dt}(r\hat{r}) = \frac{dr}{dt}\hat{r} + r\frac{d\hat{r}}{dt} \quad (20)$$

$$\vec{v} = \dot{r}\hat{r} + r\dot{\theta}\hat{\theta} \quad (21)$$

# Plane polar coordinate

Now further, one can construct acceleration.

## Plane polar coordinate

Now further, one can construct acceleration.

$$\vec{a} = \frac{d\vec{v}}{dt} = \frac{d}{dt}(\dot{r}\hat{r} + r\dot{\theta}\hat{\theta}) \quad (22)$$

$$= (\ddot{r} - r\dot{\theta}^2)\hat{r} + (r\ddot{\theta} + 2\dot{r}\dot{\theta})\hat{\theta} \quad (23)$$



# Plane polar coordinate

Now further, one can construct acceleration.


$$\vec{a} = \frac{d\vec{v}}{dt} = \frac{d}{dt}(\dot{r}\hat{r} + r\dot{\theta}\hat{\theta}) \quad (22)$$

$$= (\ddot{r} - r\dot{\theta}^2)\hat{r} + (r\ddot{\theta} + 2\dot{r}\dot{\theta})\hat{\theta} \quad (23)$$

The term  $\ddot{r}$  is *linear acceleration* in radial direction,  $r\dot{\theta}^2$  is the *centripetal acceleration*,  $\ddot{\theta}$  is the *acceleration* in the tangential direction, and  $2\dot{r}\dot{\theta}$  is the *Coriolis acceleration*.


## Example

**Acceleration construction:** Consider an object  $P$  moving on a circular path with a uniform velocity and radius  $R$ . Prove that it'll always be attracted towards the center of the circle.



## Example


**Acceleration construction:** Consider an object  $P$  moving on a circular path with a uniform velocity and radius  $R$ . Prove that it'll always be attracted towards the center of the circle.



The total acceleration can be written as:  $\vec{a} = a_r \hat{r} + a_\theta \hat{\theta}$   
where,  $a_r = \ddot{r} - r\dot{\theta}^2$  and  $a_\theta = r\ddot{\theta} + 2\dot{r}\dot{\theta}$ .

## Example

**Acceleration construction:** Consider an object  $P$  moving on a circular path with a uniform velocity and radius  $R$ . Prove that it'll always be attracted towards the center of the circle.




The total acceleration can be written as:  $\vec{a} = a_r \hat{r} + a_\theta \hat{\theta}$   
where,  $a_r = \ddot{r} - r\dot{\theta}^2$  and  $a_\theta = r\ddot{\theta} + 2\dot{r}\dot{\theta}$ .

Since the object is exhibiting circular motion with  $r = R$ . Hence,  $\dot{r} = \ddot{r} = 0$ . This implicates,  $a_r = -R\dot{\theta}^2$  ;  $a_\theta = R\ddot{\theta}$

## Example

**Acceleration construction:** Consider an object  $P$  moving on a circular path with a uniform velocity and radius  $R$ . Prove that it'll always be attracted towards the center of the circle.




The total acceleration can be written as:  $\vec{a} = a_r \hat{r} + a_\theta \hat{\theta}$

where,  $a_r = \ddot{r} - r\dot{\theta}^2$  and  $a_\theta = r\ddot{\theta} + 2\dot{r}\dot{\theta}$ .

Since the object is exhibiting circular motion with  $r = R$ . Hence,  $\dot{r} = \ddot{r} = 0$ . This implicates,  $a_r = -R\dot{\theta}^2$  ;  $a_\theta = R\ddot{\theta}$  For a uniform circular motion,  $\dot{\theta} = \omega$ . Thus  $a_\theta = 0$ . Wherease, for non-uniform circular motion,  $a_\theta = R\frac{d\omega}{dt} = r\alpha$  and  $a_r = r\omega^2$ .

## Example

**Acceleration construction:** Consider an object  $P$  moving on a circular path with a uniform velocity and radius  $R$ . Prove that it'll always be attracted towards the center of the circle.



The total acceleration can be written as:  $\vec{a} = a_r \hat{r} + a_\theta \hat{\theta}$   
where,  $a_r = \ddot{r} - r\dot{\theta}^2$  and  $a_\theta = r\ddot{\theta} + 2\dot{r}\dot{\theta}$ .

Since the object is exhibiting circular motion with  $r = R$ . Hence,  $\dot{r} = \ddot{r} = 0$ . This implicates,  $a_r = -R\dot{\theta}^2$  ;  $a_\theta = R\ddot{\theta}$  For a uniform circular motion,  $\dot{\theta} = \omega$ . Thus  $a_\theta = 0$ . Wherease, for non-uniform circular motion,  $a_\theta = R\frac{d\omega}{dt} = r\alpha$  and  $a_r = r\omega^2$ .

- ▶ As the magnitude of velocity is constant, but due to change in the direction of velocity, it changes the direction and producing non-zero acceleration towards the center.


# Cylindrical Coordinates

Now we would try to see the coordinate transformation from Cartesian to cylindrical coordinates, velocity and acceleration.

# Cylindrical Coordinates

Now we would try to see the coordinate transformation from Cartesian to cylindrical coordinates, velocity and acceleration.

Let us consider a point  $P$  on a cylinder with Cartesian coordinates  $(x, y, z)$  and cylindrical coordinates  $(\rho, \phi, z)$ .






# Cylindrical Coordinates

Now we would try to see the coordinate transformation from Cartesian to cylindrical coordinates, velocity and acceleration.

Let us consider a point  $P$  on a cylinder with Cartesian coordinates  $(x, y, z)$  and cylindrical coordinates  $(\rho, \phi, z)$ .



- The position vector for the point  $P$  is

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k} \quad (24)$$

## Cylindrical coordinates

- Now, we can write down the transformations as

$$x = \rho \cos \phi \quad ; \quad y = \rho \sin \phi \quad ; \quad z = z \quad (25)$$

## Cylindrical coordinates

- Now, we can write down the transformations as

$$x = \rho \cos \phi \quad ; \quad y = \rho \sin \phi \quad ; \quad z = z \quad (25)$$

- Now, we can express the unit vectors for cylindrical coordinates.

## Cylindrical coordinates

- Now, we can write down the transformations as

$$x = \rho \cos \phi \quad ; \quad y = \rho \sin \phi \quad ; \quad z = z \quad (25)$$

- Now, we can express the unit vectors for cylindrical coordinates.

$$\hat{\rho} = \frac{\vec{\rho}}{\rho} = \frac{x\hat{i} + y\hat{j}}{\rho} = \cos \phi \hat{i} + \sin \phi \hat{j} \quad (26)$$

$$\hat{\phi} = -\sin \phi \hat{i} + \cos \phi \hat{j} \quad (27)$$

$$\hat{z} = \hat{z} \quad (28)$$

## Cylindrical coordinates

- Now, we can write down the transformations as

$$x = \rho \cos \phi \quad ; \quad y = \rho \sin \phi \quad ; \quad z = z \quad (25)$$

- Now, we can express the unit vectors for cylindrical coordinates.

$$\hat{\rho} = \frac{\vec{\rho}}{\rho} = \frac{x\hat{i} + y\hat{j}}{\rho} = \cos \phi \hat{i} + \sin \phi \hat{j} \quad (26)$$

$$\hat{\phi} = -\sin \phi \hat{i} + \cos \phi \hat{j} \quad (27)$$

$$\hat{z} = \hat{z} \quad (28)$$

- Construct the **velocity** and **acceleration**.

## Cylindrical coordinates

- Now, we can write down the transformations as

$$x = \rho \cos \phi \quad ; \quad y = \rho \sin \phi \quad ; \quad z = z \quad (25)$$

- Now, we can express the unit vectors for cylindrical coordinates.

$$\hat{\rho} = \frac{\vec{\rho}}{\rho} = \frac{x\hat{i} + y\hat{j}}{\rho} = \cos \phi \hat{i} + \sin \phi \hat{j} \quad (26)$$

$$\hat{\phi} = -\sin \phi \hat{i} + \cos \phi \hat{j} \quad (27)$$

$$\hat{z} = \hat{z} \quad (28)$$

- Construct the **velocity** and **acceleration**.

$$\vec{v} = \dot{\vec{r}} = \dot{\rho}\hat{\rho} + \rho\dot{\phi}\hat{\phi} + \dot{z}\hat{z} \quad (29)$$



$$\vec{a} = \dot{\vec{v}} = \left(\ddot{\rho} - \rho\dot{\phi}^2\right)\hat{\rho} + \left(\rho\ddot{\phi} + 2\dot{\rho}\dot{\phi}\right)\hat{\phi} + \ddot{z}\hat{z} \quad (30)$$

# Spherical Polar Coordinates

- ▶ Now we can have overview of Spherical polar coordinates.

# Spherical Polar Coordinates



- Now we can have overview of Spherical polar coordinates.





# Spherical Polar Coordinates



- Now we can have overview of Spherical polar coordinates.



$$\text{In } \triangle OPC, \quad \cos \theta = \frac{OP}{OC} = \frac{r}{z} \Rightarrow z = r \cos \theta$$

# Spherical Polar Coordinates

- Now we can have overview of Spherical polar coordinates.





$$\text{In } \triangle OPC, \quad \cos \theta = \frac{OP}{OC} = \frac{r}{z} \Rightarrow z = r \cos \theta$$

$$\text{From } \triangle PDO, \quad \cos(90 - \theta) = \sin \theta = \frac{OD}{OP} = \frac{OD}{r} \Rightarrow OD = r \sin \theta$$

# Spherical Polar Coordinates

- Now we can have overview of Spherical polar coordinates.





$$\text{In } \triangle OPC, \quad \cos \theta = \frac{OP}{OC} = \frac{r}{z} \Rightarrow z = r \cos \theta$$

$$\text{From } \triangle PDO, \quad \cos(90 - \theta) = \sin \theta = \frac{OD}{OP} = \frac{OD}{r} \Rightarrow OD = r \sin \theta$$

$$\text{Now from } \triangle OAD, \quad \cos \phi = \frac{OA}{OD} = \frac{x}{r \sin \theta} \Rightarrow x = r \sin \theta \cos \phi$$

# Spherical Polar Coordinates

- Now we can have overview of Spherical polar coordinates.



In  $\triangle OPC$ ,  $\cos \theta = \frac{OP}{OC} = \frac{r}{z} \Rightarrow z = r \cos \theta$



From  $\triangle PDO$ ,  $\cos(90 - \theta) = \sin \theta = \frac{OD}{OP} = \frac{OD}{r} \Rightarrow OD = r \sin \theta$

Now from  $\triangle OAD$ ,  $\cos \phi = \frac{OA}{OD} = \frac{x}{r \sin \theta} \Rightarrow x = r \sin \theta \cos \phi$

Similarly, from  $\triangle OBD$  we can find,  $y = r \sin \theta \sin \phi$ .

# Spherical Polar Coordinates

- Now we can have overview of Spherical polar coordinates.



In  $\Delta OPC$ ,  $\cos \theta = \frac{OP}{OC} = \frac{r}{z} \Rightarrow z = r \cos \theta$

From  $\Delta PDO$ ,  $\cos(90 - \theta) = \sin \theta = \frac{OD}{OP} = \frac{OD}{r} \Rightarrow OD = r \sin \theta$



Now from  $\Delta OAD$ ,  $\cos \phi = \frac{OA}{OD} = \frac{x}{r \sin \theta} \Rightarrow x = r \sin \theta \cos \phi$

Similarly, from  $\Delta OBD$  we can find,  $y = r \sin \theta \sin \phi$ .

Thus,  $x = r \sin \theta \cos \phi$  ;  $y = r \sin \theta \sin \phi$  ;  $z = r \cos \theta$

# Spherical Polar Coordinates

- Now we can have overview of Spherical polar coordinates.



In  $\Delta OPC$ ,  $\cos \theta = \frac{OP}{OC} = \frac{r}{z} \Rightarrow z = r \cos \theta$

From  $\Delta PDO$ ,  $\cos(90 - \theta) = \sin \theta = \frac{OD}{OP} = \frac{OD}{r} \Rightarrow OD = r \sin \theta$

Now from  $\Delta OAD$ ,  $\cos \phi = \frac{OA}{OD} = \frac{x}{r \sin \theta} \Rightarrow x = r \sin \theta \cos \phi$

Similarly, from  $\Delta OBD$  we can find,  $y = r \sin \theta \sin \phi$ .

Thus,  $x = r \sin \theta \cos \phi$  ;  $y = r \sin \theta \sin \phi$  ;  $z = r \cos \theta$

- Find the expression for velocity & acceleration in spherical polar coordinate.