

Funktionalanalysis Hausaufgaben Blatt 2

Jun Wei Tan*

Julius-Maximilians-Universität Würzburg

(Dated: November 5, 2024)

Problem 1. For $p \in [1, \infty]$ we define the set

$$\ell^p := \begin{cases} \{x := (x_n)_{n \in \mathbb{N}} \subset \mathbb{K} \mid \|x\|_p := (\sum_{n=1}^{\infty} |x_n|^p)^{\frac{1}{p}} < \infty\} & p < \infty \\ \{x := (x_n)_{n \in \mathbb{N}} \subset \mathbb{K} \mid \|x\|_{\infty} := \sup_{n \in \mathbb{N}} |x_n| < \infty\} & p = \infty. \end{cases}$$

Show that the usual operations on sequences induce a vector space structure on ℓ^p . Moreover, show that ℓ^p is a subspace of ℓ^r for $p \leq r$.

Problem 2. In this exercise, we consider the spaces ℓ^p for $p \in (1, \infty)$. Note that for every such p there exists a conjugate number $q \in (1, \infty)$ which satisfies $\frac{1}{p} + \frac{1}{q} = 1$.

- (a) Show that the product of two non-negative real numbers $a, b \in [0, \infty)$ satisfies Young's inequality, that is

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}.$$

Hint: Use the AM-GM inequality.

- (b) Prove that Hölder's inequality

$$\|xy\|_1 := \sum_{n=1}^{\infty} |x_n y_n| \leq \|x\|_p \|y\|_q$$

holds true for any two sequences $x \in \ell^p$ and $y \in \ell^q$.

- (c) Show Minkowsky's inequality, that is

$$\|x + y\|_p \leq \|x\|_p + \|y\|_p$$

for $x, y \in \ell^p$.

* jun-wei.tan@stud-mail.uni-wuerzburg.de

- (d) Let $\lambda := (\lambda_n)_{n \in \mathbb{N}} \subset [0, 1]$ be a sequence in ℓ^1 with $\|\lambda\|_1 = 1$. Show that Jensen's inequality

$$f\left(\sum_{n=1}^{\infty} \lambda_n x_n\right) \leq \sum_{n=1}^{\infty} \lambda_n f(x_n)$$

holds true for every convex function $f \in \mathcal{C}(I)$ on an open interval $I \subseteq \mathbb{R}$ and every sequence $(x_n)_{n \in \mathbb{N}} \subset I$ such that $\sum_{n=1}^{\infty} \lambda_n x_n$ and $\sum_{n=1}^{\infty} \lambda_n f(x_n)$ converge and $\sum_{n=1}^{\infty} \lambda_n x_n \in I$. Conclude that $\|x\|_r \leq \|x\|_p$ for every $x \in \ell^p$ and $p \leq r$.

Problem 3. Let $p \in [1, \infty)$. Consider the sequences $(e_n := (\delta_{nm})_{m \in \mathbb{N}})_{n \in \mathbb{N}} \subset \ell^p$. Show that for every sequence $x = (x_n)_{n \in \mathbb{N}} \in \ell^p$ the series $\sum_{n \in \mathbb{N}} x_n e_n$ converges unconditionally towards x with respect to $\|x\|_p$. Does it converge absolutely? Moreover, show that a sequence $x = (x_n)_{n \in \mathbb{N}}$ lies in ℓ^p if the series $\sum_{n \in \mathbb{N}} x_n e_n$ converges unconditionally with respect to $\|\cdot\|_p$.

Hint: Having Minkowski's inequality, you can use that $(\ell^p, \|\cdot\|_p)$ is a normed space without proof

Problem 4. In the upcoming exercise sheets, we will prove the Stone-Weierstraß theorem in several steps. Here, we do some necessary preparation we will need for the actual proof.

By recursion, define the polynomials

$$p_0(x) = 0, \quad \text{and} \quad p_{n+1}(x) = p_n(x) + \frac{1}{2}(x - p_n^2(x)).$$

- (a) Show $p_n(0) = 0$ and the estimates

$$p_n(x) \geq 0, \quad \text{and} \quad 0 \leq \sqrt{x} - p_n(x) \leq \frac{2\sqrt{x}}{2 + n\sqrt{x}}$$

for $x \in [0, 1]$.

Hint: First show the coarser estimates $0 \leq p_n(x) \leq 1$ for $x \in [0, 1]$ by induction. Use this in a second induction to improve the estimates.

- (b) Conclude that $(p_n)_{n \in \mathbb{N}}$ converges uniformly to the square root function on the interval $[0, 1]$.
- (c) Let $\alpha > 0$. Construct a sequence of polynomials that converges uniformly to the square root function on $[0, \alpha]$.
- (d) Let $\alpha > 0$. Construct a sequence of polynomials that converges uniformly to the absolute value function on $[-\alpha, \alpha]$.