

Notes for the course in
FUNCTIONAL ANALYSIS

Held in WS24/25

At the JMU Würzburg
By Prof. Dr. Stefan Waldmann

By

Jun Wei Tan

Julius-Maximilians-Universität Würzburg

jun-wei.tan@stud-mail.uni-wuerzburg.de

Contents

1	Topology	3
1.1	Basic Concepts	3
1.2	Baire Spaces	5
2	Topological Vector Spaces	8
2.1	Continuity	10
2.2	Subspaces	11
3	Banach Spaces	12
3.1	Introductory Definitions	12
3.1.1	Norms & Seminorms	12
3.1.2	Bases	14
3.1.3	Examples	15
3.2	The Hahn-Banach Theorems	16
3.2.1	Sublinear Functionals	16
3.2.2	Geometric Hahn-Banach	19
3.3	Duality	22
3.3.1	The Weak Topology	22
3.3.2	Polars & Banach-Alaoglu Theorems	24
3.3.3	Separability & Reflexivity	28
3.4	Banach-Steinhaus	30
3.5	Open Mapping & Closed Graph Theorems	32
4	Hilbert Spaces	36
4.1	Introduction	36
4.2	Lattices	41
4.3	Bases	42
4.4	Applications	42

Topology

1.1 Basic Concepts

Theorem 1.1 (List of Useful Identities).

- | | |
|-----|----------------------------------------------------------------------|
| (a) | $(A \cup B)^{\text{cl}} = A^{\text{cl}} \cup B^{\text{cl}}.$ |
| (b) | $(A \cup B)^{\circ} \supseteq A^{\circ} \cup B^{\circ}.$ |
| (c) | $(A \cap B)^{\text{cl}} \subseteq A^{\text{cl}} \cap B^{\text{cl}}.$ |
| (d) | $(A \cap B)^{\circ} = A^{\circ} \cap B^{\circ}.$ |
| (e) | $(M \setminus A)^{\text{cl}} = M \setminus A^{\circ}.$ |
| (f) | $(M \setminus A)^{\circ} = M \setminus A^{\text{cl}}.$ |

Proof. (a) If all neighbourhoods of x intersect A , then they must certainly intersect $A \cup B$. The same thing happens if all neighbourhoods intersect B . Conversely, we suppose that x is not in A^{cl} or B^{cl} . Then we have an open neighbourhood not intersecting A , and an open neighbourhood not intersecting B . Considering the intersection of these two neighbourhoods shows that x is not in the closure of $A \cup B$ either.

(b) $A^{\circ} \cup B^{\circ}$ is an open set contained in $A \cup B$.

(c) Suppose every neighbourhood of x intersects $A \cap B$. Then every neighbourhood intersects A , and also intersects B .

(d) Clearly, $A^{\circ} \cap B^{\circ}$ is an open set contained in $A \cap B$. Conversely, it is also true that $(A \cap B)^{\circ}$ is an open set contained in A , and thus its interior, and it is also contained in B .

(e) Clearly, $M \setminus A^\circ$ is a closed set containing $M \setminus A$. This shows one inclusion.

Now suppose $x \in M \setminus A^\circ$. Since it is not in the interior, no open neighbourhood of x is completely contained in A ; in particular, every neighbourhood must intersect $M \setminus A$. This shows the reverse inclusion.

(f) Clearly, $M \setminus A^{\text{cl}}$ is an open subset of $M \setminus A$.

Conversely, suppose x is an element of $(M \setminus A)^\circ$. Then there is an open neighbourhood of x contained in $M \setminus A$, in particular not intersecting A . This shows that x is in $M \setminus A^{\text{cl}}$. \square

Definition 1.2 (Preorder). A relation \preceq on a set A is a preorder if the following conditions hold:

- (a) $\alpha \preceq \alpha$ for all $\alpha \in A$
- (b) If $\alpha \preceq \beta$ and $\beta \preceq \gamma$, then $\alpha \preceq \gamma$.

Definition 1.3. A *directed set* A is a set with a preorder and the following additional condition: For all $\alpha, \beta \in A$, there is some γ with $\alpha \preceq \gamma$ and $\beta \preceq \gamma$.

Remark 1.4. The additional condition $\alpha \preceq \beta$ and $\beta \preceq \alpha \implies \beta = \alpha$ is known as a *partial order*. This is sometimes too restrictive, and is usually not important. The important property is the defining property of a directed set.

Definition 1.5. A map from a directed set I into a set X is called a *net* (indexed by I). The elements of this net are denoted by $(x_i)_{i \in I}$.

Definition 1.6 (Convergence). Let X now be a topological space. We say that a net $(x_i)_{i \in I}$ converges to x if for every neighbourhood U of x there is $\alpha \in I$ such that $\beta \succeq \alpha \implies x_\beta \in U$.

Theorem 1.7. The limit of a net is unique if the space X is Hausdorff.

Proof. The proof is analogous to that of a sequence. \square

It is obvious that all sequences are nets. The difference is as follows: A sequence only has countably many elements. Thus, it is possible that there will be too many open sets. This leads us to our next theorem:

Theorem 1.8. Let X be a topological space and $(x_i)_{i \in I}$ be a net converging to x . If X is first countable, then there is a sequence converging to x .

In general, we have the following theorem

Theorem 1.9. Let A be a subset of the topological space X . $x \in A^{\text{cl}}$ if and only if there is a net in A converging to x .

1.2 Baire Spaces

Definition 1.10 (Nowhere Dense). A subset A of a topological space is called nowhere dense if the interior of its closure is open, $(A^{\text{cl}})^{\circ} = \emptyset$.

Theorem 1.11. A subset A of a topological space (M, \mathcal{M}) is nowhere dense iff its complement contains a dense open set.

Proof. We perform the following computation:

$$\begin{aligned}
 A \text{ nowhere dense} &\iff (A^{\text{cl}})^{\circ} = \emptyset \\
 &\iff M \setminus (A^{\text{cl}})^{\circ} = M \\
 &\iff (M \setminus A^{\text{cl}})^{\text{cl}} = M \\
 &\iff ((M \setminus A)^{\circ})^{\text{cl}} = M \\
 &\iff (M \setminus A)^{\circ} \text{ is dense.} \quad \square
 \end{aligned}$$

Definition 1.12 (Meager). A subset is called meager if it is a countable union of nowhere dense sets.

Note that the entire topological space can be meager - for example, consider \mathbb{Q} with its usual topology. Then \mathbb{Q} is a countable union of singletons, which are themselves nowhere dense. Thus, \mathbb{Q} is meager in \mathbb{Q} .

Theorem 1.13. Let (M, \mathcal{M}) be a topological space. Then the following are equivalent:

- (a) Any countable union of closed subsets of M without inner points has no inner points.
- (b) Any countable intersection of open dense subsets of M is dense.
- (c) Every non-empty open subset of M is not meager
- (d) The complement of every meager subset of M is dense.

Proof. The proof follows (a) \implies (b) \implies (c) \implies (d) \implies (a).

1. Let $(U_i)_{i \in \mathbb{N}}$ be a collection of dense open subsets of M . We consider their complements, which all have empty interior. Since

$$M \setminus \left(\bigcap_{i=1}^{\infty} U_i \right) = \bigcup_{i=1}^{\infty} (M \setminus U_i)$$

and the sets in the union on the right are all closed subsets without interior points, their union has no interior points, hence the countable intersection is dense.

2. Suppose we had a meager nonempty open subset U of M , that is, we have $U = \bigcup_{i=1}^{\infty} A_i$ with A_i nowhere dense sets. Then $M \setminus A_i$ is a dense subset for all i , by (b), their intersection is still dense. Then

$$\begin{aligned} \emptyset &= M \setminus \left(\bigcap_{i=1}^{\infty} (M \setminus A_i^{\text{cl}}) \right)^{\text{cl}} \\ &= M \setminus \left(M \setminus \bigcup_{i=1}^{\infty} A_i^{\text{cl}} \right)^{\text{cl}} \\ &= M \setminus \left(M \setminus \left(\bigcup_{i=1}^{\infty} A_i^{\text{cl}} \right)^{\circ} \right) \\ &= \left(\bigcup_{i=1}^{\infty} A_i^{\text{cl}} \right)^{\circ}, \end{aligned}$$

which is a contradiction, since we had U as a nonempty open subset of the final union.

3. Suppose we have a meager subset A such that $M \setminus A$ is not dense. Then $(M \setminus A)^{\text{cl}} = M \setminus A^{\circ} \neq M$. Then A has nonempty interior, contradicting (c).
4. Finally, let us consider a sequence of closed sets $(A_n)_{n \in \mathbb{N}}$ without interior points, and we suppose that their union has a nonempty interior. Then

$$\left(M \setminus \bigcup_{n=1}^{\infty} A_n \right)^{\text{cl}} = M \setminus \left(\bigcup_{n=1}^{\infty} A_n \right)^{\circ} \neq M,$$

contradicting (d), since $\bigcup_{n=1}^{\infty} A_n$ is meager and hence has a dense complement. \square

Definition 1.14 (Baire Spaces). A topological space X is a Baire space if one (and hence all) of the conditions of Theorem 1.13 hold.

Note that if X is a nonempty Baire space, then it cannot be meager. For if X were a Baire space, we would have

$$X = \bigcup_{n=1}^{\infty} U_n$$

with all U_n s meager. Then $X = \bigcup_{n=1}^{\infty} \overline{U_n}$, which is a contradiction, since this is a union of closed subsets without interior points, which must have empty interior, but the interior of X is X , which is nonempty.

Theorem 1.13 is clearly unwieldy, and one must first show that a space is a Baire space before even beginning to apply this theorem. Thus, we seek easier conditions with which we can verify that a space is a Baire space. These theorems are known as Baire's Theorems.

Theorem 1.15 (Baire I). A complete metric space (M, d) is a Baire space.

Proof. We seek to show the condition in Theorem 1.13(b). Let us consider a collection of dense open sets O_n and a point $x \in M$, as well as an open neighbourhood U of x . Since U intersects O_1 , we can find an open ball $B_{\epsilon_1}(p_1)$ whose closure is contained in $U \cap O_1$. Inductively, we construct open balls $B_{\epsilon_n}(p_n)$ such that $B_{\epsilon_n}(p_n)^{\text{cl}} \subseteq B_{\epsilon_{n-1}}(p_{n-1})$. Then,

$$\bigcap_{i=1}^{\infty} B_{\epsilon_n}(p_n)^{\text{cl}} \subseteq U \cap \bigcap_{i=1}^{\infty} O_n$$

But since the metric space is complete, the intersection on the left side is nonempty. Thus, the intersection on the right is nonempty. Since U was an arbitrary open set, the intersection is dense. \square

Theorem 1.16 (Baire II). A locally compact Hausdorff space (M, \mathcal{M}) is a Baire space.

CHAPTER TWO

Topological Vector Spaces

Definition 2.1. A topological vector space is a vector space with a topology such that addition and scalar multiplication are continuous.

Theorem 2.2. Translation $T_v : x \mapsto x + v$ and multiplication $\lambda : x \mapsto \lambda x$ with $\lambda \neq 0$ are homeomorphisms.

Proof. They are invertible with continuous inverse T_{-v} and $\frac{1}{\lambda}$ respectively □

It is defined in some books that a topological vector space must be T_1 (or T_0). The usefulness of this definition comes from the fact that T_0 topological vector spaces are automatically T_2 and T_3 . This is, in fact, not a statement about topological vector spaces, but about topological groups. Before we prove the result, we will need the following lemmas

Lemma 2.3. Let G be a group that is also a T_1 topological space. G is a topological group iff the map $G \times G \rightarrow G, (x, y) \mapsto xy^{-1}$ is continuous.

Proof. Clearly, if G is a topological group, then the map is continuous.

Conversely, we consider $y \mapsto (e, y) \mapsto y^{-1}$, which is continuous as a composition of continuous functions. Then $(x, y) \mapsto (x, y^{-1}) \mapsto xy$ is continuous, again as a composition of continuous functions. □

Lemma 2.4. A topological space X is Hausdorff iff the diagonal $X \times X$ is closed.

Proof. Suppose the diagonal is closed. Then, given (x, y) , there is an open set Z and a basis element $U \times V$ such that $(x, y) \in U \times V \subseteq Z$. Since this does not intersect the diagonal, it follows that U and V must be disjoint.

Conversely, given x and y , we can find disjoint open sets U and V containing x and y , and $U \times V$ does not intersect the diagonal, since they are disjoint. □

Now, we move on to our first result about regularity. This is independent of the other separation axioms:

Theorem 2.5. Let G be a topological group. It is regular. In particular, for all points $x \in G$ and closed sets A not containing x , we have a neighbourhood V of e such that $V \cdot x$ and $V \cdot A$ are disjoint.

Proof. Firstly, we show that for all neighbourhoods U of e , there is a symmetric neighbourhood V such that $V \cdot V \subseteq U$ of e .

This follows because the map $(x, y) \mapsto x \cdot y$ is continuous, hence there is a neighbourhood V such that $V \cdot V \subseteq U$. Then because $(x, y) \mapsto x \cdot y^{-1}$ is continuous, there exists W with $W \cdot W^{-1} \subseteq V$. Then $W \cdot W^{-1}$ is the desired symmetric neighbourhood.

Now, since A is closed, we have $A \cdot x^{-1}$ closed (and not containing e), thus we choose an open symmetric neighbourhood V of e such that $V \cdot V \cap (A \cdot x^{-1}) = \emptyset$.

Now we show that $V \cdot x$ and $V \cdot A$ are disjoint. Suppose not. Then we have $v \cdot x = v' \cdot a$ for $v, v' \in V, a \in A$. However, this also means that $v'^{-1} \cdot v = a \cdot x^{-1}$, a contradiction. \square

As a corollary we have, by a simple change of notation:

Corollary 2.6. Let V be a topological vector space, $A \subseteq V$ closed and $v \in V$. Then there exists a symmetric neighbourhood U of 0 such that $(A + U) \cap (v + U) = \emptyset$.

We can also extend this to compact sets:

Corollary 2.7. Let V be a topological vector space, $A \subseteq V$ closed, $K \subseteq V$ compact. Then there exists a symmetric neighbourhood U of 0 such that $(A + U) \cap (K + U) = \emptyset$.

Proof. As usual, we apply the previous corollary to each point $x \in K$. Then we choose a finite subcover of these, and take their intersections. \square

Now, we move on to the more important result:

Theorem 2.8. Let G be a T_0 topological group. Then it is T_1, T_2 and T_3 .

Proof. We will prove $T_0 \implies T_1 \implies T_2$.

The goal is: We show that $\{e\}$ is closed. Pick $g \neq e$. Because G is T_0 , either $G \setminus \{e\}$ is a neighbourhood of g , or $G \setminus \{g\}$ is a neighbourhood of e . Suppose the latter. Then applying the translation $T_{g^{-1}}$ shows that $G \setminus \{e\}$ is a neighbourhood of g^{-1} , and applying the inversion map shows that $G \setminus \{e\}$ is a neighbourhood of g . Thus $\{e\}$ is closed.

Then the translations show that every singleton set is closed.

Note: If the group is abelian, then the proof is significantly easier: We suppose U is a neighbourhood of x not containing y . Then $x + y - U$ is a neighbourhood of y not containing x .

Now let the identity be closed. The diagonal of $G \times G$ is the preimage of $\{e\}$ under the map $(x, y) \mapsto xy^{-1}$, hence the diagonal is closed too. This shows that G is Hausdorff.

Since all topological groups are regular, this shows T_3 . \square

2.1 Continuity

Definition 2.9 (Uniform Continuity). Let V, W be topological vector spaces and $\phi : V \rightarrow W$ be a map. Then ϕ is called uniformly continuous if for all neighbourhoods $U \subseteq W$ of 0 there exists a neighbourhood $Z \subseteq V$ such that $v - v' \in Z \implies \phi(v) - \phi(v') \in U$.

Theorem 2.10 (Equivalence of Continuity Conditions). Let V and W be topological vector spaces and let $\phi : V \rightarrow W$ be a linear map. Then the following are equivalent

- (a) The map ϕ is continuous at 0 .
- (b) The map ϕ is continuous at some $v \in V$.
- (c) The map ϕ is continuous.
- (d) The map ϕ is uniformly continuous.

Proof. 1. Assume ϕ is continuous at 0 .

Recall: This means that for all neighbourhoods U of 0_W , there is a neighbourhood Z of 0_V such that $\phi(Z) \subseteq U$. Now choose $v \in V$ and a neighbourhood $U \ni \phi(v)$. Then $T_{-v}(U)$ is a neighbourhood of 0_W , and we choose a neighbourhood Z of 0 such that $\phi(Z) \subseteq T_{-v}(U)$ by continuity at 0 . Then $T_v(Z) \subseteq U$ and contains v , as desired.

- 2. Repeating the same proof, we can get continuity at every point in V , which is equivalent to continuity.
- 3. We actually only need continuity at 0 . Given an open neighbourhood Z of 0_W , we choose an open neighbourhood U of 0_V that maps into Z . Then we have, for $v - u \in Z$, that $\phi(v) - \phi(u) = \phi(v - u) \in W$.
- 4. Uniform continuity directly implies continuity at 0 if we take v' to be 0 in the definition. \square

Definition 2.11 (Algebraic and Topological Dual). The algebraic dual of a vector space V is the well known space of linear maps from V to the field over which it is defined. We denote this by $V^* = \text{Hom}(V, \mathbb{K})$. The topological dual $V' = L(V, \mathbb{K})$ is the space of all *continuous* linear maps. Clearly, $L(V, \mathbb{K}) \subseteq \text{Hom}(V, \mathbb{K})$.

Note that in a complex vector space, all linear functionals are uniquely defined by their real part:

Theorem 2.12. Let f be a linear functional such that g is its real part. Then

$$f(v) = g(v) - ig(iv).$$

Proof. Define the imaginary part of the functional to be h , such that $f = g + ih$. Then, $f(iv) = g(iv) + ih(iv)$. On the other hand, by linearity, $f(iv) = if(v) = ig(v) - h(v)$. Comparing real parts, we have

$$g(iv) = -h(v)$$

or

$$f(v) = g(v) - ig(v)$$

as desired. \square

2.2 Subspaces

Theorem 2.13. Let V be a topological vector space. If $W \subseteq V$ is a subspace, then its closure W^{cl} is also a subspace of V .

Proof. Suppose $v, w \in W^{\text{cl}}$. Then there are nets $(v_i)_{i \in I}$ converging to v and $(w_i)_{i \in I}$ converging to w . Then $(v_i + w_i)_{i \in I}$ is a net converging to $v + w$, hence $v + w$ is still in W^{cl} . \square

Theorem 2.14. All proper subspaces S of a topological vector space V have empty interior.

Proof. The proof follows in two steps: First, we translate an open set contained in S to the origin. Then, since all neighbourhoods of the origin are absorbing, $S = V$ by scaling. \square

Banach Spaces

3.1 Introductory Definitions

3.1.1 Norms & Seminorms

Definition 3.1 (Norm). A norm $\|\cdot\|$ on a vector space V is a function $V \rightarrow \mathbb{R}$ that satisfies

- (a) (Homogeneity) $\|sv\| = |s| \cdot \|v\|$ for all vectors v and scalars s
- (b) (Triangle Inequality) $\|u + v\| \leq \|u\| + \|v\|$
- (c) (Positivity) $\|v\| > 0$ for all $v \neq 0$.

Remark 3.2. A norm induces a topology through the metric defined by $d(v, w) = \|v - w\|$. If a vector space has this topology, it is known as a *normed space*.

Definition 3.3 (Banach Space). A Banach space is a complete normed space.

Definition 3.4 (Seminorm). A seminorm p is a norm without the positivity condition. Instead, we have positive semidefiniteness, i.e.

$$p(v) \geq 0 \quad \forall v \in V.$$

Note that a seminorm does not form a metric space. We define the *kernel* of the seminorm as the set $\{v \in V \mid p(v) = 0\}$. Note that this is a subspace. To get an actual norm, we must “divide” by the kernel.

We can define a norm on the quotient vector space $V/\ker p$ by letting the seminorm p act on any representative of this space.

Theorem 3.5 (Quotient of Norms). The norm $\|\cdot\| : V/\ker p \rightarrow \mathbb{R}, [v] \mapsto p(v)$ is a norm.

Proof. First, we show that it is well defined. Consider $v \in V$ and let $u \in \ker p$ be arbitrary. Then we have

$$p(v + u) \leq p(v) + p(u) = p(v)$$

by the triangle inequality, and conversely

$$p(v) = p(v - u + u) \leq p(v + u) + p(u) = p(v + u)$$

which shows that the norm is independent of the choice of representative.

The triangle inequality and homogeneity follow from the same properties of the seminorm. It is also true by definition that the norm is positive. \square

Thus, given a seminorm on a vector space, we can construct a new vector space that has a norm, and thus an induced topology.

Theorem 3.6 (Operator Norm). Let V, W be normed spaces and let $A : V \rightarrow W$ be a linear map. Then the following statements are equivalent:

- (a) A is continuous.
- (b) There exists a constant $c \geq 0$ such that

$$\|A(v)\| \leq c\|v\|$$

for all $v \in V$.

Proof. We use the fact that continuity is equivalent to continuity at 0. Then, we simply unravel the definitions. A is continuous at 0 if for every neighbourhood $Z \subseteq W$ of 0_W we have a neighbourhood $U \subseteq V$ of 0_V such that $A(U) \subseteq Z$.

Equivalently, we can consider open balls in place of Z and U . This is then equivalent to the second condition. \square

Note that we could have proven a weaker theorem in the case of topological vector spaces. To do that, we would define boundedness as follows:

Definition 3.7. A function $\Lambda : V \rightarrow W$ between topological vector spaces is bounded if it maps bounded sets into bounded sets

where a bounded set is defined as

Definition 3.8. A subset U of a topological vector space V is bounded if for each neighbourhood V of 0 there exists $t \in \mathbb{K}$ such that $U \subseteq tV$.

We can see that boundedness in the normed setting implies boundedness in the general topological vector space setting. In general, however, continuity implies boundedness, and the converse is not true. Thus, we will only discuss this in Banach spaces in the future.

Definition 3.9 (Operator Norm). The operator norm $\|A\|$ is defined as

$$\|A\| = \inf\{c \mid \|Av\| \leq c\|v\| \ \forall v \in V\}$$

Corollary 3.10 (Equivalent Statements).

(a)

$$\|A\| = \sup_{V \ni v \neq 0} \frac{\|Av\|}{\|v\|}$$

(b)

$$\|A\| = \sup_{\|v\|=1} \|Av\|$$

Theorem 3.11 (Operator Composition). Let V, W, Z be normed spaces, $A \in L(V, W)$, $B \in L(W, Z)$. Then

$$\|B \circ A\| \leq \|A\| \|B\|.$$

Proof. For $v \in V$, we have

$$\|BAv\| \leq \|B\| \|Av\| \leq \|B\| \|A\| \|v\|.$$

□

Note that the above definition shows once again the continuity of composition. Also note that this norm turns $L(V, W)$ into a normed space.

3.1.2 Bases

Definition 3.12 (Hamel Basis). A Hamel basis is a set $B \subseteq V$ such that for all $v \in V$, we have

$$v = \sum_{k=1}^n a_k e_k$$

with $a_i \in \mathbb{K}$ and $e_i \in B$ for all i , and

$$\sum_{k=1}^n a_k e_k = 0 \implies a_i = 0 \forall i.$$

The existence of Hamel bases is equivalent to the axiom of choice, and can be proven by Zorn's Lemma. The proof is as follows: We construct minimal spanning sets and maximally linearly independent sets by Zorn's Lemma (partial order by inclusion), and show that they are the same. These sets are Hamel bases.

A Hamel basis has all the beloved properties of a basis from finite dimensional linear algebra. For example, a linear map is uniquely defined through its action on the Hamel basis:

$$Av = \sum_{k=1}^n a_k A e_k.$$

However, a Hamel bases are usually difficult to come by, as we see with the following theorem:

Theorem 3.13. A Banach space with a countable Hamel basis is finite dimensional.

Proof. Suppose we have a countable Hamel basis $(e_n)_{n \in \mathbb{N}}$ of a Banach space V . Denote $M_n = \text{span}(\{e_1, \dots, e_n\})$. This is a closed proper subspace of V . Thus it has empty interior. However, because by definition the e_n s form a Hamel basis, we have $V = \bigcup_{k=1}^{\infty} M_k$, contradicting Baire's category theorem. \square

Note that Hamel bases generally interact very poorly with the topology. For example, the projection maps do not have to be continuous.

Example 3.14. Let V be the space of polynomials on the interval $[0, 1]$ equipped with the norm $\|p\| = \int_0^1 |p(x)| dx$. Then the projection onto the constant term is not continuous.

Proof. The sequence

$$\int_0^1 |(x-1)|^n dx = \frac{(-1)^{n+1}}{n+1}$$

converges to 0, but the projection of the polynomials onto the constant term yields the sequence $(-1)^n$, which is not convergent. Thus, the projection cannot be continuous. \square

In fact, this poor interplay yields the most interesting application of Hamel bases - the existence of *discontinuous* linear functionals.

Theorem 3.15 (Existence of Discontinuous Functionals). Let V be an infinite dimensional normed space. Then there exist discontinuous linear functionals on V .

Proof. Choose a countable linearly independent set $\{v_n\}$. Without loss of generality, we can normalize all vectors to norm 1. This set is contained in some Hamel basis B . Define $f(v_n) = n$, and $f(v) = 0$ for $v \in B \setminus \{v_n\}$. This functional is unbounded. \square

Definition 3.16 (Schauder Basis). A Schauder basis is a countable set $\{e_n\} \subseteq V$ such that all vectors $v \in V$ can be uniquely expressed as a sum

$$v = \sum_{k=1}^{\infty} a_k e_k,$$

where the convergence is understood to be in the topology of the vector space.

The importance of a Schauder basis is that it is countable, and we are still able to define *some* linear maps by their action on the basis. In particular, for any continuous linear map A , we have

$$Av = \sum_{k=1}^{\infty} a_k A e_k.$$

Also important to note is that this basis must be *ordered*, since the sum does not necessarily converge unconditionally.

3.1.3 Examples

Example 3.17. The space c is the space of convergent sequences.

Example 3.18. The space c_0 is the space of sequences converging to 0. It is a subspace of c with codimension 1.

Example 3.19. The space c_{00} is the space of sequences that are 0 outside of finitely many elements. This is a subspace of c_0 . It is the prototype for a non complete space.

Example 3.20. There are discontinuous linear functionals.

The simplest example in a non-complete space is the functional $f(e_i) = i$ defined on c_{00} .

In a complete space, this is significantly harder and requires the axiom of choice. This cannot be done with Hahn-Banach (to come), as Hahn-Banach extends bounded functionals.

Example 3.21 (Non-Closed Subspaces). Every subspace of a Banach space with countable dimension cannot be closed, as it would then be complete, a contradiction, as Banach spaces cannot have countable dimension.

3.2 The Hahn-Banach Theorems

3.2.1 Sublinear Functionals

Definition 3.22. Let V be a vector space over \mathbb{K} . A functional p is called *sublinear*, if

- (a) (Homogeneity) $p(rx) = rp(x)$ for all real $r \geq 0$ and $v \in V$.
- (b) (Subadditivity) $p(u + v) \leq p(u) + p(v)$ for all $u, v \in V$

The main property that relates sublinear functionals to linear functionals is as follows:

Theorem 3.23. If p is a sublinear functional on a real vector space V , then the following are equivalent:

- (a) p is linear
- (b) $p(v) + p(-v) = 0$ for all $v \in V$.
- (c) $p(v) + p(-v) \leq 0$ for all $v \in V$

Proof. If p is linear, then $p(v) + p(-v) = p(v - v) = p(0) = 0$. Clearly, (b) implies (c).

Now assume (c). First, we prove (b):

$$0 = p(v - v) \leq p(v) + p(-v) \leq 0$$

or $p(v) + p(-v) = 0$, proving (b). Using (b) we have, for $r < 0$,

$$p(rv) = -p(-rv) = rp(v),$$

proving the first aspect of linearity. Then we have

$$p(v + w - w) \leq p(v + w) + p(w) = p(v + w) + p(-w),$$

implying

$$p(v) + p(w) \leq p(v + w).$$

The subadditivity yields the other inequality, completing the proof of linearity. \square

We can define a partial order on the sublinear functions by defining p to be less than q if p is less than q pointwise. Zorn's Lemma then yields minimal elements. We will prove eventually that these minimal elements are linear functionals. Before that, however, we need two lemmas:

Theorem 3.24 (Auxiliary Functionals). If p is a sublinear functional on a real vector space V , then the auxiliary functional $q(v) = \inf\{p(v + tw) - tp(w) | t \geq 0, w \in V\}$ is a sublinear functional such that $q \leq p$.

Proof. By considering $t = 0$ in the infimum, we see that $q \leq p$. We only need to show sublinearity. Consider first $r = 0$. Then

$$q(rv) = \inf\{p(rv + tw) - tp(w) | t \geq 0, w \in V\} = \inf\{p(tw) - tp(w) | t \geq 0, w \in V\} = 0.$$

For $r > 0$, we have

$$\begin{aligned} q(rv) &= \inf\{p(rv + tw) - tp(w) | t \geq 0, w \in V\} \\ &= \inf\left\{rp\left(v + \frac{t}{r}w\right) - tp(w) | t \geq 0, w \in V\right\} \\ &= r \inf\left\{p\left(v + \frac{t}{r}w\right) - \frac{t}{r}p(w) | t \geq 0, w \in V\right\} \\ &= rq(v) \end{aligned}$$

For subadditivity, we take the special points $w = \frac{1}{s+t}(sx + ty)$, or $(s+t)w = sx + ty$. Then

$$\begin{aligned} q(x + y) &\leq p(x + y + (s+t)w) - (s+t)p(w) \\ &\leq p(x + sw) - sp(w) + p(y + tw) - tp(w) \end{aligned}$$

which shows that $q(x + y) \leq q(x) + q(y)$. \square

Now, we move on to the proof of the main result.

Theorem 3.25. A sublinear functional p on a real vector space V is linear iff it is minimal.

Proof. Suppose we have $q \leq p$, with q sublinear and p linear. Since q is sublinear, we have $0 = q(v - v) \leq q(v) + q(-v)$, which implies $-q(-v) \leq q(v)$. Since $q(-v) \leq p(-v) = -p(v)$, we have $p(v) \leq -q(-v) \leq q(v)$, suggesting that $p \leq q$. Thus $p = q$.

Conversely, suppose that p is a minimal sublinear functional. Then we must have $q = p$ where q is the auxiliary sublinear functional defined in Theorem 3.24. If we let $t = 1$ and $w = -v$ in the above definition, we get

$$p(v) \leq p(v - v) - p(-v),$$

or $p(v) + p(-v) \leq 0$, as desired. Finally, we show the boundedness which we require to apply Zorn's Lemma \square

Theorem 3.26. For every sublinear functional p , we have a linear functional $f \leq p$.

Proof. We begin, as always, by considering a chain of sublinear functionals, which is also a totally ordered set of sublinear functionals P . Suppose that $q(x)$ is unbounded from below for all $q \in P$ and some $x \in V$. Then we have, for all n , a p_n such that $p_n(x) \leq -n$.

Then we construct a decreasing sequence of sublinear functionals q_n using $q_n = \min(p_1, \dots, p_n)$. Since $q_n(x) \leq -n$, we have

$$0 = q_n(x - x) \leq q_n(x) + q_n(-x) \leq -n + q_n(-x)$$

or $q_n(-x) \geq n$. Then we have, for all n , $n \leq q_n(x) \leq q_1(x)$, a contradiction. Thus we can take the infimum for all $x \in V$. This yields a functional q^* . It remains to show that q^* is sublinear.

Since $q_n(0) = 0$ for all n , we have $q^*(0) = 0$. By homogeneity, $q^*(rv) = \inf\{q_n(rv) | n \in \mathbb{N}\} = r \inf\{q_n(v) | n \in \mathbb{N}\}$. Finally, we have $q^*(u + v) = \inf\{q_n(u + v) | n \in \mathbb{N}\} \leq q^*(u) + q^*(v)$. \square

Theorem 3.27 (Dominated Extension). Let X be a real vector space, p a sublinear functional on X , M a subspace of X and $f : M \rightarrow \mathbb{R}$ a linear functional on M such that $f \leq p$. Then there is a linear functional F on X that extends f such that $F \leq p$.

Proof. The overarching idea of this proof is as follows: We seek a sublinear functional q such that $q \leq p$ on X , and $q \leq f$ on M . By the previous theorem, we will then get a linear functional $F \leq q \leq p$ on X and $F \leq f$ on M . Since f is maximal on M , we then have $f \leq F$, or $f = F$ on M .

We choose as our candidate

$$q(x) = \inf\{p(x + m) + f(m) | m \in M\}.$$

It is clear that $q \leq p$ on X . We first show that q is real valued. We do this by showing two different estimates:

First, we have $f(-m) \leq p(-m)$. Thus, for all $x \in X$, we have $-p(-x) + f(-m) \leq -p(-x) + p(m)$, or

$$-p(-x) \leq p(m) - p(-x) - f(-m).$$

Then, we estimate $p(-m) \leq p(x - m) + p(-x)$ and thus

$$p(-m) - p(-x) \leq p(x - m).$$

Combining the two equalities, we get

$$-p(-x) \leq p(x - m) + f(m)$$

which shows that q is real valued. Now we show sublinearity.

1. Since $f(-m) \leq p(-m)$, we have $p(-m) + f(m) \geq 0$. This shows that $q(0) \geq 0$. Conversely, we have $p(0) + f(0) = 0$, which shows that $q(0) \leq 0$.
2. For $a \in \mathbb{R}_0^+$, we have $q(ax) = \inf\{p(ax - m) + f(ax)\} = \inf a\{p(x - m/a) + af(m/a)\} = aq(x)$.
3. For sublinearity, we choose $x, y \in X$ and $r > 0$. There exists $m, n \in M$ such that

$$q(x) > p(x - m) + f(m) + \frac{r}{2}, \quad q(y) > p(y - n) + f(n) + \frac{r}{2}.$$

Then

$$q(x) + q(y) \geq p(x + y - (m + n)) - f(m + n) - r \geq q(x + y) - r. \quad \square$$

Theorem 3.28 (Hahn-Banach (Normed)). Let V be a normed space, $W \subseteq V$ a subspace, and f a continuous linear functional on W . Then there is a continuous extension of f to V .

Proof. We consider $g : V \rightarrow \mathbb{R}$, $v \mapsto \|f\|\|v\|$. This map is clearly sublinear and $f \leq g$. Then, by the Dominated Extension Theorem, we get an extension to all of V . Because this map is bounded, it is continuous. \square

Corollary 3.29. Let V be a normed space and $v \in V$. Then there exists a linear functional $\varphi \in V'$ such that $\varphi(v) = \|v\|$.

Proof. We let $W = \text{span}\{v\}$ and $\varphi(x) = k$ if $x = kv$ in Theorem 3.28 \square

3.2.2 Geometric Hahn-Banach

Definition 3.30. Let V be a normed space. A subset A of V is called

- (a) *Absorbing*, if for every $v \in V$ there is $\lambda > 0$ with $v \in \lambda A$.
- (b) *Balanced*, if for all $|z| \leq 1$ we have $zA \subseteq A$.
- (c) *Convex*, if for all $v, w \in A$ and $\lambda \in [0, 1]$ we have $\lambda v + (1 - \lambda)w \in A$.
- (d) *Absolutely convex*, if it is balanced and convex.

Theorem 3.31. Let V be a vector space over \mathbb{K} .

- (a) If $p : V \rightarrow \mathbb{R}_0^+$ is a seminorm then

$$B_{p,1}(0) = \{v \in V | p(v) < 1\}$$

and

$$B_{p,1}(0)^{\text{cl}} = \{v \in V | p(v) \leq 1\}$$

are absorbing and absolutely convex.

- (b) If $C \subseteq V$ is convex, balanced and absorbing then

$$p_C(V) = \inf\{\lambda | \lambda > 0, v \in \lambda C\}$$

is a seminorm.

- (c) For C absolutely convex and absorbing

$$B_{p_C,1}(0) \subseteq C \subseteq B_{p_C,1}(0)^{\text{cl}}.$$

Proof. (a) It is clear from the definition of a seminorm that both of these are absolutely convex.

If $v \in V$ and $p(v) = \alpha$, then $p(v/2\alpha) = 1/2$, which tells us that $v/2\alpha$ is an element of the ball, and thus the ball is absorbing.

- (b) First, we show absolute homogeneity. We note that $v \in \lambda C$ if and only if $|\alpha|v \in |\alpha|\lambda C$. Since C is balanced, so is $|\alpha|\lambda C$, and thus this happens iff $\alpha v \in |\alpha|\lambda C$. Since the infimum is absolutely homogeneous, this shows that the p_c is too.

Suppose $v \in \lambda_v C$ and $u \in \lambda_u C$. If we prove that $u + v \in (\lambda_u + \lambda_v)C$, we will be done. In fact, we know more - we know that $aC + bC = (a + b)C$. Let us prove this.

Suppose $v \in aC + bC$. Then $v = au + bw$ for $u, w \in C$. Since C is convex, we have

$$\frac{v}{a+b} = \frac{a}{a+b}u + \frac{b}{a+b}w \in C$$

suggesting that $v \in (a+b)C$.

- (c) Clearly, for all $v \in C$, we have $p(v) \leq 1$.

If $p(v) < 1$, then $v \in \lambda C$ for some $\lambda < 1$. Since C is balanced, this shows that $v \in C$. □

Definition 3.32 (Minkowski Functional). Let $C \subseteq V$ be an absorbing subset in a vector space V over a \mathbb{K} . Then $p_c : V \rightarrow \mathbb{R}_0^+$ defined by

$$p_c(v) = \inf\{\lambda > 0 | v \in \lambda C\}$$

is called the Minkowski functional of C .

(b) of the previous theorem shows that this is a seminorm if C is absolutely convex too.

Theorem 3.33 (Separation I). Let V be a normed space with two nonempty disjoint convex subsets $A, B \subseteq V$.

- (a) Suppose A is open. Then there exists $\Phi \in V'$ and $\alpha \in \mathbb{R}$ such that

$$\operatorname{Re}(\Phi(v)) < \alpha \leq \operatorname{Re}(\Phi(u))$$

for $v \in A, u \in B$

- (b) Suppose both A and B are open. Then there exists $\Phi \in V'$ and $\alpha \in \mathbb{R}$ with

$$\operatorname{Re}(\Phi(v)) < \alpha < \operatorname{Re}(\Phi(u))$$

for $v \in A, u \in B$.


Proof. Fix $a_0 \in A$, $b_0 \in B$. We define $x_0 = b_0 - a_0$, and let $C = A - B + x_0$. Then C is a convex neighbourhood of 0. Let p be its Minkowski functional. Since C is a neighbourhood of 0, it is absorbing. Thus, its Minkowski functional is a sublinear functional on X , with $p(x_0) \geq 1$.

Then, by Corollary 3.29, we define a linear functional on the subspace spanned by x_0 such that $f(x_0) = 1$. Since this is bounded by p , f extends to a linear functional Λ on X . Since $\Lambda(C) \leq 1$, we have $\Lambda(-C) \geq -1$ and Λ is bounded on a neighbourhood $C \cap -C$ of 0 and is hence continuous. Then, we let $a \in A, b \in B$, and note that $a - b + x_0 \in C$, thus

$$\Lambda(a) - \Lambda(b) + 1 = \Lambda(a - b + x_0) < 1,$$

or $\Lambda(a) < \Lambda(b)$. Thus the images of A and B under Λ are convex subsets of the real line, and the image of A is an open subset, since linear functionals are always open maps. However, the only convex open subsets of the real line are the intervals, and choosing the right endpoint of $\Lambda(A)$ yields (a). (b) follows almost identically. \square

The picture for this theorem is as follows: A linear functional Φ defines a hyperplane for each fixed value of α , then splitting the space into two parts, one where the functional is less than, and one where the functional is greater than α . In \mathbb{R}^2 , the picture looks as follows:



Theorem 3.34 (Separation II). Let V be a normed vector space, $K \subseteq V$ compact, convex, nonempty, $C \subseteq V$ closed, convex, nonempty, with $K \cap C = \emptyset$. Then there exists $\Phi \in V'$ and $\alpha, \beta \in \mathbb{R}$ with $\operatorname{Re} \Phi(v) < \alpha < \beta < \operatorname{Re} \Phi(u)$ for all $v \in K$ and $u \in C$.

Proof. There is a convex neighbourhood V of 0 in V such that $K + V \cap C = \emptyset$ (Corollary 2.7). Then, we apply Theorem 3.33 to $K + V$ and C to get a linear functional that satisfies all the required properties, since K is a compact subset of $K + V$.

Note that we used that V is normed in the first line, as Corollary 2.7 only yields a not necessarily convex neighbourhood. Local convexity would also have sufficed. \square

Theorem 3.35 (Separation III). Let V be a normed space, $K \subseteq V$ compact convex, $C \subseteq V$ closed convex nonempty, balanced, such that $K \cap C = \emptyset$. Then there exists $\Phi \in V'$ such that $\sup_{u \in C} |\Phi(u)| < \inf_{v \in K} |\Phi(v)|$.

Proof. Follows directly from Theorem 3.35. \square

Corollary 3.36. Let $C \subseteq V$ be a closed balanced convex subset and $v \in V \setminus C$. Then there exists $\Phi \in V'$ with $\Phi(v) > 1$, $\Phi(u) \leq 1$ for all $u \in C$.

Proof. Apply Theorem 3.35 with $K = \{v\}$ and rescale as necessary. \square

3.3 Duality

3.3.1 The Weak Topology

It is often the case that we want to work with components of vectors. However, due to the difficulty of picking a basis, we cannot define the components the same way we do in finite dimensional linear algebra. Thus, we define the components to be the outputs of linear functionals applied to vectors. This notion leads us to the weak topology.

Definition 3.37. Let V be normed.

- (a) For $\varphi \in V'$ we define the seminorm $p_\varphi : V \rightarrow \mathbb{R}_0^+$ by $p_\varphi(v) = |\varphi(v)|$ and one calls $B_{p_\varphi, r}(v) = \{w \in V \mid p_\varphi(v - w) < r\}$ the open ball.
- (b) We define the weak topology to be the topology on V generated by all open balls $B_{p_\varphi, r}(v)$.

Equivalently, we could also have defined the weak topology as the weakest topology such that all the seminorms p_φ are continuous. Note that the weak topology is locally convex as the subbasis elements are convex due to Theorem 3.31.

Theorem 3.38. Let V be a normed space.

- (a) The weak topology turns V into a topological vector space.
- (b) The weak topology is coarser than the norm topology.
- (c) $\varphi \in V^*$ is weakly continuous iff φ is norm continuous.
- (d) The weak topology is Hausdorff
- (e) p_φ for $\varphi \in V'$ is weakly continuous.

Proof. (a) We only check the continuity condition for a subbasis:

$$p_\varphi(u + u' - (v + v')) \leq p_\varphi(u - v) + p_\varphi(u' - v') < r$$

which implies that $u + u' \subseteq B_{p_\varphi, r}(v + v')$. Thus

$$B_{p_\varphi, \frac{r}{2}}(v) \times B_{p_\varphi, \frac{r}{2}}(v') \subseteq +^{-1}B_{p_\varphi, r}(v + v').$$

Thus, $+$ is continuous at (v, v') . Continuity of multiplication follows by the homogeneity of seminorms.

- (b) If $p_\varphi(v - v') < \epsilon$, we also have $p_\varphi(v - v') = |\varphi(v - v')| \leq \|\varphi\| \|v - v'\|$, thus showing that we can choose δ such that $B_\delta(v) \subseteq B_{p_\varphi, \epsilon}(v)$.
- (c) If φ is weakly continuous it is norm continuous because of (b). Conversely, we see that $\varphi^{-1}(B_r(0)) = B_{p_\varphi, r}(0)$, which is an element of the subbasis. Thus, φ is weakly continuous.
- (d) Suppose $v \neq 0$. Then, we define a continuous linear functional φ such that $r = \varphi(v) > 0$ by Hahn-Banach. Then $B_{p_\varphi, r/2}(0) \cap B_{p_\varphi, r/2}(v) = \emptyset$.
- (e) Follows by definition. □

Theorem 3.39. Let V be a topological vector space.

- (a) A net $(v_i)_{i \in I}$ in V is weakly convergent to $v \in V$ if and only if for every $\varphi \in V'$ one has $\lim_{i \in I} \varphi(v_i) = \varphi(v)$.
- (b) A net $(v_i)_{i \in I}$ in V is a weak Cauchy net if and only if for every $\varphi \in V'$ the net $(\varphi(v_i))_{i \in I}$ is a Cauchy net in \mathbb{K} .

Proof. (a) Since φ is (weakly) continuous, we have the desired equality.

Conversely, assume that the equality is satisfied. What we need to show is that the net v_i is eventually in every neighbourhood of v . Since V is locally convex, it suffices to consider absolutely convex neighbourhoods K . Consider the Minkowski functional of K , p_K . Since p_K also generates the weak topology, we can find $\alpha \in I$ such that $\beta \succeq \alpha \implies p_K(v_i - v) < 1$, or $v_i \in K$.

- (b) Follows analogously to (a). □

Theorem 3.40. Let V be a normed space and let $C \subseteq V$ be convex. Then the weak closure of C coincides with the norm closure.

Proof. Since the weak topology is weaker, $C^{\text{cl}} \subseteq C^{\text{wcl}}$.

Conversely, we choose $v_0 \in V$, $v_0 \notin C^{\text{cl}}$. Then Theorem 3.34 yields $\gamma \in \mathbb{R}$ and $\Lambda \in V'$ such that

$$\operatorname{Re} \Lambda x_0 < \gamma < \operatorname{Re} \Lambda x.$$

Thus, the set $\{x \mid \operatorname{Re} \Lambda x < \gamma\}$ is a weak neighbourhood of x_0 that does not intersect C , showing that $x \notin C^{\text{wcl}}$. □

Corollary 3.41. Let $U \subseteq V$ be a subspace of a normed space. Then $U^{\text{cl}} = U^{\text{wcl}}$.

Proof. Subspaces are automatically convex. □

Theorem 3.42. Let V be a normed space. Then the canonical map $i : v \rightarrow V''$ is norm preserving and hence injective.

Proof. We define this injection by

$$i(v)\varphi = \varphi(v).$$

We have $i \in V''$ and $\|i(v)\| \leq \|v\|$ for a given $v \in V$. This implies that there is $\varphi \in V'$ with $\|\varphi\| = 1$ and $\varphi(v) = \|v\|$. Thus,

$$|i(v)\varphi| = |\varphi(v)| = \|v\| = \|v\|\|\varphi\|.$$

Thus, $\|i(v)\| \geq \|v\|$. Together, we have $\|i(v)\| \geq \|v\|$. □

Theorem 3.43 (Completion of normed space). Every normed space V can be completed to a Banach space. More precisely $i(v)^{\text{cl}} \subseteq V''$ is a completion.

Proof. Note that the dual space of *any* TVS is always complete. As a closed subspace of a complete space, we have a complete space. \square

The canonical map i allows us to define a weak topology on V' generated by the functionals $i(V) \in V''$. We call this the weak *-topology. Convergence in the weak *-topology is defined by the same conditions as in the weak topology. In particular, a net $(v_i)_{i \in I} \subseteq V'$ converges to $v \in V'$ if and only if $v_i(x) \rightarrow v(x)$ for all $x \in V$ - that is, the linear functionals converge pointwise.

Theorem 3.44. Let V, W be normed spaces with $A : V \rightarrow W$ continuous and linear. Then

$$\|A\| = \sup_{\varphi \neq 0, w \neq 0} \frac{|\varphi(Av)|}{\|\varphi\|_{W'} \|v\|_V} = \sup_{\|\varphi\| = \|v\| = 1} |\varphi(Av)| = \|A'\|.$$

3.3.2 Polars & Banach-Alaoglu Theorems

Definition 3.45. Let V be a topological vector space with topological dual V' .

(a) The polar A^* of a subset $A \subseteq V$ is defined by

$$A^* = \{\varphi \in V' \mid |\varphi(v)| \leq 1 \text{ for all } v \in A\}.$$

(b) The polar B_* of a subset $B \subseteq V'$

$$B_* = \{v \in V \mid |\varphi(v)| \leq 1 \text{ for all } \varphi \in B\}.$$

The polar A^* of a set $A \subseteq V$ is the set of linear functionals which are small on A . Note that the 1 in the definition comes without loss of generality, as

$$\varphi(v) < r \iff \frac{\varphi}{r}(v) < 1.$$

Theorem 3.46. Let V be a topological vector space and $B \subseteq V'$. Then $B_* \subseteq V$ is absolutely convex and weakly closed.

Proof. For $z \in \mathbb{K}$ with $|z| \leq 1$ and $v \in B_*$, then for $\varphi \in B$ we have $|\varphi(zv)| = |z||\varphi(v)| \leq 1$, which shows that B_* is balanced.

Then we choose $\lambda \in [0, 1]$ and $v, w \in B_*$. Then for $\varphi \in B$ we have

$$\varphi(\lambda v + (1 - \lambda)w) \leq \lambda|\varphi(v)| + (1 - \lambda)|\varphi(w)| \leq 1.$$

This implies that B_* is convex.

To show that it is closed, we have

$$\begin{aligned}
 B_{**} &= \{v \in V \mid |\varphi(v)| \leq 1 \text{ for all } \varphi \in B\} \\
 &= \{v \in V \mid p_\varphi(v) \leq 1 \text{ for all } \varphi \in B\} \\
 &= \bigcap_{\varphi \in B} B_{p_\varphi, 1}(0)^{\text{cl}} \\
 &= \bigcap_{\varphi \in B} p_\varphi^{-1}([0, 1])
 \end{aligned}$$

which is an intersection of closed sets and hence closed. \square

We also have the following facts

Theorem 3.47. Let V be a topological vector space.

- (a) If $A \subseteq B \subseteq V$, then $A^* \supseteq B^*$.
- (b) For $a \neq 0$, we have $(aA)^* = a^{-1}A^* = |a|^{-1}A^*$.
- (c) $A \subseteq A^{***}$ and $A^* = A^{****}$.

Proof. (a) If $|\varphi(v)| < 1$ for all $v \in B$, then it holds true for all $v \in A$.

(b) We have $\varphi(av) = a\varphi(v)$, as well as $|\varphi(av)| = |a||\varphi(v)|$.

(c) We have $A \subseteq A^{***}$ by definition. Thus, $A^* \subseteq (A^*)^{**}$. However, by (a), $A \subseteq A^{***}$ implies that $A^* \supseteq A^{****}$. \square

Next, we will consider what happens to polar sets of hulls. However, before we continue, we first need to define what hulls are.

Definition 3.48. The convex hull C_c of a subset $C \subseteq V$ is the intersection of all convex sets containing C .

The most useful characterisation of convex hulls is as follows:

Lemma 3.49. The convex hull C_c of a set C is the set consisting of all convex combinations of vectors of C .

Proof. Let K be the set of all convex combinations of elements of C . Since K is convex, we have $C_c \subseteq K$. Conversely, all linear combinations of elements of C must be in C_c , and thus $K \subseteq C_c$. \square

We also have balanced hulls:

Definition 3.50. The balanced hull C_b of a subset $C \subseteq V$ is the intersection of all balanced sets containing C .

Again, this is characterised by the following Lemma:

Lemma 3.51. The balanced hull C_b of a subset $C \subseteq V$ is the union of all sets zC , with $|z| \leq 1$.

Proof. We define

$$K := \bigcup_{|z| \leq 1} zC.$$


Clearly, K is balanced, and thus $C_b \subseteq K$. Conversely, we must have $K \subseteq C_b$. \square

We also have the absolutely convex hull of a set

Definition 3.52. The absolutely convex hull A_{bc} of a subset $A \subseteq V$ is defined by $(A_b)_c$.

We note that this is convex by definition, and we can easily verify that it is also balanced. It is also important to note that this is not the same as $(A_c)_b$, as $(A_c)_b$ may not be convex.

Example 3.53. Let A be the set in \mathbb{R}^2 consisting of $(0,0)$, $(1,0)$ and $(1,1)$. Its convex hull is the triangle consisting of these 3 points, and its balanced hull is its convex hull plus its mirror image, which is not convex.



Now we are ready to discuss their polar sets:

Theorem 3.54 (Polars of Hulls). Let V be a topological vector space, $A \subseteq V$.

- (a) $A^* = (A_b)^*$.
- (b) $A^* = (A_c)^*$.
- (c) $A^* = (A^{\text{wcl}})^*$.
- (d) $A^* = ((A_{bc})^{\text{wcl}})^*$.

Proof. For all of these, the reverse inclusion follows due to A being a subset of the other set.

- (a) Now suppose $|\varphi(v)| \leq 1$ for all $v \in A$. Then consider $u \in A_b$. We have $u = zv$ for some $v \in A$, and thus $\varphi(u) = \varphi(zv) = z\varphi(v)$.

(b) Consider $u \in A_c$, meaning that $u = \sum_{k=1}^n \alpha_k v_k$ for $\sum_{k=1}^n \alpha_k = 1$, $v_k \in A$ for all k . We have

$$\varphi(u) = \sum_{k=1}^n \alpha_k \varphi(v_k) \leq \sum_{k=1}^n \alpha_k = 1.$$

(c) Suppose $|\varphi(v)| \leq 1$ for all $v \in A$. Then, if $w \in A^{\text{wcl}}$, it means that there is a net $(v_i)_{i \in I} \subseteq A$ such that $v_i \rightarrow w$. In the weak topology, this means that $\varphi(v_i) \rightarrow \varphi(w) \leq 1$.

(d) We note that $A \subseteq (A_{bc})^{\text{wcl}}$. Since A^{***} is absolutely convex and weakly closed by Theorem 3.46, we have $(A_{bc})^{\text{wcl}} \subseteq A^{***}$. This implies that

$$A^* \supseteq (A_{bc})^{\text{wcl}} \supseteq A^{***} = A^*.$$

□

Theorem 3.55 (Bipolar Theorem). Let A be a subset of a topological vector space V . Then $A^{***} = (A_{bc})^{\text{wcl}}$.

Proof. As in 3.54 we have $A \subseteq (A_{bc})^{\text{wcl}} \subseteq A^{***}$. Choose $w \notin (A_{bc})^{\text{wcl}}$. Then, by Theorem 3.35, there is a weakly continuous linear functional Φ such that $\sup_{v \in (A_{bc})^{\text{wcl}}} |\Phi(v)| < \Phi(w)$. Since $(A_{bc})^{\text{wcl}}$ is balanced, we have $0 = \Phi(0) < a$. Hence, we may rescale the functional to get a linear functional Ψ with $\sup_{v \in (A_{bc})^{\text{wcl}}} |\Psi(v)| = 1$. This implies that $\Psi \in A^*$. Since $\Psi(w) > 1$, we have $w \notin A^{***}$, completing the proof. □

Theorem 3.56 (Banach-Alaoglu). If U is a balanced neighbourhood of 0 in the topological vector space V then U^* is weak *-compact.

Proof. Since neighbourhoods of 0 are absorbing, we choose $\gamma(v)$ for each $v \in V$ such that $v \in \gamma(v)U$. Then we define a set P by $P = \prod_{v \in V} K_{\gamma(v)}(0)^{\text{cl}}$. Since the disks are compact, P is compact by Tychonoff's Theorem. P consists of all functions $f : V \rightarrow \mathbb{K}$ such that $|f(x)| \leq \gamma(x)$. Thus, we have $U^* \subseteq V' \cap P$.

We now have two topologies on U^* , induced by V' and P respectively. We show that these topologies coincide, and U^* is closed in the topology induced by P . Since P is compact, this will imply that U^* is compact.

Fix $\Lambda_0 \in A^*$, and choose $v_i \in V$ for $i = 1, \dots, n$ as well as $\delta > 0$

$$W_1 = \{\Lambda \in V' \mid |\Lambda v_i - \Lambda_0 v_i| < \delta, i = 1, \dots, n\}$$

$$W_2 = \{f \in P \mid |f v_i - \Lambda_0 v_i| < \delta, i = 1, \dots, n\}$$

The W s form a local basis for the topology. Since $A^* \cap W_1 = A^* \cap W_2$, the topologies coincide.

Now suppose that we have f_0 in the P -closure of A^* . Then we choose scalars α, β , and $u, v \in V$. The set of all f such that $|f - f_0| < \epsilon$ at x, y and $\alpha x + \beta y$ is a P -neighbourhood of f_0 . Thus, A^* contains such an f . Since f is linear, we have

$$\begin{aligned} & f_0(\alpha x + \beta y) - \alpha f_0(x) - \beta f_0(y) \\ &= (f_0 - f)(\alpha x + \beta y) + \alpha(f - f_0)(x) + \beta(f - f_0)(y). \end{aligned}$$

Approximating each of these by ϵ , we have

$$|f_0(\alpha x + \beta y) - \alpha f_0(x) - \beta f_0(y)| = \epsilon(1 + |\alpha| + |\beta|)$$

showing that f_0 is linear. Similarly, we conclude that $|f_0(v)| \leq 1$ for all $v \in A$. Thus, we have $f_0 \in A^*$, completing the proof. □

The importance of Banach-Alaoglu comes from sequential compactness, which we investigate in the following theorems:

Definition 3.57. A space X is called *sequentially compact* if every sequence has a convergent subsequence.

In a metric space, for example, a normed space, sequential compactness is equivalent to compactness, which we state without proof:

Theorem 3.58. Let X be a metric space. X is sequentially compact if and only if X is compact.

However, the unit ball in the dual space is not, in general, metrizable. We require an additional condition, that the original vector space is separable:

Theorem 3.59 (Sequential Compactness of Unit Ball). Let V be a separable Banach space. Then $B_1(0)^{\text{cl}} \subseteq V'$ is first countable in the weak *-topology.

Proof. First, we begin by letting $\{v_1, \dots\}$ be a countable dense subset of V . □

This implies that it is sequentially compact by the next theorem:

Theorem 3.60. A first countable compact space X is sequentially compact.

Proof. Suppose not. Then we can find a sequence $(x_n)_n \subseteq X$ such that it has no convergent subsequence.

First, we note that this means the set $\{x_n\}$ has no limit points. Then, for each $y \in X$ we choose an open set U_y that only contains finitely many points x_n . By hypothesis, we have a finite subcover U_{y_1}, \dots, U_{y_n} .

Since they cover the space, we cannot have all of them only contain finitely many points of x_n . This is a contradiction. □

3.3.3 Separability & Reflexivity

Definition 3.61. The closed linear span of a subset of V is the smallest closed linear subspace containing that subset.

Lemma 3.62. The closed linear span of a set $\{v_n\}$ is the closure of the linear span of v_n , i.e. the set of finite linear combinations of v_n .

Proof. We know that the closure of a linear space is also a linear space; this shows that the closed linear span is a subset of the closure of the linear span.

Conversely, the closure of the linear span must also be contained in the closed linear span. □

Theorem 3.63 (Spanning Criterion). A point v belongs to the closed linear span Y of a subset $\{y_n\}$ if and only if every continuous linear functional that vanishes on $\{y_n\}$ vanishes at v — that is,

$$\ell(y_j) = 0 \ \forall y_j \implies \ell(v) = 0.$$

Proof. One implication is obvious: Since ℓ is linear, it vanishes at finite linear combinations of the y_n s. Since it is continuous, it vanishes on their closure.

Conversely, suppose that v does not belong to the closed linear span of $\{y_n\}$. Then

$$\inf_{y \in Y} \|v - y\| = d > 0.$$

Then, define the subspace V to consist of all points of the form $y + av$, $y \in Y$, and define the linear functional ℓ_0 on Z by $\ell_0(y + av) = a$. Since we have

$$\|y + av\| \geq d|a|,$$

we can conclude that ℓ_0 is bounded by d^{-1} . Thus, we can extend this using the Hahn-Banach theorem to the entire Banach space. By definition, the extension vanishes on all of Y but has the value 1 at z . \square

Theorem 3.64. Let V be a normed space with separable dual V' in the norm topology. Then V is separable.

Proof. We choose a countable dense subset ℓ_n of V' . By definition of the norm, we choose v_n such that

$$\|v_n\| = 1, \ell_n(v_n) > \frac{1}{2}\|\ell_n\|.$$

We claim that the closed linear span of v_n is V . By 3.63, this means that we must show that every continuous linear functional that vanishes on v_n vanishes everywhere. Suppose otherwise. Suppose that there exists ℓ with $\ell(v_n) = 0$ for all v_n and $\|\ell\| = 1$.

Since the ℓ_n are dense, we can find ℓ_n with $\|\ell - \ell_n\| > \frac{1}{3}$. Since $\|\ell\| = 1$, this means that $\ell_n > \frac{2}{3}$. However, since $\ell(v_n) = 0$, we have

$$\frac{1}{3} > |(\ell - \ell_n)(v_n)| = |\ell_n(v_n)| > \frac{1}{2}\|\ell_n\|$$

a contradiction. Since finite linear combinations of the v_n s are dense, we can also consider finite linear combinations with rational coefficients; these are also dense, and countable. Thus, V is separable. \square

Now, we move on to considering the question of the double dual. In finite dimensions, there is only one real or complex vector space of dimension n up to isometry, namely \mathbb{R}^n or \mathbb{C}^n . Its dual is also isomorphic to itself, and thus its double dual is simply itself. In infinite dimensions, this is no longer the case. We call such spaces *reflexive*.

Definition 3.65. A Banach space V is called reflexive if $V'' = V$.

A natural question is why we talk about Banach spaces here. It seems that this definition can be extended to *any* normed space, where the dual inherits the operator norm topology. However, as we shall see, if a normed space is not Banach, it cannot be reflexive:

Theorem 3.66. The dual V' of a normed space V is a Banach space under the operator norm topology.

Proof. It was proven earlier that it is a normed space. For completeness, we let ℓ_n be a Cauchy sequence in V' . Then, since the reals are complete, we must have pointwise convergence of ℓ_n . We call this limit function ℓ .

By pointwise convergence, ℓ must also be linear and bounded, and thus continuous. It is also clear that $\ell_n \rightarrow \ell$. \square

Example 3.67. $V = c_0$ is separable but $V' = \ell^\infty$ is not.

Theorem 3.68. Pulling back the weak * topology from V'' to V reproduces the weak topology, but the weak topology is finer than the weak * topology on V' .

Theorem 3.69. Let V be a normed space.

- (a) If V is reflexive then V is a Banach space
- (b) If V is reflexive then the weak and weak * topology on V' coincide.
- (c) If U is a closed subspace of a reflexive space V then U is reflexive too.
- (d) Suppose V is complete. Then V is reflexive iff V' is reflexive.

Proof. (a) Clear since V'' is complete.

- (b) We consider $\Psi \in V''$. We have $p_v(\varphi) = |\varphi(v)| = |i(v)\varphi| = \|\Psi(\varphi)\|$. But since $E \mapsto nE$ is a homeomorphism, E is nonmeager. However, as an intersection of closed sets, E is closed. Together, this implies that E has an interior point x . Then $x - E$ is a neighbourhood of 0, meaning that there exists

□

Theorem 3.70. Let V be reflexive. Then $B_1(0)^{\text{cl}}$ is sequentially compact in the weak topology.

Corollary 3.71. Let V be reflexive.

- (a) A norm-bounded sequence in V has a weakly convergent subsequence.
- (b) Every weak Cauchy sequence converges weakly.

3.4 Banach-Steinhaus

Definition 3.72 (Equicontinuity). Let V and W be topological vector spaces, and Γ a collection of linear maps $V \rightarrow W$. Γ is called equicontinuous if for all neighbourhoods Z of 0_W , we have a neighbourhood V of 0_V such that $\Lambda(V) \subseteq Z$ for all $\Lambda \in \Gamma$.

Theorem 3.73. Let V and W be topological vector spaces, Γ an equicontinuous collection of linear maps, and E a bounded subset of V . Then there exists a bounded F in W such that $\Lambda(E) \subseteq F$ for all $\Lambda \in \Gamma$.

Proof. Let F be the union of the sets $\Lambda(E)$ for all $\Lambda \in \Gamma$. We show that this is bounded. By equicontinuity, we choose $Z \ni 0_W$ open, $U \ni 0_V$ open such that $\Lambda(U) \subseteq Z$ for all $\Lambda \in \Gamma$. Since E is bounded, we must have $E \subseteq tU$ for some $t \in \mathbb{R}$. Then $\Lambda(E) \subseteq t\Lambda(U) \subseteq tZ$ and thus F is bounded. \square

Theorem 3.74 (Banach-Steinhaus). Suppose V and W are topological vector spaces, and Γ is a collection of linear maps $V \rightarrow W$. Let B be the set of all $v \in V$ such that the orbits

$$\Gamma(v) = \{\Lambda v | \Lambda \in \Gamma\}$$

are bounded in W .

If B is nonmeager, then $B = V$ and Γ is equicontinuous.

Proof. We pick balanced neighbourhoods Z and U of 0_W such that $\overline{U} + \overline{U} \subseteq Z$. Then we set

$$E = \bigcap_{\Lambda \in \Gamma} \Lambda^{-1}(\overline{U}).$$

If $v \in B$, then $\Gamma(v) \in nU$ for some n , thus $v \in nE$. Hence, we have

$$B \subseteq \bigcup_{n=1}^{\infty} nE.$$

Since B is nonmeager, we have at least one nE nonmeager. However, since $E \mapsto nE$ is a homeomorphism, E must itself be nonmeager. Since E is closed, this means it has an interior point v . Then $v - E$ contains an open neighbourhood V of 0 , and

$$\Lambda(V) \subseteq \Lambda(v) - \Lambda(E) \subseteq \overline{U} - \overline{U} \subseteq W$$

for all $\Lambda \in \Gamma$. This shows equicontinuity and hence uniform boundedness by Theorem 3.73. \square

Often, the fact that B is nonmeager comes from properties of the space. Again, note as we did in 1.2 that even being pointwise bounded everywhere does not necessarily imply that B is nonmeager, as a topological space X needs not be nonmeager unless it is a Baire space. By Theorem 1.15, we have the following corollary:

Corollary 3.75. Let V be a Banach space, W a topological vector space and Γ be a collection of functions $V \rightarrow W$. If the sets

$$\{\Lambda v | \Lambda \in \Gamma\}$$

are bounded for all $v \in V$, then Γ is equicontinuous.

As another corollary, let us relate this to the operator norm:

Corollary 3.76. Let V and W be Banach spaces, Γ a collection of linear functions $V \rightarrow W$. If

$$\sup_{\Lambda \in \Gamma} \|\Lambda v\| < \infty$$

for all $v \in V$, then there exists M such that $\|\Lambda v\| \leq M\|v\|$ for all $v \in V, \Lambda \in \Gamma$

Proof. The theorem directly yields, for $\|v\| \leq 1$ and $\Lambda \in \Gamma$, a constant M such that

$$\|\Lambda v\| < M.$$

Linearity yields the relation

$$\|\Lambda v\| < M\|v\|$$

for all $v \in V$. □

3.5 Open Mapping & Closed Graph Theorems

All continuous bijections $\mathbb{R} \rightarrow \mathbb{R}$ are homeomorphisms, as they must either be increasing or decreasing. In general, it is not true that continuous bijections are homeomorphisms, as we shall see:

Example 3.77 (Continuous bijections are not homeomorphisms). The map $[0, 2\pi) \rightarrow S^1, t \mapsto e^{it}$ is continuous and bijective, but not a homeomorphism.

However, the situation for *linear* maps is much better.

Theorem 3.78. Let $\Phi : V \rightarrow W$ be a linear map between topological vector spaces. Then Φ is an open map iff for every zero neighbourhood $Z \subseteq V$ then $\Phi(Z) \subseteq W$ is a zero neighbourhood. It suffices to test this for a basis of zero neighbourhoods in V .

The condition above is also known as the property of being open at 0. Because of the translational invariance, if Φ is open at 0, it is also open at all points in v , and hence it is open. Open maps in Banach spaces are completely characterised through the following theorem

Theorem 3.79 (Open Mapping Theorem). Let $L : V \rightarrow W$ be a continuous linear transformation between Banach spaces V and W . Then the following are equivalent:

- (a) L is surjective.
- (b) L is open.
- (c) For every $w \in W$ there exists a solution $v \in V$ to the equation $Lv = w$.
- (d) There is a constant $C > 0$ such that for every $w \in W$ there is a solution $v \in V$ to the equation $Lv = w$ that satisfies $\|v\| \leq C\|w\|$.
- (e) There is a constant $C > 0$ and a dense subspace of vectors $P \subseteq W$ such that the previous condition applies to this dense subspace.

Proof. Clearly, (c) is equivalent to (a). (d) also clearly implies (c) and (e).

Now we show equivalence of (b) and (d). Suppose (b). Then we know that $L(B_1(0_V))$ contains an open ball $B_r(0_W)$. This then implies (d) by linearity. Conversely, we know that the image of the open ball with radius C includes the open ball of radius 1 around the origin, which shows openness at 0.

Now suppose (e). We show (d) by observing that we can approximate $w \in W$ arbitrarily well by vectors from P . Thus, we approximate w using a sequence w_n such that $\sum w_n = w$. For each w_n , we can solve the

equation to get v_n , with $\|v_n\| \leq C\|w_n\|$. By the triangle inequality and completeness of V , we have $\|v\| \leq C\|w\|$ and $Lv = w$.

Thus it only remains to show that (c) implies (d). For each $n \in \mathbb{N}$, we let $E_n \subseteq W$ be the subset of W such that there exists a solution to $Lv = w$ with $\|v\| \leq n\|w\|$. By (c), we know that the union of the E_n s is the entire space. Thus, the closure of some E_n contains a ball $B_r(w_0)$.

Then, the proof follows a sequence of approximations

1. Since E_n is dense in $B_r(w_0)$, we have *approximate solvability* — that is, for each $\epsilon > 0$ and $w \in B_r(w_0)$, we have v with $\|Lv - w\|_W \leq \epsilon$ and $\|v\|_V \leq n\|Lv\|_W$. Thus, $\|v\|_V \leq n\|w_0\| + nr + n\epsilon$.

2. By subtracting two such approximate solutions, we get:

For any $w \in B_{2r}(0)$ and $\epsilon > 0$, we have $v \in V$ with $\|Lv - w\|_W \leq 2\epsilon$ and $\|v\|_V \leq 2nr + 2n\epsilon$.

3. By rescaling L , we can conclude that for any $\epsilon > 0$, there exists $v \in V$ with $\|Lv - w\|_W \leq 2\epsilon$ and $\|v\|_V \leq 2n\|w\|_W + 2n\epsilon$.

4. In particular, setting $\epsilon = \frac{1}{4}\|w\|_W$, we conclude:

For any $w \in W$, we have $w = Lv + w'$, where $\|w'\|_W \leq \frac{1}{2}\|w\|_W$ and $\|v\|_V \leq \frac{5}{2}n\|w\|_W$.

5. By iterating this and using the completeness of V , we can find a solution that satisfies the bounds. \square

The key point of the theorem (and what is commonly known as the open mapping theorem) is that surjectivity is equivalent to openness. Note that the reverse direction is actually trivial, as the only open subspace of W is W itself.

As a corollary, we have

Corollary 3.80 (Inverse Mapping Theorem). A continuous bijective linear operator between Banach spaces has continuous inverse.

Another consequence of the open mapping theorem is a theorem on the equivalence of norms

Corollary 3.81. Suppose V is Banach with respect to the norms $\|\cdot\|_1$ and $\|\cdot\|_2$. If there exists C such that

$$\|v\|_1 \leq C\|v\|_2$$

for all $v \in V$, then the two norms are equivalent.

Proof. The inequality shows that the identity map is continuous. By the open mapping theorem, it is also open and thus a homeomorphism. \square

The next theorem follows almost trivially from the open mapping theorem:

Theorem 3.82 (Closed Graph Theorem). Let $L : V \rightarrow W$ be a linear map between Banach spaces. Then L is continuous iff its graph

$$\Gamma_L = \{(v, Lv) | v \in V\} \subseteq V \times W$$

is closed in the product topology.

Proof. Let f be a map $f : V \rightarrow W$. It factors as

$$V \xrightarrow{i} \Gamma_f \xrightarrow{\text{pr}_W} W$$

where i is the inverse of the projection map. The projection map is continuous. If we can apply the open mapping theorem to pr_V , then it is open and hence a homeomorphism. Thus, the inverse i is continuous, and f is continuous as a composition of continuous maps.

In particular, pr_V is always surjective. Since L is linear by hypothesis, pr_V is also linear, since $\text{pr}_V((v, Lv) + (w, Lw)) = v + w$. The last step we need is that Γ_L is a Banach space. This follows because it is a closed subspace of a Banach space.

The other direction of the proof is a well known property of topological spaces. If L is continuous (not even necessarily linear), $v_n \rightarrow v$ and $Lv_n \rightarrow w$, we have $L(\lim v_n) = \lim Lv_n \rightarrow w$, thus the graph is closed. \square

As somewhat of a converse of the inverse mapping theorem, we have the following:

Theorem 3.83. Let V be a normed space and W a Banach space. If there exists a bijective bounded linear operator $\phi : V \rightarrow W$, then V is Banach.

Proof. We construct a second norm on V $\|\cdot\|'_V$ using ϕ : We define $\|v\|'_V = \|\phi v\|_W$. Clearly, this is a norm. It is homogeneous, positive (since ϕ is bijective), and since ϕ is linear, we have

$$\|v + w\|'_V = \|\phi v + \phi w\|_W \leq \|\phi v\|_W + \|\phi w\|_W = \|v\|'_V + \|w\|'_V.$$

Now let us think about completeness relative to $\|\cdot\|'_V$. Consider a Cauchy sequence v_n with respect to $\|\cdot\|'_V$. Then ϕv_n is Cauchy in W , meaning it converges to some limit w . Thus, we have

$$\|v_n - \phi^{-1}w\|'_V = \|\phi v_n - w\|_W \rightarrow 0$$

showing that v_n converges to $\phi^{-1}w$. This means that V is complete with respect to $\|\cdot\|'_V$. Now we relate the two norms on V . Since ϕ is continuous with respect to $\|\cdot\|$, we have $\|v\|'_V = \|\phi v\|_W \leq C\|v\|_V$. TODO \square

Now, we show that the hypotheses of the open mapping theorem are all necessary. First, we construct an example with the domain being incomplete.

Example 3.84. There exists a surjective continuous linear function $\phi : V \rightarrow W$, with V incomplete, W complete that is not open.

Proof. Let $(W, \|\cdot\|)$ be any Banach space, and define f to be a discontinuous linear functional (see Theorem 3.15). Then, let V be the set W equipped with the norm $\|v\|_f = \|v\| + |f(v)|$.

By the triangle inequality on the real numbers, this is a norm. We now consider the identity map ι from $(W, \|\cdot\|_f)$ to $(W, \|\cdot\|)$. Since $\|v\| \leq \|v\|_f$, this map is bounded and hence continuous.

However, it cannot be open. Note that f is continuous with respect to $\|\cdot\|_f$, as $|f(v)| \leq \|v\|_f$. Thus, f is bounded as a linear map between $(W, \|\cdot\|_f)$ and \mathbb{R} and hence continuous.

If ι were open, then it would be a homeomorphism, and thus f would be continuous relative to $\|\cdot\|$ too, a contradiction. \square

In general, the above argument required the axiom of choice to prove existence of a Hamel basis. Here, we provide an example that does not

Example 3.85. Let V be $C([0, 1])$. The identity map $(V, \|\cdot\|_\infty) \rightarrow (V, \|\cdot\|_1)$ is not open.

Proof. We show that the image of the ∞ -ball with radius 1 is not open. In particular, we show that 0 is not in the interior of the ∞ -ball in the L^1 topology.

We do this by constructing a sequence of vectors that converge in L^1 to 0, but which are all outside the ∞ -ball. Intuitively, since the integral is a global average, we can construct continuous functions with spikes that get thinner, such that the integral converges to 0, while the peak remains high.

With this idea in mind, we construct

$$f_n(x) = \begin{cases} nx & 0 \leq x \leq \frac{1}{n} \\ 2 - nx & \frac{1}{n} < x \leq \frac{2}{n} \\ 0 & \text{otherwise.} \end{cases}$$

The graph of this function is an isosceles triangle between 0 and $2/n$ that peaks at height 1. Thus, it has ∞ norm 1, while it has L^1 norm equal to

$$\|f_n\|_1 = \frac{1}{n}.$$

Thus, $f_n \rightarrow 0$ in the L^1 norm, while lying in the exterior of the ∞ -ball with radius 1. □

Hilbert Spaces

4.1 Introduction

Definition 4.1 (Scalar Product). A scalar product in a vector space V over \mathbb{C} is a sesquilinear map $V \times V \rightarrow \mathbb{C}$, denoted by $\langle \cdot, \cdot \rangle$ such that

- (a) $\langle u, v + w \rangle = \langle u, v \rangle + \langle u, w \rangle$, $\langle u, aw \rangle = a\langle u, w \rangle$
- (b) $\langle u, v \rangle = \overline{\langle v, u \rangle}$
- (c) $\langle u, u \rangle > 0$ for $u \neq 0$.

Clearly, all inner products induce a norm through $\|v\| = \sqrt{\langle v, v \rangle}$. When we speak of the *topology* of a scalar product space, we will speak of the topology induced by this norm. The most important inequality is the Cauchy-Schwarz inequality:

Theorem 4.2 (Cauchy-Schwarz). A scalar product satisfies

$$|\langle u, v \rangle|^2 \leq \|u\|^2 \|v\|^2.$$

Proof. First, we use sesquilinearity to expand

$$\|u + tv\|^2 = \|u\|^2 + 2t \operatorname{Re} \langle u, v \rangle + t^2 \|v\|^2. \quad (4.1)$$

Then, we set $t = -\operatorname{Re} \langle u, v \rangle / \|v\|^2$ and multiply by $\|v\|^2$ to get

$$(\operatorname{Re} \langle u, v \rangle)^2 \leq \|u\|^2 \|v\|^2.$$

Rescaling u by a unit complex number to make $\langle u, v \rangle$ real, we get the desired inequality. \square


The next question is what kind of norms come from a scalar product. This question is answered using the *parallelogram law*

Proposition 4.3. Let $\|\cdot\|$ be a norm arising from a scalar product. Then

$$\|u + v\|^2 + \|u - v\|^2 = 2\|u\|^2 + 2\|v\|^2.$$

Proof. We set $t = \pm 1$ in Eq. (4.1) and sum up the two resulting equations. □

The geometric intuition behind this identity comes from the scalar product in \mathbb{R}^2 :



The construction of a scalar product from the norm then comes from the *polarization identity*, which we only discuss in the real case:

Theorem 4.4 (Polarization Identity). Suppose $\|\cdot\|$ is a norm satisfying the parallelogram law. Then

$$\begin{aligned} \langle u, v \rangle &= \frac{1}{4} (\|u + v\|^2 - \|u - v\|^2) \\ &= \frac{1}{2} (\|u + v\|^2 - \|u\|^2 - \|v\|^2) \\ &= \frac{1}{2} (\|u\|^2 + \|v\|^2 - \|u - v\|^2) \end{aligned}$$

defines a scalar product.

Proof. First, we note that all 3 forms are equivalent by the parallelogram law. Symmetry is clear. Positivity is clear by substituting $u = v$ into the first form. Finally, we show linearity. We attempt to equate $\langle u, v + w \rangle$ with $\langle u, v \rangle + \langle u, w \rangle$:

$$\|u + v + w\|^2 - \|u - v - w\|^2 \stackrel{?}{=} \|u + v\|^2 - \|u - v\|^2 + \|u + w\|^2 - \|u - w\|^2$$

□

Now that we understand how scalar products work, we move on to the theory of Hilbert spaces:

Definition 4.5 (Hilbert Space). A Hilbert space is a complete scalar product space.

Theorem 4.6 (Convex Minimizer). Given a nonempty closed, convex subset K of a Hilbert space \mathcal{H} and $x \in H$, there is a point y in K that has minimum distance to x .

Proof. First, we define the infimum distance

$$d = \inf_{y \in K} \|y - x\|.$$

We let y_n be a sequence that minimizes distance — that is,

$$\lim_{n \rightarrow \infty} d_n = d, \quad d_n = \|x - y_n\|$$

We will show that this sequence is Cauchy. Because K is closed, it will then converge in K . By continuity, it will converge to a minimum of distance.

We do this by applying the parallelogram law:

$$\left\| x - \frac{y_n + y_m}{2} \right\|^2 + \frac{1}{4} \|y_n - y_m\|^2 = \frac{1}{2} (d_n^2 + d_m^2).$$

Since K is convex, $(y_n + y_m)/2$ is in K , and thus

$$d \leq \left\| x - \frac{y_n + y_m}{2} \right\| < d + \epsilon.$$

Using this, we can bound $\|y_n - y_m\|$ and show that it is Cauchy. □

Definition 4.7. If W is a linear subspace, its orthogonal complement is the set of vectors orthogonal to W - that is,

$$W^\perp = \{v \in V \mid \langle v, w \rangle = 0 \ \forall w \in W\}.$$

We now ask the question of when we have the standard result in linear algebra - that is, when we can decompose a vector space as the direct sum of a subspace and its complement. It turns out that this is not true for all subspaces, as we now see

Theorem 4.8. Let \mathcal{H} be a Hilbert space, Y a closed subspace of \mathcal{H} , Y^\perp the orthogonal complement of Y . We claim that

- (a) Y^\perp is a closed subspace of Y ,
- (b) $Y \oplus Y^\perp = \mathcal{H}$,
- (c) $(Y^\perp)^\perp = Y$.

Proof. By the sesquilinearity of the scalar product, Y^\perp is a vector subspace. Now we show that it is closed. Since \mathcal{H} is Hilbert, being closed is equivalent to being complete.

Let $(v_n)_n \subseteq Y^\perp$ be convergent, $v_n \rightarrow v$. We have

$$\langle v, u \rangle = \langle v - v_n, u \rangle + \langle v_n, u \rangle = \langle v - v_n, u \rangle.$$

Then, by applying the Schwarz inequality, we have

$$\langle v, u \rangle \leq \|v - v_n\| \|u\|$$

for $u \in Y$. Since the first norm tends to 0, v is perpendicular to u for all $u \in Y$, or $v \in Y^\perp$.

The second part is done analogously to the finite dimensional case, but with much more machinery required. Consider $v \in \mathcal{H}$. By Theorem 4.6, we have a point u in Y closest to v . We show that $v - u \in Y^\perp$.

By the minimal property of u , we have, for any $y \in Y$,

$$\|v - u\|^2 \leq \|v - u + ty\|^2.$$

Defining $z = v - u$, we can rewrite the right hand side as

$$\|z + ty\|^2 = \|z\|^2 + 2t \operatorname{Re}\langle z, y \rangle + t^2 \|y\|^2.$$

Since this is minimised, we must have $\langle z, y \rangle = 0$.

The decomposition is unique, as if we had $v = y + z = y' + z'$, then $y - y' = z - z'$ would be orthogonal to each other, as they belong to Y and Y^\perp respectively, but that implies that they are 0.

The direct sum decomposition shows (c) directly. \square

We finally move on to what is the core theorem of scalar product spaces. In finite dimensional scalar product spaces, given a basis for a vector space, we can define a dual basis. This then yields an isomorphism between a vector space and its dual. In a scalar product space, this can be done without even the choice of a basis. We now seek to do the same. Before this proof, we first need a lemma:

Lemma 4.9. Let V be a Banach space.

- (a) The nullspace of a linear functional that is not identically 0 is a linear subspace of codimension 1.
- (b) If two linear functionals have the same nullspace, they are a constant multiple of each other.
- (c) The nullspace of a bounded linear functional is a closed subspace.

Proof. (a) Follows from the fundamental theorem of homomorphisms for groups:

$$V / \ker f \cong \mathbb{R}$$

which has dimension 1.

- (b) Let $v \in V$. We decompose v into the sum $v = u + w$, where u is in the (shared) kernel and w is not (this requires that the kernel is closed, see (c)).

Then, the two linear functionals are uniquely determined through their action on w . But since w is in a 1-dimensional subspace, it follows that the two functionals are a constant multiple of each other.

- (c) A bounded linear functional is continuous; the preimage of 0 is hence closed. The subspace property follows from linearity as it does in finite dimensions. \square

Now the important theorem follows immediately:

Theorem 4.10 (Riesz Representation Theorem). Let f be a bounded (and hence continuous) linear functional on a Hilbert space \mathcal{H} . Then f has the form

$$f = \langle v, \cdot \rangle.$$

The point v is uniquely determined.

Proof. The nullspace Y of f is a closed subspace of \mathcal{H} . Its orthogonal complement Y^\perp is one-dimensional. Let p be nonzero in Y^\perp , and define the linear functional m by $m = \langle p, \cdot \rangle$. This is a linear functional with the same nullspace as f , and by rescaling appropriately, we regain f . \square

Corollary 4.11. Every Hilbert space is reflexive.

A generalization is the Lax-Milgram Theorem

Theorem 4.12 (Lax-Milgram). Let \mathcal{H} be a Hilbert space and $B(u, v)$ be a function of two variables with the following properties:

(a) $B(x, y)$ is linear in y and skew-linear in x ,

(b) There exists c such that

$$|B(u, v)| \leq c\|u\|\|v\|,$$

(c) There exists $b > 0$ such that

$$|B(v, v)| \geq b\|v\|^2$$

for all $v \in \mathcal{H}$.

Then every bounded linear functional ℓ has the form

$$\ell(v) = B(u, v),$$

where u is a uniquely determined vector in \mathcal{H} .

Proof. Since $B(u, \cdot)$ is linear, we can write it as $\langle u', \cdot \rangle$ for some $u' \in \mathcal{H}$. Since u' is uniquely determined by u , it is a function of u . By the linearity of B , it must be a linear function of u . Thus, the set of possible u' is a linear subspace of \mathcal{H} . We claim that it is a closed subspace.

To do this, we consider the “norm”

$$B(u, u) = \langle u', u \rangle.$$

Then, we approximate their absolute values through the Cauchy-Schwarz inequality and condition (c):

$$b\|u\|^2 \leq \|u'\|\|u\|$$

or

$$b\|u\| \leq \|u'\|.$$

Note here again that u' is a linear function of u . Now we consider a sequence u'_n with corresponding u_n . We have, by skew-linearity,

$$B(u_n - u_m, v) = \langle u'_n - u'_m, v \rangle.$$

By the previous argument, this means that

$$b\|u_n - u_m\| \leq \|u'_n - u'_m\|.$$

This means that if the u'_n s form a Cauchy sequence, so do the u_n s. By completeness, this converges. Thus, if $u'_n \rightarrow u'$ and $u \rightarrow u'$, we have, by continuity both of B and of the scalar product,

$$B(u, v) = \langle u', v \rangle.$$

Thus, u' is still a candidate vector that may be produced by the Riesz Representation theorem, showing that the subspace is closed. Now, we show that this closed subspace is all of \mathcal{H} . Otherwise, there would be a vector w orthogonal to all u' . Then $B(u, w) = 0$ for all $u \in \mathcal{H}$. Letting $u = w$ yields $B(w, w) = 0$, or $w = 0$.

Now, we put the pieces of the puzzle in place. By the Riesz theorem, we know that

$$\ell = \langle u', \cdot \rangle$$

for some $u' \in \mathcal{H}$. By the previous argument, we know that

$$\langle u', \cdot \rangle = B(u, \cdot)$$

for some $u \in \mathcal{H}$. This completes the proof. \square

4.2 Lattices

The previous results about closed subspaces of Hilbert spaces can be generalized using the concept of a *lattice*:

Definition 4.13 (Lattice). A set V with two operations $\wedge, \vee : V \times V \rightarrow V$ is called a lattice if for all $a, b, c \in V$ we have

- (a) (Commutativity and associativity of \vee): $a \vee b = b \vee a$, $a \vee (b \vee c) = (a \vee b) \vee c$,
- (b) (Commutativity and associativity of \wedge): $a \wedge b = b \wedge a$, $a \wedge (b \wedge c) = (a \wedge b) \wedge c$,
- (c) (Symmetry) $a \vee a = a = a \wedge a$,
- (d) $(a \vee b) \wedge a = a = (a \wedge b) \vee a$.

The last part can be understood by thinking of \wedge as a logical AND and thinking of \vee as a logical OR. On a lattice, we can create a partial order:

Lemma 4.14. Let V be a lattice.

- (a) $a \vee b = b$ iff $a \wedge b = a$ for all $a, b \in V$.
- (b) We define $a \leq b$ if $a \vee b = b$. This turns V into a partially ordered set.
- (c) The partial order admits infimums and supremums for two elements $a, b \in V$, given by

$$\inf(a, b) = a \wedge b, \quad \sup(a, b) = a \vee b.$$

Proof. (a) Suppose $a \vee b = b$. Then

$$(a \vee b) \wedge a = b \wedge a = a \wedge b = a$$

by (d) of the definition. Conversely, if $a \wedge b = a$, then

$$(b \wedge a) \vee b = a \vee b = b.$$

- (b) We verify the three properties

- (1) (Reflexivity) Since $a \vee a = a$, we have $a \leq a$.
- (2) (Symmetry) Suppose $a \vee b = b$, and $b \vee a = a$. Then $a = b$ by commutativity.
- (3) (Transitivity) Suppose $a \vee b = b$, and $b \vee c = c$. Then

$$a \vee c = (a \wedge b) \vee c = (a \wedge b \wedge c) \vee c = [c \wedge (a \wedge b)] \vee c = c.$$

- (c) First, we show that these are actually lower and upper bounds respectively. We have

$$(a \wedge b) \vee a = a, \quad (a \wedge b) \vee b = b$$

directly by definition. Similarly,

$$a \vee (a \vee b) = (a \vee a) \vee b = a \vee b, \quad b \vee (a \vee b) = (b \vee b) \vee a = a \vee b.$$

Then, we show that these are minimal and maximal lower and upper bounds. □

4.3 Bases

As one might expect, the next step up from a (Schauder) basis in a Banach space would be an orthonormal basis:

Definition 4.15 (Orthonormal Basis). An orthonormal (Schauder) basis $\{v_n\}$ is a set of vectors whose closed linear span is the entire space, such that $\langle v_i, v_j \rangle = 0$ for $i \neq j$, and $\|v_n\| = 1$.

Such bases always exist, by the classic Zorn's Lemma argument

Theorem 4.16. Every Hilbert space has an orthonormal Schauder basis.

Proof. Consider all orthonormal sets, partially ordered by inclusion. By Zorn's Lemma, there is a maximal orthonormal set. □

As in finite dimensions, we have the Gram-Schmidt procedure for orthogonalizing a finite set of vectors:

Theorem 4.17 (Gram-Schmidt). Let $\{u_n\}$ be a basis for the Hilbert space \mathcal{H} . Then there exists an orthonormal basis $\{v_n\}$ such that the span of v_1, \dots, v_n is the span of u_1, \dots, u_n .

4.4 Applications

As a first application, we prove the Radon-Nikodym theorem as a consequence of the Riesz Representation theorem. First, we need to define a few terms

Definition 4.18 (Absolute Continuity). Let μ and ν be measures defined on the same measurable space (X, \mathcal{A}) . We say that ν is absolutely continuous with respect to μ , or $\nu \ll \mu$, if for all $A \in \mathcal{A}$, $\mu(A) = 0 \implies \nu(A) = 0$.

Before we move on to the full Radon-Nikodym theorem, we prove a slightly weaker version:

Theorem 4.19 (Finite Radon-Nikodym). Let μ and ν be finite measures defined on the same measurable space (X, \mathcal{A}) such that $\nu \ll \mu$. Then there exists measurable $g : X \rightarrow \mathbb{R}$ such that

$$\nu(A) = \int_A g \, d\mu$$

for all $A \in \mathcal{A}$.

Proof. We consider the Hilbert space $L^2(\mu + \nu)$. Since both measures are finite, the constant 1-function is integrable, and hence by the Schwarz inequality we have

$$\left| \int v \, d(\mu + \nu) \right|^2 \leq \left(\int v^2 \, d(\mu + \nu) \right) \left(\int 1 \, d(\mu + \nu) \right).$$

This shows that all square integrable functions are integrable. Then, we consider the linear functional

$$\ell(v) = \int v \, d\mu$$

which is bounded with respect to the $L^2(\mu)$ norm, and thus is even more strongly bounded with respect to the $L^2(\mu + \nu)$ norm. Thus, by the Riesz representation theorem we have

$$\ell(v) = \int v \, d\mu = \int vu \, d(\mu + \nu)$$

for a unique $u \in L^2(\mu + \nu)$. This can be rewritten as

$$\int v(1 - u) \, d\mu = \int vu \, d\nu. \quad (4.2)$$

Now we show that $0 < u \leq 1$ μ -almost everywhere.

We begin by considering the set $F = \{x \in X | u \leq 0\}$. As the inverse image of a measurable set under a measurable function, it is itself measurable. Then, by letting v be its characteristic function (which is L^2), Equation (4.2) becomes

$$\int_F (1 - u) \, d\mu = \int u \, d\nu.$$

Since $u \leq 0$ on F , the right hand side is ≤ 0 , while the left hand side is $\geq \mu(F)$. Thus, we must have $\mu(F) = 0$.

Similarly, we consider $F = \{x \in X | u > 1\}$. Here, we have

$$\int (1 - u) \, d\mu \leq 0, \quad \int u \, d\nu \geq \nu(F).$$

The only possibility for equality happens when F is a μ -null set. Thus, we can modify u on a μ -null set such that $0 < u \leq 1$ everywhere.

Finally, we claim that

$$g = \frac{1 - u}{u}.$$

To see this, we write $p = uv$ and rewrite Equation (4.2) as

$$\int pg \, d\mu = \int p \, d\nu.$$

We now choose v such that p becomes the characteristic function of a set A . Then, we have

$$\nu(A) = \int g \, d\mu$$

as desired. □

Now, we prove the full Radon-Nikodym theorem. This is a purely measure theoretic exercise, with no need to resort to functional analysis. As one might expect, it proceeds by cutting up the σ -finite space into finite spaces, applying the finite version of the Radon-Nikodym theorem, and pasting them together using characteristic functions.

Theorem 4.20 (Radon-Nikodym). Let μ and ν be σ -finite measures defined on the same measurable space (X, \mathcal{A}) such that $\nu \ll \mu$. Then there exists a measurable function $g : X \rightarrow \mathbb{R}$ such that

$$\nu(A) = \int_A g \, d\mu$$

for all $A \in \mathcal{A}$.

Proof. Since X is σ -finite, we have $X = \bigcup_{n=1}^{\infty} A_n$, with $\mu(A_n) < \infty$, and the A_n s pairwise disjoint. Then, we can apply the finite version of the Radon-Nikodym theorem to find $g_n : A_n \rightarrow \mathbb{R}$ such that

$$\nu(E \cap A_n) = \int_{A_n} g_n \, d\mu$$

for measurable E . Then we let $g = \sum_{n=1}^{\infty} g_n \chi_{A_n}$. By the monotone convergence theorem, this is also positive and measurable. Then

$$\nu(E) = \sum_{n=1}^{\infty} \nu(E \cap A_n) = \sum_{n=1}^{\infty} \int_{A_n} g \, d\mu = \int g \, d\mu. \quad \square$$

List of Definitions

1.2	Definition (Preorder)	4
1.6	Definition (Convergence)	4
1.10	Definition (Nowhere Dense)	5
1.12	Definition (Meager)	5
1.14	Definition (Baire Spaces)	6
2.9	Definition (Uniform Continuity)	10
2.11	Definition (Algebraic and Topological Dual)	10
3.1	Definition (Norm)	12
3.3	Definition (Banach Space)	12
3.4	Definition (Seminorm)	12
3.9	Definition (Operator Norm)	13
3.12	Definition (Hamel Basis)	14
3.16	Definition (Schauder Basis)	15
3.32	Definition (Minkowski Functional)	20
3.72	Definition (Equicontinuity)	30
4.1	Definition (Scalar Product)	36
4.5	Definition (Hilbert Space)	37
4.13	Definition (Lattice)	41
4.15	Definition (Orthonormal Basis)	42
4.18	Definition (Absolute Continuity)	42

List of Theorems

1.1	Theorem (List of Useful Identities)	3
1.15	Theorem (Baire I)	6
1.16	Theorem (Baire II)	7
2.10	Theorem (Equivalence of Continuity Conditions)	10
3.5	Theorem (Quotient of Norms)	12
3.6	Theorem (Operator Norm)	13
3.11	Theorem (Operator Composition)	14
3.15	Theorem (Existence of Discontinuous Functionals)	15
3.24	Theorem (Auxiliary Functionals)	17
3.27	Theorem (Dominated Extension)	18
3.28	Theorem (Hahn-Banach (Normed))	19
3.33	Theorem (Separation I)	20
3.34	Theorem (Separation II)	21
3.35	Theorem (Separation III)	21
3.43	Theorem (Completion of normed space)	24
3.54	Theorem (Polars of Hulls)	26
3.55	Theorem (Bipolar Theorem)	27
3.56	Theorem (Banach-Alaoglu)	27
3.59	Theorem (Sequential Compactness of Unit Ball)	28
3.63	Theorem (Spanning Criterion)	28
3.74	Theorem (Banach-Steinhaus)	31
3.79	Theorem (Open Mapping Theorem)	32
3.82	Theorem (Closed Graph Theorem)	33
4.2	Theorem (Cauchy-Schwarz)	36
4.4	Theorem (Polarization Identity)	37
4.6	Theorem (Convex Minimizer)	37
4.10	Theorem (Riesz Representation Theorem)	39
4.12	Theorem (Lax-Milgram)	40
4.17	Theorem (Gram-Schmidt)	42
4.19	Theorem (Finite Radon-Nikodym)	43
4.20	Theorem (Radon-Nikodym)	44