Tanmay Garg

CS20BTECH11063


Data Science Analysis Assignment 1

Q1

```
In [14]: import numpy as np
         import matplotlib.pyplot as plt
         import scipy.stats as stats
          import astroML
         from astroML.stats import sigmaG
         mu_gauss = 1.5
          sigma gauss = 0.5
          # generate 1000 random numbers from a Gaussian distribution with mean of 1.5 and standard deviation of 0.5
          # generated samples = np.random.normal(mu gauss, sigma gauss, 1000)
         normal distribution = stats.norm(mu gauss, sigma gauss)
          sampled_points = normal_distribution.rvs(1000)
          generated_samples = sampled_points
         y sampled points = normal distribution.pdf(sampled points)
          plt.plot(sampled points, y sampled points, '.')
         plt.xlabel('x')
          plt.ylabel('y')
         plt.title('Sampled points from a Gaussian distribution')
         plt.hist(generated_samples, bins=19, density=True)
          plt.show()
          # sample mean
          sample mean = np.mean(generated samples)
          print('sample mean: {}'.format(sample mean))
          # sample variance
          sample variance = np.var(generated samples)
          print('sample variance: {}'.format(sample variance))
          # sample skewness
```

```
sample skewness = stats.skew(generated samples)
print('sample skewness: {}'.format(sample skewness))
# sample kurtosis
sample kurtosis = stats.kurtosis(generated samples)
print('sample kurtosis: {}'.format(sample kurtosis))
# sample standard deviation using MAD
sample std mad = 1.4826 * np.median(np.abs(generated samples - np.median(generated samples)))
print('sample standard deviation using MAD: {}'.format(sample std mad))
# sample standard deviation using sigma G
sample std sigma g = sigmaG(generated samples)
print('sample standard deviation using sigma G: {}'.format(sample std sigma g))
# sample standard deviation using sigma G formula
q25 = np.percentile(generated samples, 25)
q75 = np.percentile(generated_samples, 75)
sample_std_sigma_g_form = 0.7413*(q75 - q25)
print('sample standard deviation using sigma G formula: {}'.format(sample std sigma g form))
# Plot pdf of the Gaussian distribution
x = np.arange(0, 5, 0.01)
y = stats.norm.pdf(x, mu_gauss, sigma_gauss)
plt.plot(x, y)
plt.xlabel('x')
plt.ylabel('y')
plt.title('Gaussian distribution')
plt.show()
```

Sampled points from a Gaussian distribution




sample mean: 1.459868961658885

sample variance: 0.25674636600112427
sample skewness: -0.24203944301296926
sample kurtosis: 0.02638822955238984

sample standard deviation using MAD: 0.5096849842543005 sample standard deviation using sigma G: 0.509207380726251

sample standard deviation using sigma G formula: 0.5092066187690567



Q2

```
In [9]: import numpy as np
import matplotlib.pyplot as plt
import scipy.stats as stats


mu_cauchy = 0
gamma_cauchy = 1.5
x = np.arange(-10, 10, 0.01)
y_cauchy = stats.cauchy.pdf(x, mu_cauchy, gamma_cauchy)

mu_gauss = 0
sigma_gauss = 1.5
y_gauss = stats.norm.pdf(x, mu_gauss, sigma_gauss)

# Plot the two distributions
```

```
plt.plot(x, y_cauchy, label='Cauchy')
plt.plot(x, y_gauss, label='Gaussian')
plt.legend()
plt.xlabel('x')
plt.ylabel('pdf')
plt.title('Cauchy vs Gaussian distributions')
plt.show()
```

Cauchy vs Gaussian distributions



Q3

```
import numpy as np
import matplotlib.pyplot as plt
import scipy.stats as stats


x = np.arange(0, 10, 0.01)
```

```
mu_gauss = 5
sigma_gauss = np.sqrt(5)
y_gauss = stats.norm.pdf(x, mu_gauss, sigma_gauss)

mu_poisson = 5
y_poisson = stats.poisson.pmf([0,1,2,3,4,5,6,7,8,9,10], mu_poisson)

# Plot the two distributions
plt.step([0,1,2,3,4,5,6,7,8,9,10], y_poisson, label='Poisson')
plt.plot(x, y_gauss, label='Gaussian')
plt.legend()
plt.xlabel('x')
plt.ylabel('pdf')
plt.title('Poisson vs Gaussian distributions')
plt.show()
```

Poisson vs Gaussian distributions



```
In [11]: measure_val = [0.8920, 0.881, 0.8913, 0.9837, 0.8958]
    measured_uncertainty = [0.00044, 0.009, 0.00032, 0.00048, 0.00045]
    measure_val = np.array(measure_val)
    measured_uncertainty = np.array(measured_uncertainty)

weighted_mean = np.average(measure_val, weights=1/(measured_uncertainty**2))
    print('weighted mean: {}'.format(weighted_mean))
    weighted_uncertainty = np.sqrt(1/np.sum(1/(measured_uncertainty**2)))
    print('weighted uncertainty: {}'.format(weighted_uncertainty))

weighted mean: 0.9089185199574896
```

Q5

weighted uncertainty: 0.00020318737026848627

```
In [12]: import numpy as np
         import matplotlib.pyplot as plt
         import scipy.stats as stats
         import pandas as pd
         # read the data from the csv file
         data = pd.read_csv('exoplanet.eu_catalog.csv')
         tmp dat = data
         data = data['eccentricity'].dropna()
         print(len(data))
         # print only negative values of eccentricity
         # print(data[data < 0])</pre>
         # histogram of the eccentricity data
         plt.hist(data, bins=20)
         plt.xlabel('eccentricity')
         plt.title('Eccentricity data')
         plt.ylabel('frequency')
         plt.show()
         # gaussianize the eccentricity data
         data_gaussianized = stats.boxcox(data[data>0])[0]
         # histogram of the gaussianized eccentricity data
         plt.hist(data_gaussianized, bins=19)
         plt.xlabel('gaussianized eccentricity')
         plt.ylabel('frequency')
```

```
plt.title('Gaussianized eccentricity data')
plt.show()
```

2144

