1

Assignment 4

Tanmay Goyal - AI20BTECH11021

Download all python codes from

https://github.com/tanmaygoyal258/EE3900-Linear -Systems-and-Signal-processing/blob/main/ Assignment4/code.py

Download all latex codes from

https://github.com/tanmaygoyal258/EE3900-Linear -Systems-and-Signal-processing/blob/main/ Assignment4/main.tex

1 Problem

(Linear_Forms/Q.2.15) Find the equation of the line equidistant from parallel lines

$$(9 \ 6) \mathbf{x} = 7 \tag{1.0.1}$$

$$(3 \ 2)\mathbf{x} = -6$$
 (1.0.2)

2 Solution

In general, we can obtain the following lemma:

Lemma 2.1. Given the two following parallel lines:

$$a\mathbf{n}^T\mathbf{x} - c_1 = 0 \tag{2.0.1}$$

$$b\mathbf{n}^T\mathbf{x} - c_2 = 0 \tag{2.0.2}$$

The line equidistant from both parallel lines would be given by:

$$\mathbf{n}^{T}\mathbf{x} - \frac{1}{2} \left(\frac{c_1}{a} + \frac{c_2}{b} \right) = 0$$
 (2.0.3)

Proof. The distance between a point **A** and a line $L = \mathbf{n}^T \mathbf{x} - c$ is given by:

$$\|\mathbf{P} - \mathbf{A}\| = \frac{\left|\mathbf{n}^T \mathbf{A} - c\right|}{\|\mathbf{n}\|}$$
 (2.0.4)

where \mathbf{P} is the foot of perpendicular from \mathbf{A} onto \mathbf{L} .

Consider a point \mathbf{x} equidistant from both parallel lines, then:

$$\frac{\left|a\mathbf{n}^{T}\mathbf{x}-c_{1}\right|}{\left|\left|a\mathbf{n}\right|\right|}=\frac{\left|b\mathbf{n}^{T}\mathbf{x}-c_{2}\right|}{\left|\left|b\mathbf{n}\right|\right|}$$
(2.0.5)

$$\frac{\left|a\mathbf{n}^{T}\mathbf{x}-c_{1}\right|}{\left|a\right|}=\frac{\left|b\mathbf{n}^{T}\mathbf{x}-c_{2}\right|}{\left|b\right|}$$
(2.0.6)

$$\left| ab\mathbf{n}^T \mathbf{x} - bc_1 \right| = \left| ab\mathbf{n}^T \mathbf{x} - ac_2 \right| \tag{2.0.7}$$

$$2ab\mathbf{n}^{T}\mathbf{x} - bc_1 - ac_2 = 0 (2.0.8)$$

$$\mathbf{n}^{T}\mathbf{x} - \frac{1}{2} \left(\frac{c_1}{a} + \frac{c_2}{b} \right) = 0$$
 (2.0.9)

The two given parallel lines can be written as:

$$3(3 2)\mathbf{x} - 7 = 0 (2.0.10)$$

$$(3 2)\mathbf{x} + 6 = 0 (2.0.11)$$

On comparing the equations with (2.0.2),

$$\mathbf{n} = \begin{pmatrix} 3 & 2 \end{pmatrix} \tag{2.0.12}$$

$$a = 3$$
 (2.0.13)

$$b = 1 (2.0.14)$$

$$c_1 = 7 (2.0.15)$$

$$c_2 = -6 (2.0.16)$$

On substituting these values into (2.0.9),

$$(3 2)\mathbf{x} - \frac{1}{2}(\frac{7}{3} - 6) = 0 (2.0.17)$$

$$(3 \quad 2)\mathbf{x} - \frac{11}{6} = 0 \tag{2.0.18}$$




Fig. 0: The equidistant line