

10/04/2019  
Evening

**Answers & Solutions**  
*for*  
**JEE (MAIN)-2019 (Online) Phase-2**  
(Physics, Chemistry and Mathematics)

Time : 3 hrs.

M.M. : 360

**Important Instructions :**

1. The test is of **3 hours** duration.
2. The Test Booklet consists of **90** questions. The maximum marks are **360**.
3. There are **three** parts in the question paper A, B, C consisting of **Physics, Chemistry and Mathematics** having 30 questions in each part of equal weightage.
4. Each question is allotted 4 (**four**) marks for each correct response.  $\frac{1}{4}$  (*one-fourth*) marks will be deducted for indicating incorrect response of each question. No deduction from the total score will be made if no response is indicated for an item in the answer sheet.
5. There is only one correct response for each question.

## PART-A : PHYSICS

1. One mole of an ideal gas passes through a process where pressure and volume obey the

relation  $P = P_0 \left[ 1 - \frac{1}{2} \left( \frac{V_0}{V} \right)^2 \right]$ . Here  $P_0$  and  $V_0$  are

constants. Calculate the change in the temperature of the gas if its volume changes from  $V_0$  to  $2V_0$ .

- (1)  $\frac{1}{4} \frac{P_0 V_0}{R}$  (2)  $\frac{5}{4} \frac{P_0 V_0}{R}$   
 (3)  $\frac{1}{2} \frac{P_0 V_0}{R}$  (4)  $\frac{3}{4} \frac{P_0 V_0}{R}$

Answer (2)

Sol. If  $V_1 = V_0 \Rightarrow P_1 = P_0 \left[ 1 - \frac{1}{2} \right] = \frac{P_0}{2}$


If  $V_2 = 2V_0 \Rightarrow P_2 = P_0 \left[ 1 - \frac{1}{2} \left( \frac{1}{4} \right) \right] = \left( \frac{7P_0}{8} \right)$

$$\left( T = \frac{PV}{nR} \right) \Rightarrow \Delta T = \left| \frac{P_1 V_1}{nR} - \frac{P_2 V_2}{nR} \right|$$

$$\Delta T = \left| \left( \frac{1}{nR} \right) (P_1 V_1 - P_2 V_2) \right| = \left( \frac{1}{nR} \right) \left( \frac{P_0 V_0}{2} - \frac{7P_0 V_0}{4} \right)$$

$$= \frac{5P_0 V_0}{4nR}$$

2. The correct figure that shows, schematically, the wave pattern produced by superposition of two waves of frequencies 9 Hz and 11 Hz, is :



Answer (3)

Sol. Beat frequency =  $|f_1 - f_2| = 11 - 9 = 2$  Hz

3. A spaceship orbits around a planet at a height of 20 km from its surface. Assuming that only gravitational field of the planet acts on the spaceship, what will be the number of complete revolutions made by the spaceship in 24 hours around the planet?

[Given: Mass of planet =  $8 \times 10^{22}$  kg,

Radius of planet =  $2 \times 10^6$  m,

Gravitational constant  $G = 6.67 \times 10^{-11} \text{ Nm}^2/\text{kg}^2$ ]

- (1) 17 (2) 13  
 (3) 11 (4) 9

Answer (3)

Sol.  $T = \frac{2\pi r}{v}$ ,  $v = \sqrt{\frac{GM}{r}}$

$$T = 2\pi r \sqrt{\frac{r}{GM}} = 2\pi \sqrt{\frac{r^3}{GM}}$$


$$T = 2\pi \sqrt{\frac{(202)^3 \times 10^{12}}{6.67 \times 10^{-11} \times 8 \times 10^{22}}} \text{ sec}$$

$$T = 7812.2 \text{ s}$$

$$T \approx 2.17 \text{ hr} \Rightarrow 11 \text{ revolutions}$$

4. A plane is inclined at an angle  $\alpha = 30^\circ$  with respect to the horizontal. A particle is projected with a speed  $u = 2 \text{ ms}^{-1}$ , from the base of the plane, making an angle  $\theta = 15^\circ$  with respect to the plane as shown in the figure. The distance from the base, at which the particle hits the plane is close to :

(Take  $g = 10 \text{ ms}^{-2}$ )




- (1) 18 cm (2) 20 cm  
 (3) 14 cm (4) 26 cm

Answer (2)

Sol. Time of flight (T) =  $\frac{2u \sin \alpha}{g \cos \beta}$

$$T = \frac{(2)(2 \sin 15)}{g \cos 30} = \frac{4 \sin 15}{10 \cos 30}$$



$$\begin{aligned}\text{Range (R)} &= (2 \cos 15)T - \frac{1}{2}g \sin 30(T)^2 \\ &= (2 \cos 15) \frac{4 \sin 15}{10 \cos 30} - \left( \frac{1}{3} \times 10 \sin 30 \right) \frac{16 \sin^2 15}{100 \cos^2 30} \\ &= \frac{16\sqrt{3} - 16}{60} \approx 0.1952 \text{ m} \approx 20 \text{ cm}\end{aligned}$$

5. A bullet of mass 20 g has an initial speed of  $1 \text{ ms}^{-1}$ , just before it starts penetrating a mud wall of thickness 20 cm. If the wall offers a mean resistance of  $2.5 \times 10^{-2} \text{ N}$ , the speed of the bullet after emerging from the other side of the wall is close to :

- (1)  $0.4 \text{ ms}^{-1}$  (2)  $0.7 \text{ ms}^{-1}$   
(3)  $0.3 \text{ ms}^{-1}$  (4)  $0.1 \text{ ms}^{-1}$

Answer (2)

Sol.  $v^2 = u^2 - 2aS$

$$v^2 = (1)^2 - (2) \left[ \frac{2.5 \times 10^{-2}}{20 \times 10^{-3}} \right] \frac{20}{100}$$

$$v^2 = 1 - \frac{1}{2}$$

$$\Rightarrow v = \frac{1}{\sqrt{2}} \text{ m/s} = 0.7 \text{ m/s}$$

6. The time dependence of the position of a particle of mass  $m = 2$  is given by  $\vec{r}(t) = 2t\hat{i} - 3t^2\hat{j}$ . Its angular momentum, with respect to the origin, at time  $t = 2$  is :

- (1)  $-34(\hat{k} - \hat{i})$  (2)  $48(\hat{i} + \hat{j})$   
(3)  $36\hat{k}$  (4)  $-48\hat{k}$

Answer (4)

Sol.  $\vec{r} = 2t\hat{i} - 3t^2\hat{j}$

$$\vec{v} = \frac{d\vec{r}}{dt} = 2\hat{i} - 6t\hat{j}$$


$$\vec{a} = \frac{d\vec{v}}{dt} = -6\hat{j}$$

$$\vec{F} = m\vec{a} = -12\hat{j}$$

$$\vec{r}(\text{at } t = 2) = 4\hat{i} - 12\hat{j}$$


$$\vec{L} = m(\vec{r} \times \vec{v}) = 2(4\hat{i} - 12\hat{j}) \times (2\hat{i} - 12\hat{j}) = -48\hat{k}$$

7. The figure represents a voltage regulator circuit using a Zener diode. The breakdown voltage of the Zener diode is 6 V and the load resistance is,  $R_L = 4 \text{ k}\Omega$ . The series resistance of the circuit is  $R_i = 1 \text{ k}\Omega$ . If the battery voltage  $V_B$  varies from 8 V to 16 V, what are the minimum and maximum values of the current through Zener diode?



- (1) 0.5 mA; 6 mA (2) 0.5 mA; 8.5 mA  
(3) 1.5 mA; 8.5 mA (4) 1 mA; 8.5 mA


Answer (2)



$$I_1 = \left( 8 - 6 - \frac{3}{2} \right) = \frac{1}{2} = 0.5 \text{ mA}$$

$$I_2 = \left( 16 - 6 - \frac{3}{2} \right) = 8.5 \text{ mA}$$

8. A metal coin of mass 5 g and radius 1 cm is fixed to a thin stick AB of negligible mass as shown in the figure. The system is initially at rest. The constant torque, that will make the system rotate about AB at 25 rotations per second in 5 s, is close to :



- (1)  $4.0 \times 10^{-6} \text{ Nm}$  (2)  $7.9 \times 10^{-6} \text{ Nm}$   
(3)  $2.0 \times 10^{-5} \text{ Nm}$  (4)  $1.6 \times 10^{-5} \text{ Nm}$

Answer (3)

**Sol.**  $\tau = I\alpha$

$$\omega = \omega_0 + \alpha t$$

$$\Rightarrow 25 \times 2\pi = (\alpha)5$$

$$\alpha = 10\pi$$

$$\Rightarrow \tau = \left(\frac{5}{4} m R^2\right) \alpha$$

$$\approx \left(\frac{5}{4}\right) (5 \times 10^{-3}) (10^{-4}) 10\pi$$

$$= 2.0 \times 10^{-5} \text{ Nm}$$

9. In a Young's double slit experiment, the ratio of the slit's width is 4 : 1. The ratio of the intensity of maxima to minima, close to the central fringe on the screen, will be :

(1)  $(\sqrt{3} + 1)^4 : 16$       (2) 4 : 1

(3) 25 : 9      (4) 9 : 1

**Answer (4)**

**Sol.**  $I_1 = 4 I_0$

$$I_2 = I_0$$

$$\frac{I_{\max}}{I_{\min}} = \frac{(\sqrt{I_1} + \sqrt{I_2})^2}{(\sqrt{I_1} - \sqrt{I_2})^2} = \left(\frac{9}{1}\right)$$

10. Two radioactive substances A and B have decay constants  $5\lambda$  and  $\lambda$  respectively. At  $t = 0$ , a sample has the same number of the two nuclei. The time taken for the ratio of the

number of nuclei to become  $\left(\frac{1}{e}\right)^2$  will be :

(1)  $\frac{1}{2\lambda}$       (2)  $\frac{1}{4\lambda}$

(3)  $\frac{1}{\lambda}$       (4)  $\frac{2}{\lambda}$

**Answer (1)**

**Sol.**  $N_x(\text{at } t) = N_0 e^{-5\lambda t}$

$$N_y(\text{at } t) = N_0 e^{-\lambda t}$$

$$\frac{N_x}{N_y} = \frac{1}{e^2} = e^{-4\lambda t}$$

$$\Rightarrow 4\lambda t = 2$$

$$\Rightarrow t = \frac{2}{4\lambda} = \left(\frac{1}{2\lambda}\right)$$

11. A cubical block of side 0.5 m floats on water with 30% of its volume under water. What is the maximum weight that can be put on the block without fully submerging it under water?

[Take, density of water =  $10^3 \text{ kg/m}^3$ ]

(1) 30.1 kg      (2) 46.3 kg

(3) 87.5 kg      (4) 65.4 kg

**Answer (3)**

**Sol.** Given  $(50)^3 \times \frac{30}{100} \times (1) \times g = M_{\text{cube}} g \quad \dots(i)$

Let  $m$  mass should be placed


Hence  $(50)^3 \times (1) \times g = (M_{\text{cube}} + m) g \quad \dots(ii)$

equation (ii) – equation (i)

$$\Rightarrow mg = (50)^3 \times g(1 - 0.3) = 125 \times 0.7 \times 10^3 g$$

$$\Rightarrow m = 87.5 \text{ kg}$$

12. The graph shows how the magnification  $m$  produced by a thin lens varies with image distance  $v$ . What is the focal length of the lens used?



(1)  $\frac{b^2}{ac}$

(2)  $\frac{b^2 c}{a}$

(3)  $\frac{a}{c}$

(4)  $\frac{b}{c}$

**Answer (4)**

**Sol.** As the graph between magnification ( $m$ ) and image distance ( $v$ ) varies linearly, then

$$m = k_1 v + k_2$$

$$\Rightarrow \frac{v}{u} = k_1 v + k_2$$

$$\Rightarrow \frac{1}{u} = k_1 + \frac{k_2}{v}$$

$$\Rightarrow \frac{k_2}{v} - \frac{1}{u} = k_1$$

Clearly,  $k_1 = \frac{1}{f}$  and  $k_2 = 1$  here

$$\therefore f = \frac{1}{\text{slope of } m-v \text{ graph}} = \frac{b}{c}$$

13. In  $\text{Li}^{++}$ , electron in first Bohr orbit is excited to a level by a radiation of wavelength  $\lambda$ . When the ion gets deexcited to the ground state in all possible ways (including intermediate emissions), a total of six spectral lines are observed. What is the value of  $\lambda$ ?

(Given:  $h = 6.63 \times 10^{-34} \text{ Js}$ ;  $c = 3 \times 10^8 \text{ ms}^{-1}$ )

- (1) 11.4 nm  
(2) 12.3 nm  
(3) 9.4 nm  
(4) 10.8 nm

Answer (4)


Sol.  $\Delta E = \frac{hc}{\lambda}$

$$= (13.4)(3)^2 \left[ 1 - \frac{1}{16} \right] \text{ eV}$$

$$\Rightarrow \lambda = \frac{1242 \times 16}{(13.4) \times (9)(15)} \text{ nm} \approx 10.8 \text{ nm}$$

14. Two blocks A and B of masses  $m_A = 1 \text{ kg}$  and  $m_B = 3 \text{ kg}$  are kept on the table as shown in figure. The coefficient of friction between A and B is 0.2 and between B and the surface of the table is also 0.2. The maximum force  $F$  that can be applied on B horizontally, so that the block A does not slide over the block B is:

[Take  $g = 10 \text{ m/s}^2$ ]



- (1) 40 N  
(2) 12 N  
(3) 16 N  
(4) 8 N

Answer (3)

Sol.  $a_c = \left( \frac{F - f}{M + m} \right)$

$$a = \frac{F - (0.2)4 \times 10}{4} = \left( \frac{F - 8}{4} \right)$$

We have  $\frac{F - 8}{4} \leq (0.2)10$

$$\Rightarrow F - 8 \leq 8$$


$$\Rightarrow F \leq 16$$

15. In free space, a particle A of charge  $1 \mu\text{C}$  is held fixed at a point P. Another particle B of the same charge and mass  $4 \mu\text{g}$  is kept at a distance of 1 mm from P. If B is released, then its velocity at a distance of 9 mm from P is :

$$\left[ \text{Take } \frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ Nm}^2\text{C}^{-2} \right]$$

- (1)  $2.0 \times 10^3 \text{ m/s}$  (2)  $3.0 \times 10^4 \text{ m/s}$   
(3)  $1.5 \times 10^2 \text{ m/s}$  (4)  $1.0 \text{ m/s}$

Answer (Bonus)



Conservation of energy

$$\frac{kq_1q_2}{r_1} = \frac{kq_1q_2}{r_2} + \frac{1}{2}mv^2$$

$$v^2 = \frac{2kq_1q_2}{m} \left[ \frac{1}{r_1} - \frac{1}{r_2} \right]$$

$$= \frac{2 \times 9 \times 10^9 \times 10^{-12}}{4 \times 10^{-9} \times 10^{-3}} \left[ 1 - \frac{1}{9} \right] = 4 \times 10^9$$

$$v = \sqrt{40} \times 10^4 \text{ m/s} = 6.32 \times 10^4 \text{ m/s}$$

16. Water from a tap emerges vertically downwards with an initial speed of  $1.0 \text{ ms}^{-1}$ . The cross-sectional area of the tap is  $10^{-4} \text{ m}^2$ . Assume that the pressure is constant throughout the stream of water and that the flow is streamlined. The cross-sectional area of stream, 0.15 m below the tap would be:

(Take  $g = 10 \text{ ms}^{-2}$ )

- (1)  $1 \times 10^{-5} \text{ m}^2$  (2)  $5 \times 10^{-5} \text{ m}^2$   
(3)  $5 \times 10^{-4} \text{ m}^2$  (4)  $2 \times 10^{-5} \text{ m}^2$

Answer (2)

Sol. Using Bernoulli's equation  $v_2 = \sqrt{v_1^2 + 2gh}$

Equation of continuity

$$A_1V_1 = A_2V_2$$

$$(1 \text{ cm}^2)(1 \text{ m/s}) = (A_2) \left( \sqrt{(1)^2 + 2 \times 10 \times \frac{15}{100}} \right)$$

$$\Rightarrow A_2 (1 \text{ cm}^2) = \frac{1}{2}$$

$$\Rightarrow A_2 = 5 \times 10^{-5} \text{ m}^2$$

17. When heat  $Q$  is supplied to a diatomic gas of rigid molecules, at constant volume its temperature increases by  $\Delta T$ . The heat required to produce the same change in temperature, at a constant pressure is:

- (1)  $\frac{3}{2}Q$  (2)  $\frac{2}{3}Q$   
 (3)  $\frac{7}{5}Q$  (4)  $\frac{5}{3}Q$

Answer (3)

Sol. Heat supplied at constant volume

$$Q = nC_v\Delta T$$


and heat supplied at constant pressure

$$Q_1 = nC_p\Delta T$$

$$\therefore \frac{Q_1}{Q} = \frac{C_p}{C_v}$$

$$\Rightarrow Q_1 = (Q)\left(\frac{7}{5}\right)$$

18. A simple pendulum of length  $L$  is placed between the plates of a parallel plate capacitor having electric field  $E$ , as shown in figure. Its bob has mass  $m$  and charge  $q$ . The time period of the pendulum is given by:



- (1)  $2\pi\sqrt{\frac{L}{(g - \frac{qE}{m})}}$  (2)  $2\pi\sqrt{\frac{L}{(g + \frac{qE}{m})}}$   
 (3)  $2\pi\sqrt{\frac{L}{g^2 + \left(\frac{qE}{m}\right)^2}}$  (4)  $2\pi\sqrt{\frac{L}{g^2 - \frac{q^2E^2}{m^2}}}$

Answer (3)

$$\text{Sol. } t = 2\pi\sqrt{\frac{L}{g_{\text{eff}}}}$$

$$\Rightarrow g_{\text{eff}} = \sqrt{g^2 + \left(\frac{qE}{m}\right)^2}$$

$$\Rightarrow t = 2\pi\sqrt{\frac{L}{g^2 + \left(\frac{qE}{m}\right)^2}}$$

19. A source of sound  $S$  is moving with a velocity of 50 m/s towards a stationary observer. The observer measures the frequency of the source as 1000 Hz. What will be the apparent frequency of the source when it is moving away from the observer after crossing him? (Take velocity of sound in air is 350 m/s)

- (1) 857 Hz (2) 1143 Hz  
 (3) 807 Hz (4) 750 Hz

Answer (4)

$$\text{Sol. } f_{\text{app}} = f_{\text{act}} \left( \frac{V \pm V_0}{V \mp V_s} \right)$$

$$1000 = f_{\text{act}} \left( \frac{350 - 0}{350 + (-50)} \right) \text{ and } f' = f_{\text{act}} \left( \frac{350}{350 + 50} \right)$$

$$\Rightarrow f_{\text{act}} = \frac{1000 \times 300}{400}$$

$$f_{\text{act}} \approx 750 \text{ Hz}$$

20. A 2 mW laser operates at a wavelength of 500 nm. The number of photons that will be emitted per second is:

[Given Planck's constant  $h = 6.6 \times 10^{-34}$  Js, speed of light  $c = 3.0 \times 10^8$  m/s]

- (1)  $2 \times 10^{16}$   
 (2)  $1.5 \times 10^{16}$   
 (3)  $1 \times 10^{16}$   
 (4)  $5 \times 10^{15}$

Answer (4)

$$\text{Sol. } E = \frac{hc}{\lambda}$$

Let no. of photons per sec. is  $N$

$$\Rightarrow N \frac{hc}{\lambda} = 2 \text{ mW}$$


$$\Rightarrow N = \frac{2 \times \lambda}{hc} = \frac{2 \times 5000 \times 10^{-3}}{12420 \times 1.6 \times 10^{-19}}$$

$$N = 5 \times 10^{15}$$

21. Space between two concentric conducting spheres of radii  $a$  and  $b$  ( $b > a$ ) is filled with a medium of resistivity  $\rho$ . The resistance between the two spheres will be:

(1)  $\frac{\rho}{4\pi} \left( \frac{1}{a} - \frac{1}{b} \right)$       (2)  $\frac{\rho}{2\pi} \left( \frac{1}{a} - \frac{1}{b} \right)$   
 (3)  $\frac{\rho}{2\pi} \left( \frac{1}{a} + \frac{1}{b} \right)$       (4)  $\frac{\rho}{4\pi} \left( \frac{1}{a} + \frac{1}{b} \right)$

Answer (1)



$$dR = \frac{(\rho)(dx)}{4\pi x^2}$$

$$R = \int dR$$

$$= \left( \frac{\rho}{4\pi} \right) = \int_a^b \frac{dx}{x^2}$$


$$= \left( \frac{\rho}{4\pi} \right) \cdot \left( \frac{1}{a} - \frac{1}{b} \right)$$

22. In an experiment, brass and steel wires of length 1 m each with areas of cross section  $1 \text{ mm}^2$  are used. The wires are connected in series and one end of the combined wire is connected to a rigid support and other end is subjected to elongation. The stress required to produce a net elongation of 0.2 mm is,

[Given, the Young's Modulus for steel and brass are, respectively,  $120 \times 10^9 \text{ N/m}^2$  and  $60 \times 10^9 \text{ N/m}^2$ ]

(1)  $1.8 \times 10^6 \text{ N/m}^2$       (2)  $1.2 \times 10^6 \text{ N/m}^2$   
 (3)  $4.0 \times 10^6 \text{ N/m}^2$       (4)  $0.2 \times 10^6 \text{ N/m}^2$

Answer (Bonus)



Corresponding to the stress ( $\sigma$ )

$$\text{Total elongation } \Delta l_{\text{net}} = \frac{\sigma L_1}{Y_1} + \frac{\sigma L_2}{Y_2}$$

$$\sigma = \Delta l \left( \frac{Y_1 Y_2}{Y_1 + Y_2} \right)$$

$$= 0.2 \times 10^{-3} \times \left( \frac{120 \times 60}{180} \right) \times 10^9$$

$$= 8 \times 10^6 \frac{\text{N}}{\text{m}^2} \quad (\text{Answer is not matching})$$

23. A coil of self inductance 10 mH and resistance  $0.1 \Omega$  is connected through a switch to a battery of internal resistance  $0.9 \Omega$ . After the switch is closed, the time taken for the current to attain 80% of the saturation value is: [take  $\ln 5 = 1.6$ ]

(1) 0.002 s      (2) 0.324 s  
 (3) 0.103 s      (4) 0.016 s

Answer (4)

Sol.  $I = I_{\text{sat}} \left( 1 - e^{-\frac{Rt}{L}} \right)$  Here  $R = R_L + r = 1 \Omega$

$$0.8 I_{\text{sat}} = I_{\text{sat}} \left( 1 - e^{-\frac{t}{.01}} \right)$$

$$\Rightarrow \frac{4}{5} = 1 - e^{-100t}$$

$$\Rightarrow e^{-100t} = \left( \frac{1}{5} \right)$$

$$\Rightarrow 100t = \ln 5$$

$$\Rightarrow t = \frac{1}{100} \ln 5$$

$$= 0.016 \text{ sec}$$


24. The magnitude of the magnetic field at the center of an equilateral triangular loop of side 1 m which is carrying a current of 10 A is:

[Take  $\mu_0 = 4\pi \times 10^{-7} \text{ NA}^{-2}$ ]

(1)  $18 \mu\text{T}$       (2)  $1 \mu\text{T}$   
 (3)  $3 \mu\text{T}$       (4)  $9 \mu\text{T}$

Answer (1)

Sol.



$$d = \left( \frac{1}{3} \right) (a \sin 60^\circ)$$

$$d = \frac{a}{3} \times \frac{\sqrt{3}}{2} = \left( \frac{a}{2\sqrt{3}} \right)$$

$$B_0 = 3 \left[ \frac{\mu_0 I}{4\pi d} (\sin 60 + \sin 60) \right]$$

$$= \frac{3\mu_0 I}{4\pi \left( \frac{a}{2\sqrt{3}} \right)} = (2) \left( \frac{\sqrt{3}}{2} \right) \quad \left| \quad d = \left( \frac{1}{3} \right) (a \sin 60) \right.$$

$$= \frac{9 \left( \frac{\mu_0 I}{\pi a} \right)}{2} \quad \left| \quad d = \frac{a}{3} \times \frac{\sqrt{3}}{2} \right.$$

$$= \frac{9 \times 2 \times 10^{-7} \times 10}{1} \quad \left| \quad = \left( \frac{a}{2\sqrt{3}} \right) \right.$$


$$= 18 \mu T$$

25. A square loop is carrying a steady current  $I$  and the magnitude of its magnetic dipole moment is  $m$ . If this square loop is changed to a circular loop and it carries the same current, the magnitude of the magnetic dipole moment of circular loop will be :

- (1)  $\frac{2m}{\pi}$  (2)  $\frac{4m}{\pi}$   
 (3)  $\frac{m}{\pi}$  (4)  $\frac{3m}{\pi}$

Answer (2)

Sol.



$$2\pi r = 4a \Rightarrow r = \left( \frac{2a}{\pi} \right)$$

$$m = (I) a^2$$

$$m_1 = (I) \pi r^2$$

$$m_1 = (I)(\pi) \left( \frac{4a^2}{\pi^2} \right)$$

$$m_1 = \frac{4Ia^2}{\pi}$$

$$m_1 = \frac{4m}{\pi}$$

26. A submarine experiences a pressure of  $5.05 \times 10^6$  Pa at a depth of  $d_1$  in a sea. When it goes further to a depth of  $d_2$ , it experiences a pressure of  $8.08 \times 10^6$  Pa. Then  $d_2 - d_1$  is approximately (density of water =  $10^3$  kg/m<sup>3</sup> and acceleration due to gravity =  $10$  ms<sup>-2</sup>) :

- (1) 600 m (2) 500 m  
 (3) 300 m (4) 400 m

Answer (3)

Sol.  $\Delta P = P_2 - P_1 = \rho g \Delta H$

$$3.03 \times 10^6 = 10^3 \times 10 \times \Delta H$$

$$\Rightarrow \Delta H \approx 300 \text{ m}$$

27. The elastic limit of brass is 379 MPa. What should be the minimum diameter of a brass rod if it is to support a 400 N load without exceeding its elastic limit?

- (1) 0.90 mm  
 (2) 1.16 mm  
 (3) 1.00 mm  
 (4) 1.36 mm

Answer (2)

Sol. Stress =  $\frac{400}{\pi r^2} \leq 379 \times 10^6 \text{ N/m}^2$

$$\Rightarrow r^2 \geq \frac{400}{379 \times 10^6 \pi}$$

$$2r \geq 1.15 \text{ mm}$$

28. Light is incident normally on a completely absorbing surface with an energy flux of  $25 \text{ Wcm}^{-2}$ . If the surface has an area of  $25 \text{ cm}^2$ , the momentum transferred to the surface in 40 min time duration will be :

- (1)  $3.5 \times 10^{-6} \text{ Ns}$   
 (2)  $6.3 \times 10^{-4} \text{ Ns}$   
 (3)  $5.0 \times 10^{-3} \text{ Ns}$   
 (4)  $1.4 \times 10^{-6} \text{ Ns}$

Answer (3)

Sol.  $P = \frac{\Delta E}{C}$

$$= \frac{(25 \times 25) \times 40 \times 60}{3 \times 10^8} \text{ N-s}$$

$$= 5 \times 10^{-3} \text{ N-s}$$


29. A solid sphere of mass  $M$  and radius  $R$  is divided into two unequal parts. The first part

has a mass of  $\frac{7M}{8}$  and is converted into a uniform disc of radius  $2R$ . The second part is converted into a uniform solid sphere. Let  $I_1$  be the moment of inertia of the disc about its axis and  $I_2$  be the moment of inertia of the new sphere about its axis. The ratio  $I_1/I_2$  is given by :

- (1) 140 (2) 185  
 (3) 285 (4) 65

Answer (1)





$$V = \frac{4}{3}\pi R^3$$

$$\left(\frac{7V}{8}\right)$$

$$\frac{V}{8}$$

$$I_1 = \left(\frac{7M}{8}\right)(2R)^2 \frac{1}{2} = \frac{14}{8}MR^2$$

$$I_2 = \frac{2}{5}\left(\frac{M}{8}\right)r^2$$

$$\Rightarrow I_2 = \frac{2}{5}\left(\frac{M}{8}\right)\left(\frac{R^2}{4}\right) = \frac{MR^2}{80}$$

$$\frac{I_1}{I_2} = \frac{14 \times 80}{(8)} = 140$$

$$\left[ \begin{array}{l} \text{We have} \\ \frac{4}{3}\pi r^3 \rho = \frac{M}{8} \\ \Rightarrow r = \left(\frac{R}{2}\right) \end{array} \right]$$

30. In the formula  $X = 5YZ^2$ ,  $X$  and  $Z$  have dimensions of capacitance and magnetic field, respectively. What are the dimensions of  $Y$  in SI units?

$$(1) [M^{-2}L^{-2}T^6A^3]$$

$$(2) [M^{-1}L^{-2}T^4A^2]$$

$$(3) [M^{-2}L^0T^{-4}A^{-2}]$$

$$(4) [M^{-3}L^{-2}T^8A^4]$$

Answer (4)

$$\text{Sol. } X = 5YZ^2$$

$$Y \propto \frac{X}{Z^2}$$

$$X = C = \frac{Q^2}{E} = \frac{[A^2T^2]}{[ML^2T^{-2}]}$$

$$X = [M^{-1}L^{-2}T^4A^2]$$

$$Z = B = \frac{F}{IL}$$

$$Z = [MT^{-2}A^{-1}]$$

$$Y = \frac{[M^{-1}L^{-2}T^4A^2]}{[MT^{-2}A^{-1}]^2}$$

$$Y = [M^{-3}L^{-2}T^8A^4]$$

## PART-B : CHEMISTRY

1. The correct statement is :

- (1) Zincite is a carbonate ore.
- (2) Zone refining process is used for the refining of titanium.
- (3) Aniline is a froth stabilizer.
- (4) Sodium cyanide cannot be used in the metallurgy of silver.


Answer (3)

Sol. Ti is refined by Van Arkel method. Ag is leached by dilute solution of NaCN. Zincite is ZnO. Aniline is a froth stabilizer.

2. The pH of a 0.02 M  $NH_4Cl$  solution will be [given  $K_b(NH_4OH) = 10^{-5}$  and  $\log 2 = 0.301$ ]

- (1) 2.65
- (2) 5.35
- (3) 4.35
- (4) 4.65

Answer (2)



$$\begin{array}{ccccccc} 0.02 - x & & x & & x & & K_h = \frac{10^{-14}}{10^{-5}} = 10^{-9} \\ \approx 0.02 & & & & & & \end{array}$$

$$K_h = \frac{x^2}{0.02}$$

$$10^{-9} \times 2 \times 10^{-2} = x^2$$

$$x = \sqrt{20} \times 10^{-6}$$

$$pH = -\log(\sqrt{20} \times 10^{-6})$$

$$pH = 5.35$$

3. The INCORRECT statement is :

- (1) The spin-only magnetic moments of  $[Fe(H_2O)_6]^{2+}$  and  $[Cr(H_2O)_6]^{2+}$  are nearly similar.
- (2) The gemstone, ruby, has  $Cr^{3+}$  ions occupying the octahedral sites of beryl.
- (3) The spin-only magnetic moment of  $[Ni(NH_3)_4(H_2O)_2]^{2+}$  is 2.83 BM.
- (4) The color of  $[CoCl(NH_3)_5]^{2+}$  is violet as it absorbs the yellow light.

Answer (2)


Sol. Ruby is aluminium oxide ( $Al_2O_3$ ) containing about 0.5 – 1%  $Cr^{3+}$  ions which are randomly distributed in the position normally occupied by  $Al^{3+}$  ions.

4. Number of stereo centers present in linear and cyclic structures of glucose are respectively :


- (1) 5 & 5  
(2) 4 & 4  
(3) 5 & 4  
(4) 4 & 5

Answer (4)

Sol.



4 stereogenic centres




5 stereogenic centres

5. The difference between  $\Delta H$  and  $\Delta U$  ( $\Delta H - \Delta U$ ), when the combustion of one mole of heptane(l) is carried out at a temperature T, is equal to :

- (1)  $-3RT$   
(2)  $4RT$   
(3)  $3RT$   
(4)  $-4RT$

Answer (4)







$$\Delta H - \Delta U = \Delta n_g RT$$

$$\therefore \Delta n_g = -4$$


$$\therefore \Delta H - \Delta U = -4RT$$

6. Which one of the following graphs between molar conductivity ( $\Lambda_m$ ) versus  $\sqrt{C}$  is correct?



Answer (3)


Sol. KCl is more conducting than NaCl




7. The crystal field stabilization energy (CFSE) of  $[\text{Fe}(\text{H}_2\text{O})_6]\text{Cl}_2$  and  $\text{K}_2[\text{NiCl}_4]$ , respectively, are :

- (1)  $-2.4\Delta_o$  and  $-1.2\Delta_t$   
(2)  $-0.6\Delta_o$  and  $-0.8\Delta_t$   
(3)  $-0.4\Delta_o$  and  $-0.8\Delta_t$   
(4)  $-0.4\Delta_o$  and  $-1.2\Delta_t$

Answer (3)



8. For the reaction,




$$\Delta H = -57.2 \text{ kJ mol}^{-1} \text{ and } K_c = 1.7 \times 10^{16}.$$

Which of the following statement is INCORRECT?

- (1) The equilibrium constant is large suggestive of reaction going to completion and so no catalyst is required.
- (2) The addition of inert gas at constant volume will not affect the equilibrium constant.
- (3) The equilibrium will shift in forward direction as the pressure increases.
- (4) The equilibrium constant decreases as the temperature increases.

Answer (1)





$K_c = 1.7 \times 10^{16}$  i.e. reaction goes to completion. Equilibrium constant has no relation with catalyst. Catalyst only affects the rate with which a reaction proceeds.

For the given reaction, catalyst  $\text{V}_2\text{O}_5$  is used to speed up the reaction (Contact process).

9. A hydrated solid X on heating initially gives a monohydrated compound Y. Y upon heating above 373 K leads to an anhydrous white powder Z. X and Z, respectively are :

- (1) Baking soda and dead burnt plaster.
- (2) Baking soda and soda ash.
- (3) Washing soda and soda ash.
- (4) Washing soda and dead burnt plaster.

Answer (3)



X = Washing soda

Z = Soda ash

10. Which of these factors does not govern the stability of a conformation in acyclic compounds?

- (1) Angle strain
- (2) Steric interactions
- (3) Electrostatic forces of interaction
- (4) Torsional strain

Answer (1)

Sol. Angle strain is not present in acyclic compounds.

11. The highest possible oxidation states of uranium and plutonium, respectively, are :

- (1) 6 and 7
- (2) 7 and 6
- (3) 6 and 4
- (4) 4 and 6

Answer (1)

Sol. Maximum oxidation state shown by

Uranium = + 6

Plutonium = + 7


12. The correct option among the following is :





- (1) Colloidal medicines are more effective because they have small surface area.
- (2) Colloidal particles in lyophobic sols can be precipitated by electrophoresis.
- (3) Brownian motion in colloidal solution is faster if the viscosity of the solution is very high.
- (4) Addition of alum to water makes it unfit for drinking.

Answer (2)


Sol. Electrophoresis is used to coagulate lyophobic colloids.

13. The major product obtained in the given reaction is :



- (1) 
- (2) 
- (3) 
- (4) 

Answer (1)



14. The number of pentagons in  $C_{60}$  and trigons (triangles) in white phosphorus, respectively, are :

- (1) 20 and 3
- (2) 12 and 3
- (3) 12 and 4
- (4) 20 and 4

Answer (3)

Sol. Pentagons in  $C_{60} = 12$

Triangles in  $P_4 = 4$

15. For the reaction of  $H_2$  with  $I_2$ , the rate constant is  $2.5 \times 10^{-4} \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$  at  $327^\circ\text{C}$  and  $1.0 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$  at  $527^\circ\text{C}$ . The activation energy for the reaction, in  $\text{kJ mol}^{-1}$  is :

( $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$ )

- (1) 150
- (2) 59
- (3) 72
- (4) 166

Answer (4)

Sol.  $\log \frac{K_2}{K_1} = \frac{E_a}{2.303R} \left( \frac{1}{T_1} - \frac{1}{T_2} \right)$


$$\log \frac{1}{2.5 \times 10^{-4}} = \frac{E_a}{8.314 \times 2.303} \left( \frac{1}{600} - \frac{1}{800} \right)$$

$$3.6 = \frac{E_a}{8.314 \times 2.303} \times \frac{200}{600 \times 800}$$

$$E_a = 165.4 \text{ kJ/mol}$$


$$\approx 166 \text{ kJ/mol}$$

16. The major product 'Y' in the following reaction is :




Answer (3)

Sol.




17. The major product 'Y' in the following reaction is :



Answer (3)

Sol.



18. 1 g of a non-volatile non-electrolyte solute is dissolved in 100 g of two different solvents A and B whose ebullioscopic constants are in the ratio of 1 : 5. The ratio of the elevation in their

boiling points,  $\frac{\Delta T_b(A)}{\Delta T_b(B)}$ , is :

- (1) 1 : 5
- (2) 10 : 1
- (3) 5 : 1
- (4) 1 : 0.2


Answer (1)

Sol.  $\Delta T_b = k_b \times m$

$$\frac{(k_b)_A}{(k_b)_B} = \frac{1}{5}$$


$$\therefore \frac{(\Delta T_b)_A}{(\Delta T_b)_B} = \frac{(k_b)_A}{(k_b)_B} = \frac{1}{5}$$

19. Compound A ( $C_9H_{10}O$ ) shows positive iodoform test. Oxidation of A with  $KMnO_4/KOH$  gives acid B ( $C_8H_6O_4$ ). Anhydride of B is used for the preparation of phenolphthalein. Compound A is:



Answer (2)

Sol.



20. The ratio of the shortest wavelength of two spectral series of hydrogen spectrum is found to be about 9. The spectral series are :

- (1) Paschen and Pfund
- (2) Brackett and Pfund
- (3) Lyman and Paschen
- (4) Balmer and Brackett

Answer (3)

Sol. Shortest wavelength means  $n_2 = \infty$

$$\text{Lyman series } \bar{\nu}_L = \frac{1}{\lambda_L} = -1312 \times \frac{1}{1^2} \times 1^2$$

$$\text{Paschen series } \bar{\nu}_P = \frac{1}{\lambda_P} = -1312 \times \frac{1}{3^2} \times 1^2$$

$$\frac{\bar{\nu}_L}{\bar{\nu}_P} = \frac{\lambda_P}{\lambda_L} = 9$$

21. The correct match between Item-I and Item-II is:

	Item – I		Item – II
(a)	High density polythene	(I)	Peroxide catalyst
(b)	Polyacrylonitrile	(II)	Condensation at high temperature and pressure
(c)	Novolac	(III)	Ziegler – Natta Catalyst
(d)	Nylon 6	(IV)	Acid or base catalyst

- (1) (a)  $\rightarrow$  (III), (b)  $\rightarrow$  (I), (c)  $\rightarrow$  (IV), (d)  $\rightarrow$  (II)  
 (2) (a)  $\rightarrow$  (III), (b)  $\rightarrow$  (I), (c)  $\rightarrow$  (II), (d)  $\rightarrow$  (IV)  
 (3) (a)  $\rightarrow$  (IV), (b)  $\rightarrow$  (II), (c)  $\rightarrow$  (I), (d)  $\rightarrow$  (III)  
 (4) (a)  $\rightarrow$  (II), (b)  $\rightarrow$  (IV), (c)  $\rightarrow$  (I), (d)  $\rightarrow$  (III)

Answer (1)

- Sol. a. HDPE – Ziegler-Natta Catalyst  
 b. Polyacrylonitrile – Peroxide Catalyst  
 c. Novolac – Catalysed by acid or base  
 d. Nylon-6 – Condensation at High T and P

22. Air pollution that occurs in sunlight is :

- (1) Fog
- (2) Oxidising smog
- (3) Acid rain
- (4) Reducing smog

Answer (2)


Sol. Air pollution caused by sunlight is photochemical smog and it is oxidising.

23. The increasing order of nucleophilicity of the following nucleophiles is :

- (a)  $CH_3CO_2^-$
  - (b)  $H_2O$
  - (c)  $CH_3SO_3^-$
  - (d)  $OH^-$
- (1) (d)  $<$  (a)  $<$  (c)  $<$  (b)  
 (2) (b)  $<$  (c)  $<$  (d)  $<$  (a)  
 (3) (a)  $<$  (d)  $<$  (c)  $<$  (b)  
 (4) (b)  $<$  (c)  $<$  (a)  $<$  (d)

Answer (4)

Sol. Greater the negative charge Present on a nucleophilic centre greater would be its nucleophilicity.



24. The correct order of the first ionization enthalpies is :

- (1)  $Mn < Ti < Zn < Ni$     (2)  $Ti < Mn < Zn < Ni$   
 (3)  $Ti < Mn < Ni < Zn$     (4)  $Zn < Ni < Mn < Ti$

Answer (3)

Sol. Order for I.E. is



25. The noble gas that does NOT occur in the atmosphere is :

- (1) Ne                                      (2) Kr  
 (3) He                                      (4) Ra


Answer (4)

Sol. Radon is not present in atmosphere.

26. Which of the following is NOT a correct method of the preparation of benzylamine from cyanobenzene?

- (1) (i)  $SnCl_2 + HCl(gas)$     (ii)  $NaBH_4$   
 (2)  $H_2/Ni$   
 (3) (i)  $LiAlH_4$                       (ii)  $H_3O^+$   
 (4) (i)  $HCl/H_2O$                       (ii)  $NaBH_4$

Answer (4)



27. The minimum amount of  $O_2(g)$  consumed per gram of reactant is for the reaction:


(Given atomic mass : Fe = 56, O = 16, Mg = 24, P = 31, C = 12, H = 1)

- (1)  $2Mg(s) + O_2(g) \rightarrow 2MgO(s)$   
 (2)  $4Fe(s) + 3O_2(g) \rightarrow 2Fe_2O_3(s)$   
 (3)  $C_3H_8(g) + 5O_2(g) \rightarrow 3CO_2(g) + 4H_2O(l)$   
 (4)  $P_4(s) + 5O_2(g) \rightarrow P_4O_{10}(s)$

Answer (2)

Sol. (1)  $2Mg + O_2 \rightarrow 2MgO$


$$1 \text{ g requires } \frac{32}{48} \text{ g} = 0.66 \text{ g of } O_2$$



1 g Fe requires = 0.43 g of oxygen




1 g of  $C_3H_8$  requires = 3.6 g of  $O_2$



1 g of P requires = 1.3 g of oxygen

28. Points I, II and III in the following plot respectively correspond to

( $V_{mp}$  : most probable velocity)



- (1)  $V_{mp}$  of  $N_2$  (300 K);  $V_{mp}$  of  $O_2$  (400 K);  
 $V_{mp}$  of  $H_2$  (300 K)  
 (2)  $V_{mp}$  of  $H_2$  (300 K);  $V_{mp}$  of  $N_2$  (300 K);  
 $V_{mp}$  of  $O_2$  (400 K)  
 (3)  $V_{mp}$  of  $N_2$  (300 K);  $V_{mp}$  of  $H_2$  (300 K);  
 $V_{mp}$  of  $O_2$  (400 K)  
 (4)  $V_{mp}$  of  $O_2$  (400 K);  $V_{mp}$  of  $N_2$  (300 K);  
 $V_{mp}$  of  $H_2$  (300 K)

Answer (1)

$$\text{Sol. } V_{mp} = \sqrt{\frac{2RT}{M}}$$

$\therefore$  as  $\frac{T}{M}$  increases,  $V_{mp}$  increases

From curve

$$(V_{mp})_I < (V_{mp})_{II} < (V_{mp})_{III}$$

$$(V_{mp})_{N_2} \propto \sqrt{\frac{300}{28}}, (V_{mp})_{O_2} \propto \sqrt{\frac{400}{32}}, (V_{mp})_{H_2} \propto \sqrt{\frac{300}{2}}$$

$$\therefore (V_{mp})_{N_2} < (V_{mp})_{O_2} < (V_{mp})_{H_2}$$

(Under given Condition)

29. The correct statements among (a) to (d) are :
- Saline hydrides produce  $H_2$  gas when reacted with  $H_2O$ .
  - Reaction of  $LiAlH_4$  with  $BF_3$  leads to  $B_2H_6$ .
  - $PH_3$  and  $CH_4$  are electron - rich and electron - precise hydrides, respectively.
  - $HF$  and  $CH_4$  are called as molecular hydrides.
- (a), (c) and (d) only.
  - (c) and (d) only.
  - (a), (b) and (c) only.
  - (a), (b), (c) and (d).

**Answer (4)**

**Sol.** – With water saline hydrides produce  $H_2$  gas

$$3LiAlH_4 + 4BF_3 \longrightarrow 2B_2H_6 + 3LiF + 3AlF_3$$

- $PH_3$  is electron rich while  $CH_4$  is electron precise hydride
  - $HF$  and  $CH_4$  are molecular hydrides
30. In chromatography, which of the following statements is INCORRECT for  $R_f$  ?
- The value of  $R_f$  cannot be more than one.
  - Higher  $R_f$  value means higher adsorption.
  - $R_f$  value is dependent on the mobile phase.
  - $R_f$  value depends on the type of chromatography.

**Answer (2)**

**Sol.**  $R_f$  represents retardation factor in chromatography.

$$R_f = \frac{\text{Distance moved by the substance from base line}}{\text{Distance moved by the solvent from baseline}}$$

- Higher  $R_f$  value means lower adsorption

## PART-C : MATHEMATICS

1. Let  $f(x) = \log_e(\sin x)$ , ( $0 < x < \pi$ ) and  $g(x) = \sin^{-1}(e^{-x})$ , ( $x \geq 0$ ). If  $\alpha$  is a positive real number such that  $a = (f \circ g)'(\alpha)$  and  $b = (f \circ g)(\alpha)$ , then :

- $a\alpha^2 - b\alpha - a = 1$
- $a\alpha^2 + b\alpha + a = 0$
- $a\alpha^2 - b\alpha - a = 0$
- $a\alpha^2 + b\alpha - a = -2\alpha^2$

**Answer (1)**

**Sol.**  $f(x) = \ln(\sin x)$ ,  $g(x) = \sin^{-1}(e^{-x})$

$$f(g(x)) = \ln(\sin(\sin^{-1} e^{-x}))$$

$$= -x$$

$$\Rightarrow -\alpha = b$$

$$f'(g(\alpha)) = a$$

$$\text{i.e., } a = -1$$

$$\therefore a\alpha^2 - b\alpha + 1 = -\alpha^2 + \alpha^2 + 1 = -a$$

2. The angles A, B and C of a triangle ABC are in A.P. and  $a : b = 1 : \sqrt{3}$ . If  $c = 4$  cm, then the area (in sq.cm) of this triangle is :

- $\frac{2}{\sqrt{3}}$
- $4\sqrt{3}$
- $2\sqrt{3}$
- $\frac{4}{\sqrt{3}}$

**Answer (3)**

**Sol.**  $\because$  A, B, C, are in A.P

$$\Rightarrow 2B = A + C$$

$$\Rightarrow B = \frac{\pi}{3}$$

$$\text{Area} = \frac{1}{2}(4x)\sin 60^\circ$$

$$= \sqrt{3}x$$

$$\text{Now } \cos 60^\circ = \frac{16 + x^2 - 3x^2}{8x}$$

$$\Rightarrow 4x = 16 - 2x^2$$

$$x = 2 \text{ (as } -4 \text{ is rejected)}$$


$$\text{Hence, area} = 2\sqrt{3} \text{ sq. cm}$$

3. The sum of the real roots of the equation

$$\begin{vmatrix} x & -6 & -1 \\ 2 & -3x & x-3 \\ -3 & 2x & x+2 \end{vmatrix} = 0, \text{ is equal to :}$$

- 6
- 0
- 4
- 1

**Answer (2)**





Sol.  $\begin{vmatrix} x & -6 & -1 \\ 2 & -3x & x-3 \\ -3 & 2x & x+2 \end{vmatrix} = 0$

$$\Rightarrow x(-3x^2 - 6x - 2x^2 + 6x) - 6(-3x + 9 - 2x - 4) - (4x - 9x) = 0$$

$$\Rightarrow x(-5x^2) - 6(-5x + 5) - 4x + 9x = 0$$

$$\Rightarrow x^3 - 7x + 6 = 0$$

All the roots are real

$$\therefore \text{Sum of real roots} = \frac{0}{1} = 0$$

4. If  $z$  and  $w$  are two complex numbers such that

$$|zw| = 1 \text{ and } \arg(z) - \arg(w) = \frac{\pi}{2}, \text{ then :}$$

$$(1) z\bar{w} = \frac{1-i}{\sqrt{2}} \quad (2) \bar{z}w = i$$

$$(3) z\bar{w} = \frac{-1+i}{\sqrt{2}} \quad (4) \bar{z}w = -i$$

Answer (4)

Sol.  $|zw| = 1 \quad \dots(i)$

$$\arg\left(\frac{z}{w}\right) = \frac{\pi}{2} \quad \dots(ii)$$

$$\therefore \frac{z}{w} + \frac{\bar{z}}{\bar{w}} = 0 \Rightarrow z\bar{w} = -\bar{z}w$$

$$\text{from (i)} \quad z\bar{z}w\bar{w} = 1$$

$$(\bar{z}w)^2 = -1 \Rightarrow \bar{z}w = \pm i$$

$$\text{from (ii)} \quad -\arg(\bar{z}) - \arg w = \frac{\pi}{2}$$

$$\Rightarrow \arg(\bar{z}w) = \frac{-\pi}{2}$$

$$\text{Hence, } \bar{z}w = -i$$

5. The locus of the centres of the circles, which touch the circle,  $x^2 + y^2 = 1$  externally, also touch the  $y$ -axis and lie in the first quadrant, is :


$$(1) y = \sqrt{1+2x}, x \geq 0 \quad (2) x = \sqrt{1+4y}, y \geq 0$$

$$(3) x = \sqrt{1+2y}, y \geq 0 \quad (4) y = \sqrt{1+4x}, x \geq 0$$

Answer (1)

Sol. Let centre of required circle is  $(h, k)$ .

$$\therefore OO' = r + r'$$



$$\Rightarrow \sqrt{h^2 + k^2} = 1 + h$$

$$h^2 + k^2 = 1 + h^2 + 2h$$

$$k^2 = 1 + 2h$$

$$\text{Locus is } y = \sqrt{1+2x}$$

6. The sum  $1 + \frac{1^3+2^3}{1+2} + \frac{1^3+2^3+3^3}{1+2+3} + \dots$   
 $+ \frac{1^3+2^3+3^3+\dots+15^3}{1+2+3+\dots+15} - \frac{1}{2}(1+2+3+\dots+15)$

is equal to :

$$(1) 1860$$

$$(2) 620$$

$$(3) 660$$

$$(4) 1240$$

Answer (2)

Sol.  $S = 1 + \frac{1^3+2^3}{1+2} + \frac{1^3+2^3+3^3}{1+2+3} + \dots 15 \text{ terms}$

$$T_n = \frac{1^3+2^3+\dots+n^3}{1+2+\dots+n} = \frac{\frac{n^2(n+1)^2}{4}}{\frac{n(n+1)}{2}} = \frac{n(n+1)}{2}$$

$$S = \frac{1}{2} \left( \sum_{n=1}^{15} n^2 + \sum_{n=1}^{15} n \right) = \frac{1}{2} \left( \frac{15(16)(31)}{6} + \frac{15(16)}{2} \right)$$

$$= 680$$

$$\Rightarrow 680 - \frac{1}{2} \frac{15(16)}{2} = 680 - 60 = 620$$

7. The smallest natural number  $n$ , such that the coefficient of  $x$  in the expansion of  $\left(x^2 + \frac{1}{x^3}\right)^n$  is  ${}^nC_{23}$ , is :

$$(1) 58$$

$$(2) 35$$

$$(3) 38$$

$$(4) 23$$

Answer (3)



Sol.  $\left(x^2 + \frac{1}{x^3}\right)^n$

General term  $T_{r+1} = {}^nC_r (x^2)^{n-r} \left(\frac{1}{x^3}\right)^r$

${}^nC_r \cdot x^{2n-5r}$

for coefficient of  $x$ ,  $2n - 5r = 1$

Given  ${}^nC_r = {}^nC_{23}$

$r = 23$  or  $n - r = 23$

$\Rightarrow n = 58$  or  $n = 38$

Minimum value is  $n = 38$

8. If both the mean and the standard deviation of 50 observations  $x_1, x_2, \dots, x_{50}$  are equal to 16, then the mean of  $(x_1 - 4)^2, (x_2 - 4)^2, \dots, (x_{50} - 4)^2$  is :

- (1) 380 (2) 480  
(3) 400 (4) 525

Answer (3)

Sol.  $\frac{x_1 + x_2 + \dots + x_{50}}{50} = 16$

$16^2 = \frac{x_1^2 + x_2^2 + \dots + x_{50}^2}{50} - 16^2$

$2(16)^2 \cdot 50 = x_1^2 + x_2^2 + \dots + x_{50}^2$

Required mean =  $\frac{(x_1 - 4)^2 + (x_2 - 4)^2 + \dots + (x_{50} - 4)^2}{50}$   
 $= \frac{16^2(100) + 4^2(50) - 8(16 \times 50)}{50}$   
 $= 16^2(2) + 16 - 8(16) = 400$

9. If  $\int x^5 e^{-x^2} dx = g(x) e^{-x^2} + c$ , where  $c$  is a constant of integration, then  $g(-1)$  is equal to :

- (1) -1 (2)  $-\frac{1}{2}$   
(3) 1 (4)  $-\frac{5}{2}$

Answer (4)

Sol.  $I = \int x^5 \cdot e^{-x^2} dx$

Put  $-x^2 = t \Rightarrow -2x dx = dt$

$I = \int \frac{t^2 \cdot e^t dt}{(-2)} = \frac{-1}{2} e^t (t^2 - 2t + 2) + c$

$\therefore g(x) = \frac{-1}{2} (x^4 + 2x^2 + 2)$

$g(-1) = \frac{-5}{2}$

10. Let  $a, b$  and  $c$  be in G.P. with common ratio  $r$ , where  $a \neq 0$  and  $0 < r \leq \frac{1}{2}$ . If  $3a, 7b$  and  $15c$  are the first three terms of an A.P., then the 4<sup>th</sup> term of this A.P. is :

- (1)  $\frac{2}{3}a$  (2)  $a$   
(3)  $\frac{7}{3}a$  (4)  $5a$

Answer (2)

Sol. Let  $b = ar, c = ar^2$

AP :  $3a, 7ar, 15ar^2$

$14ar = 3a + 15ar^2$

$\Rightarrow 15r^2 - 14r + 3 = 0$

$\Rightarrow r = \frac{1}{3}$  or  $\frac{3}{5}$  (rejected)

Fourth term =  $15ar^2 + 7ar - 3a$

$= a(15r^2 + 7r - 3)$

$= a\left(\frac{15}{9} + \frac{7}{3} - 3\right)$


$= a$

11. The area (in sq. units) of the region bounded by the curves  $y = 2^x$  and  $y = |x + 1|$ , in the first quadrant is :

- (1)  $\frac{3}{2} - \frac{1}{\log_e 2}$  (2)  $\frac{1}{2}$   
(3)  $\log_e 2 + \frac{3}{2}$  (4)  $\frac{3}{2}$

Answer (1)

Sol.



Area =  $\int_0^1 ((x+1) - 2^x) dx$

$= \left[ \frac{x^2}{2} + x - \frac{2^x}{\ln 2} \right]_0^1$


$= \left( \frac{1}{2} + 1 - \frac{2}{\ln 2} \right) - \left( \frac{-1}{\ln 2} \right)$

$= \frac{3}{2} - \frac{1}{\ln 2}$

12. A perpendicular is drawn from a point on the line  $\frac{x-1}{2} = \frac{y+1}{-1} = \frac{z}{1}$  to the plane  $x + y + z = 3$  such that the foot of the perpendicular Q also lies on the plane  $x - y + z = 3$ . Then the co-ordinates of Q are :

- (1) (1, 0, 2)                      (2) (2, 0, 1)  
(3) (4, 0, -1)                    (4) (-1, 0, 4)

Answer (2)



Let Q be  $(\alpha, \beta, \gamma)$

$$\alpha + \beta + \gamma = 3 \quad \dots(i)$$

$$\alpha - \beta + \gamma = 3 \quad \dots(ii)$$

$$\therefore \alpha + \gamma = 3 \text{ and } \beta = 0$$

Equating DR's of PQ :

$$\frac{\alpha - 2\lambda - 1}{1} = \frac{\lambda + 1}{1} = \frac{\gamma - \lambda}{1}$$

$$\Rightarrow \alpha = 3\lambda + 2, \gamma = 2\lambda + 1$$

Substituting in equation (i) we get

$$\Rightarrow 5\lambda + 3 = 3$$

$$\lambda = 0$$

Point is Q(2, 0, 1)

13. If the tangent to the curve  $y = \frac{x}{x^2 - 3}$ ,  $x \in \mathbb{R}$ , ( $x \neq \pm\sqrt{3}$ ), at a point  $(\alpha, \beta) \neq (0, 0)$  on it is parallel to the line  $2x + 6y - 11 = 0$ , then :

- (1)  $|6\alpha + 2\beta| = 19$                       (2)  $|2\alpha + 6\beta| = 19$   
(3)  $|6\alpha + 2\beta| = 9$                         (4)  $|2\alpha + 6\beta| = 11$

Answer (1)

Sol.  $y = \frac{x}{x^2 - 3}$

$$\frac{dy}{dx} = \frac{(x^2 - 3) - x(2x)}{(x^2 - 3)^2} = \frac{-x^2 - 3}{(x^2 - 3)^2}$$

$$\left. \frac{dy}{dx} \right|_{(\alpha, \beta)} = \frac{-\alpha^2 - 3}{(\alpha^2 - 3)^2} = -\frac{1}{3}$$

$$3(\alpha^2 + 3) = (\alpha^2 - 3)^2 \quad \dots(i)$$

i.e.  $\alpha^2 = 9$

$$\text{Also, } \beta = \frac{\alpha}{\alpha^2 - 3} \Rightarrow \alpha^2 - 3 = \frac{\alpha}{\beta} \Rightarrow \frac{\alpha}{\beta} = 6$$

$$\Rightarrow \alpha = \pm 3, \beta = \pm \frac{1}{2}$$

Which satisfies  $|6\alpha + 2\beta| = 19$

14. Let  $y = y(x)$  be the solution of the differential equation,  $\frac{dy}{dx} + y \tan x = 2x + x^2 \tan x$ ,  $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ , such that  $y(0) = 1$ . Then :

$$(1) y\left(\frac{\pi}{4}\right) + y\left(-\frac{\pi}{4}\right) = \frac{\pi^2}{2} + 2$$

$$(2) y\left(\frac{\pi}{4}\right) - y\left(-\frac{\pi}{4}\right) = \sqrt{2}$$

$$(3) y'\left(\frac{\pi}{4}\right) + y'\left(-\frac{\pi}{4}\right) = -\sqrt{2}$$

$$(4) y'\left(\frac{\pi}{4}\right) - y'\left(-\frac{\pi}{4}\right) = \pi - \sqrt{2}$$

Answer (4)

Sol.  $\frac{dy}{dx} + y \tan x = 2x + x^2 \tan x$

$$P = \tan x, Q = 2x + x^2 \tan x$$

$$\text{I.F.} = e^{\int \tan x dx} = e^{\ln|\sec x|} = |\sec x|$$

$$\begin{aligned} y(\sec x) &= \int (2x + x^2 \tan x) \sec x dx \\ &= \int x^2 \tan x \sec x dx + \int 2x \sec x dx \\ &= x^2 \sec x - \int 2x \sec x dx + \int 2x \sec x dx \\ &= x^2 \sec x + c \end{aligned}$$

$$\text{As } y(0) = 1, c = 1$$

$$\therefore y = x^2 + \cos x$$

$$\text{At } x = \frac{\pi}{4}, y\left(\frac{\pi}{4}\right) = \frac{\pi^2}{16} + \frac{1}{\sqrt{2}}$$

$$y\left(-\frac{\pi}{4}\right) = \frac{\pi^2}{16} + \frac{1}{\sqrt{2}}$$

$$y\left(\frac{\pi}{4}\right) - y\left(-\frac{\pi}{4}\right) = 0$$

$$\frac{dy}{dx} = 2x - \sin x$$

$$y'\left(\frac{\pi}{4}\right) = \frac{\pi}{2} - \frac{1}{\sqrt{2}}, y'\left(-\frac{\pi}{4}\right) = -\frac{\pi}{2} + \frac{1}{\sqrt{2}}$$

$$y'\left(\frac{\pi}{4}\right) - y'\left(-\frac{\pi}{4}\right) = \pi - \sqrt{2}$$

15. Let  $a_1, a_2, a_3, \dots$  be an A.P. with  $a_6 = 2$ . Then the common difference of this A.P., which maximises the product  $a_1 a_4 a_5$ , is :

- (1)  $\frac{2}{3}$  (2)  $\frac{8}{5}$   
(3)  $\frac{3}{2}$  (4)  $\frac{6}{5}$

**Answer (1)**

**Sol.**  $a + 5d = 2$

$$\begin{aligned} \text{Let } A &= a_1 a_4 a_5 = a(a + 3d)(a + 4d) \\ &= a(2 - 2d)(2 - d) \end{aligned}$$

$$A = (2 - 5d)(4 - 6d + 2d^2)$$

$$\frac{dA}{dd} = 0$$

$$(2 - 5d)(-6 + 4d) + (4 - 6d + 2d^2)(-5) = 0$$

$$\Rightarrow 15d^2 - 34d + 16 = 0$$

$$d = \frac{8}{5}, \frac{2}{3}$$

$$\text{For } d = \frac{2}{3}, \frac{d^2 A}{dd^2} < 0$$

$$\text{Hence } d = \frac{2}{3}$$

16. If the plane  $2x - y + 2z + 3 = 0$  has the distances  $\frac{1}{3}$  and  $\frac{2}{3}$  units from the planes  $4x - 2y + 4z + \lambda = 0$  and  $2x - y + 2z + \mu = 0$ , respectively, then the maximum value of  $\lambda + \mu$  is equal to :

- (1) 13 (2) 15  
(3) 5 (4) 9

**Answer (1)**

**Sol.**  $P_1 : 2x - y + 2z + 3 = 0$

$$P_2 : 2x - y + 2z + \frac{\lambda}{2} = 0$$

$$P_3 : 2x - y + 2z + \mu = 0$$

$$\text{Given } \frac{1}{3} = \frac{\left| 3 - \frac{\lambda}{2} \right|}{\sqrt{9}} \Rightarrow \left| 3 - \frac{\lambda}{2} \right| = 1$$

$$\lambda_{\max} = 8$$

$$\text{Also, } \frac{2}{3} = \frac{|\mu - 3|}{\sqrt{9}} \Rightarrow \mu_{\max} = 5$$

$$(\lambda + \mu)_{\max} = 13$$

17. If  $\lim_{x \rightarrow 1} \frac{x^2 - ax + b}{x - 1} = 5$ , then  $a + b$  is equal to:

- (1) 5 (2) -4  
(3) 1 (4) -7

**Answer (4)**

$$\text{Sol. } \lim_{x \rightarrow 1} \frac{x^2 - ax + b}{x - 1} = 5$$

As limit is finite,  $1 - a + b = 0$

$$\Rightarrow \lim_{x \rightarrow 1} \frac{2x - a}{1} = 5 \quad \left( \frac{0}{0} \text{ form} \right)$$

$$\text{i.e., } 2 - a = 5$$

$$\text{or } a = -3$$

$$\therefore b = -4$$

$$a + b = -3 - 4 = -7$$

18. The number of real roots of the equation  $5 + |2^x - 1| = 2^x(2^x - 2)$  is :

- (1) 4 (2) 2  
(3) 1 (4) 3

**Answer (3)**

**Sol.** Let  $2^x - 1 = t$

$$5 + |t| = (t + 1)(t - 1)$$

$$\Rightarrow |t| = t^2 - 6$$

$$\text{For } t > 0, t^2 - t - 6 = 0$$

$$\text{i.e., } t = 3 \text{ or } -2 \text{ (rejected)}$$

$$\text{For } t < 0, t^2 + t - 6 = 0$$

$$\text{i.e., } t = -3 \text{ or } 2 \text{ (both rejected)}$$

$$\therefore 2^x - 1 = 3$$

$$\Rightarrow x = 2$$

19. Lines are drawn parallel to the line  $4x - 3y + 2 = 0$ , at a distance  $\frac{3}{5}$  from the origin. Then which one of the following points lies on any of these lines?

- (1)  $\left( \frac{1}{4}, -\frac{1}{3} \right)$  (2)  $\left( \frac{1}{4}, \frac{1}{3} \right)$   
(3)  $\left( -\frac{1}{4}, \frac{2}{3} \right)$  (4)  $\left( -\frac{1}{4}, -\frac{2}{3} \right)$

**Answer (3)**

**Sol.** Let straight line be  $4x - 3y + \alpha = 0$

$$\text{Given } \frac{3}{5} = \left| \frac{\alpha}{5} \right|$$

$$\Rightarrow \alpha = \pm 3$$

Line is  $4x - 3y + 3 = 0$  or  $4x - 3y - 3 = 0$

Clearly  $\left(-\frac{1}{4}, \frac{2}{3}\right)$  satisfies  $4x - 3y + 3 = 0$

20. Suppose that 20 pillars of the same height have been erected along the boundary of a circular stadium. If the top of each pillar has been connected by beams with the top of all its non-adjacent pillars, then the total number of beams is :

- (1) 210 (2) 180  
(3) 170 (4) 190

**Answer (3)**

**Sol.** Required number of beams  $= {}^{20}C_2 - 20$   
 $= 190 - 20 = 170$

21. A spherical iron ball of radius 10 cm is coated with a layer of ice of uniform thickness that melts at a rate of  $50 \text{ cm}^3/\text{min}$ . When the thickness of the ice is 5 cm, then the rate at which the thickness (in cm/min) of the ice decreases, is :

- (1)  $\frac{5}{6\pi}$  (2)  $\frac{1}{36\pi}$   
(3)  $\frac{1}{9\pi}$  (4)  $\frac{1}{18\pi}$

**Answer (4)**

**Sol.**  $\frac{dV_{\text{ice}}}{dt} = 50$

$$V_{\text{ice}} = \frac{4}{3}\pi(10+r)^3 - \frac{4}{3}\pi(10)^3$$


$$\frac{dV}{dt} = \frac{4}{3}\pi 3(10+r)^2 \frac{dr}{dt}$$

$$= 4\pi(10+r)^2 \frac{dr}{dt}$$

$$\text{At } r = 5, 50 = 4\pi(225) \frac{dr}{dt}$$

$$\frac{dr}{dt} = \frac{50}{4\pi(225)}$$

$$= \frac{1}{18\pi} \text{ cm/min}$$



22. Minimum number of times a fair coin must be tossed so that the probability of getting at least one head is more than 99% is :

- (1) 8 (2) 6  
(3) 5 (4) 7

**Answer (4)**

**Sol.**  $1 - \left(\frac{1}{2}\right)^n > \frac{99}{100}$

$$\left(\frac{1}{2}\right)^n < \frac{1}{100}$$

$$\therefore n \geq 7$$

Minimum value is 7.

23. If  $5x + 9 = 0$  is the directrix of the hyperbola  $16x^2 - 9y^2 = 144$ , then its corresponding focus is :


- (1)  $\left(\frac{5}{3}, 0\right)$  (2)  $\left(-\frac{5}{3}, 0\right)$   
(3)  $(-5, 0)$  (4)  $(5, 0)$

**Answer (3)**

**Sol.**  $16x^2 - 9y^2 = 144$

$$\text{i.e. } \frac{x^2}{9} - \frac{y^2}{16} = 1$$

Focus  $S'(-ae, 0)$



$$a = 3, b = 4$$

$$e^2 = 1 + \frac{16}{9} = \frac{25}{9}$$

$$S' \equiv \left(-3 \times \frac{5}{3}, 0\right) \equiv (-5, 0)$$

24. Let  $\lambda$  be a real number for which the system of linear equations

$$x + y + z = 6$$

$$4x + \lambda y - \lambda z = \lambda - 2$$

$$3x + 2y - 4z = -5$$

has infinitely many solutions. Then  $\lambda$  is a root of the quadratic equation :

- (1)  $\lambda^2 + 3\lambda - 4 = 0$  (2)  $\lambda^2 - \lambda - 6 = 0$   
(3)  $\lambda^2 + \lambda - 6 = 0$  (4)  $\lambda^2 - 3\lambda - 4 = 0$

**Answer (2)**

Sol.  $\Delta = \Delta_1 = \Delta_2 = \Delta_3 = 0$

$$\Delta = \begin{vmatrix} 1 & 1 & 1 \\ 4 & \lambda & -\lambda \\ 3 & 2 & -4 \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} 0 & 0 & 1 \\ 4-\lambda & 2\lambda & -\lambda \\ 1 & 6 & -4 \end{vmatrix} = 0 \Rightarrow \lambda = 3$$

$$\Delta_1 = \begin{vmatrix} 6 & 1 & 1 \\ \lambda-2 & \lambda & -\lambda \\ -5 & 2 & -4 \end{vmatrix} = 0 \quad \text{for } \lambda = 3$$

$$\Delta_2 = \begin{vmatrix} 1 & 6 & 1 \\ 4 & \lambda-2 & -\lambda \\ 3 & -5 & -4 \end{vmatrix} = 0 \quad \text{for } \lambda = 3$$

$$\Delta_3 = \begin{vmatrix} 1 & 1 & 6 \\ 4 & \lambda & \lambda-2 \\ 3 & 2 & -5 \end{vmatrix} = 0 \quad \text{for } \lambda = 3$$

$\therefore$  For  $\lambda = 3$ , infinitely many solutions is obtained.

25. The integral  $\int_{\pi/6}^{\pi/3} \sec^{2/3} x \operatorname{cosec}^{4/3} x dx$  is equal to :

- (1)  $3^{7/6} - 3^{5/6}$  (2)  $3^{5/3} - 3^{1/3}$   
(3)  $3^{5/6} - 3^{2/3}$  (4)  $3^{4/3} - 3^{1/3}$

Answer (1)

Sol.  $I = \int_{\pi/6}^{\pi/3} \sec^{2/3} x \cdot \operatorname{cosec}^{4/3} x dx$

$$= \int_{\pi/6}^{\pi/3} \frac{1 \cdot dx}{\cos^{2/3} x \cdot \sin^{4/3} x}$$

$$= \int_{\pi/6}^{\pi/3} \frac{1}{\cos^2 x \cdot \tan^{4/3} x} dx = \int_{\pi/6}^{\pi/3} \frac{\sec^2 x dx}{\tan^{4/3} x}$$

Let  $\tan x = t$

$$I = \int_{\frac{1}{\sqrt{3}}}^{\sqrt{3}} t^{-4/3} dt = \frac{3 \left[ t^{-1/3} \right]_{\frac{1}{\sqrt{3}}}^{\sqrt{3}}}{-1}$$

$$= -3 \left[ 3^{-1/6} - \frac{1}{3^{-1/6}} \right]$$

$$= -3(3^{-1/6} - 3^{1/6})$$

$$= 3(3^{1/6} - 3^{-1/6})$$

$$= 3^{7/6} - 3^{5/6}$$

26. If the line  $ax + y = c$ , touches both the curves  $x^2 + y^2 = 1$  and  $y^2 = 4\sqrt{2}x$ , then  $|c|$  is equal to :

- (1)  $\frac{1}{\sqrt{2}}$   
(2)  $\frac{1}{2}$   
(3) 2  
(4)  $\sqrt{2}$

Answer (4)

Sol. Tangent on  $y^2 = 4\sqrt{2}x$  is  $yt = x + \sqrt{2}t^2$

As it is tangent on circle also,

$$\left| \frac{\sqrt{2}t^2}{\sqrt{1+t^2}} \right| = 1$$

$$2t^4 = 1 + t^2 \quad \text{i.e. } t^2 = 1$$

$$\text{Equation is } \pm y = x + \sqrt{2}$$

$$\text{Hence } |c| = \sqrt{2}$$


27. The tangent and normal to the ellipse  $3x^2 + 5y^2 = 32$  at the point  $P(2, 2)$  meet the x-axis at  $Q$  and  $R$ , respectively. Then the area (in sq. units) of the triangle  $PQR$  is :

- (1)  $\frac{16}{3}$   
(2)  $\frac{14}{3}$   
(3)  $\frac{34}{15}$   
(4)  $\frac{68}{15}$

Answer (4)

Sol. For  $\frac{3x^2}{32} + \frac{5y^2}{32} = 1$

Tangent at  $P$  is



$$\frac{3(2)x}{32} + \frac{5(2)y}{32} = 1$$

$$\frac{3x}{16} + \frac{5y}{16} = 1$$

$$Q \equiv \left( \frac{16}{3}, 0 \right)$$

$$\text{Normal at P is } \frac{32x}{3(2)} - \frac{32y}{5(2)} = \frac{32}{3} - \frac{32}{5}$$

$$R \equiv \left( \frac{4}{5}, 0 \right)$$


$$\begin{aligned} \text{area of } \triangle PQR &= \frac{1}{2} (PQ) (PR) = \frac{1}{2} \sqrt{\frac{136}{3}} \cdot \sqrt{\frac{136}{5}} \\ &= \frac{68}{15} \end{aligned}$$

28. The distance of the point having position vector  $-\hat{i} + 2\hat{j} + 6\hat{k}$  from the straight line passing through the point  $(2, 3, -4)$  and parallel to the vector,  $6\hat{i} + 3\hat{j} - 4\hat{k}$  is :

- (1) 7 (2)  $4\sqrt{3}$   
(3) 6 (4)  $2\sqrt{13}$

Answer (1)

$$\text{Sol. Equation of l is } \frac{x-2}{6} = \frac{y-3}{3} = \frac{z+4}{-4}$$



$$\text{Let } M(6\lambda + 2, 3\lambda + 3, -4\lambda - 4)$$

$$\text{DR's of PM is } \langle 6\lambda + 3, 3\lambda + 1, -4\lambda - 10 \rangle$$

$$\Rightarrow (6\lambda + 3)(6) + (3\lambda + 1)(3) + (-4\lambda - 10)(-4) = 0$$

$$\Rightarrow \lambda = -1$$

$$\text{i.e. } M \equiv (-4, 0, 0)$$

$$\therefore PM = \sqrt{9+4+36} = 7$$

29. If  $\cos^{-1}x - \cos^{-1}\frac{y}{2} = \alpha$ , where  $-1 \leq x \leq 1$ ,  $-2 \leq y \leq 2$ ,

$x \leq \frac{y}{2}$ , then for all  $x, y$ ,  $4x^2 - 4xy \cos \alpha + y^2$  is equal to :

- (1)  $2 \sin^2 \alpha$   
(2)  $4 \sin^2 \alpha - 2x^2y^2$   
(3)  $4 \cos^2 \alpha + 2x^2y^2$   
(4)  $4 \sin^2 \alpha$

Answer (4)

$$\text{Sol. } \cos^{-1}x - \cos^{-1}\frac{y}{2} = \alpha$$

$$\Rightarrow \cos^{-1}\left(\frac{xy}{2} + \sqrt{1-x^2} \cdot \sqrt{1-\frac{y^2}{4}}\right) = \alpha$$

$$\Rightarrow \frac{xy}{2} + \frac{\sqrt{1-x^2} \sqrt{4-y^2}}{2} = \cos \alpha$$

$$\Rightarrow xy + \sqrt{1-x^2} \sqrt{4-y^2} = 2 \cos \alpha$$

$$(xy - 2 \cos \alpha)^2 = (1-x^2)(4-y^2)$$

$$x^2y^2 + 4\cos^2 \alpha - 4xycos \alpha = 4 - y^2 - 4x^2 + x^2y^2$$

$$4x^2 - 4xycos \alpha + y^2 = 4\sin^2 \alpha$$

30. The negation of the Boolean expression  $\sim s \vee (\sim r \wedge s)$  is equivalent to :

- (1)  $s \wedge r$   
(2)  $r$   
(3)  $\sim s \wedge \sim r$   
(4)  $s \vee r$

Answer (1)

$$\text{Sol. } \sim s \vee (\sim r \wedge s)$$

$$\equiv (\sim s \vee \sim r) \wedge (\sim s \vee s)$$

$$\equiv (\sim s \vee \sim r) \quad (\because (\sim s \vee s) \text{ is tautology})$$

$$\equiv \sim(s \wedge r)$$

Hence its negation is  $s \wedge r$

