TASK 5 | INDIVIDUAL CASE SAFETY REPORT (ICSR) PROCESSING

Case-based exercise • Drug Safety Associate simulation • End-to-end ICSR workflow | FICTIONAL CASE

CASE VALIDITY & TRIAGE

Minimum criteria
• Identifiable patient ✓
• Identifiable reporter ✓
• Suspect product ✓
• Adverse event ✓

FINAL: VALID ICSR
All four minimum criteria documented.

PATIENT & REPORTER INFORMATION

Patient: RS-02
Age / sex: 35 years / Male
Weight: 74 kg
Reporter: Dr. P. Das, physician
Report type: Spontaneous
Contact: Not reported

PRODUCT EXPOSURE

Suspect: Cefixime
Dose: 200 mg
Route: Oral
Frequency: Twice daily
Indication: Acute bacterial pharyngitis

EVENT EVALUATION

Event: Generalized urticaria and facial swelling
Onset: About 3 hours after first dose
Severity: Severe
Clinical impact: Emergency assessment and treatment required
Outcome: Recovered after treatment and withdrawal

Case feature: Rapid onset; no documented airway compromise.

SERIOUSNESS ASSESSMENT

Life-threatening: Not documented
Hospitalization: Yes — admitted for observation
Persistent disability: No
Other medically important event: Yes

OVERALL: SERIOUS
Hospitalization and medical importance are the stated criteria.

DECHALLENGE • RECHALLENGE • FOLLOW-UP

Dechallenge: Positive — symptoms improved after stopping cefixime
Rechallenge: Not performed
Investigation: Clinical examination; no specific laboratory test
Alternative causes: No clear alternative documented
Follow-up: Confirm onset/treatment times, admission/discharge dates, airway symptoms, allergy history and final outcome.

CAUSALITY ASSESSMENT

Reporter assessment: Possible / probable pending full review
Temporal relationship: Reasonable
Dechallenge: Positive
Rechallenge: Not performed
Alternative causes: None clearly documented
WHO–UMC: Confirm category after full review; limited details do not justify overstatement.

EXPECTEDNESS & MEDDRA CODING

Expectedness: Pending verification against current product information for the exact cefixime product and reaction term.
Proposed event term: Urticaria — verify the appropriate MedDRA Preferred Term and hierarchy in the current browser.
Important: Do not assign a MedDRA code without hierarchy verification.

PROFESSIONAL CASE NARRATIVE & DISPOSITION

A 35-year-old male receiving oral cefixime 200 mg twice daily for acute bacterial pharyngitis developed generalized urticaria and facial swelling approximately three hours after the first dose. Cefixime was stopped, emergency assessment and treatment were provided, and symptoms improved. The patient was admitted for observation and recovered. Rechallenge was not performed, and no clear alternative cause was documented. The report includes an identifiable patient, reporter, suspected product and adverse event, meeting the four minimum ICSR validity criteria. The event is serious due to hospitalization and medical importance. Follow-up and verification of expectedness and MedDRA coding are required.

Fictional training case only — not a real patient report. Final validity, seriousness, causality, expectedness and MedDRA coding should be reviewed using complete source information and applicable guidance.