

Nine-Species MEA ePC-SAFT Parameter Bundle

Status: active MEA parameter bundle. This repository owns the live MEA bundle recorded here. A bounded, direct Engine campaign retains Uyan solvent-only dielectric mixing and selects the displayed R4 carbamate-hydrolysis correlation and ϵ_{CO_2}/k . Relative to the previous bundle, the selected result improves full-data pressure $\log_{10}$ RMSE from 0.3974 to 0.3540, temperature-balanced pressure RMSE from 0.3569 to 0.3185, and pressure coverage from 125/161 to 130/161 while retaining all 131/131 observed speciation targets. Value color alone identifies one of the four evidence classes; parameter values use regular weight so typography does not imply a second status system. This PDF is the live present-state view: new accepted evidence replaces the corresponding prior interpretation in place, while Git preserves history.

1. Question and scope

What full parameter bundle is being used for the current calculations? This notebook records the nine-species model structure, pure-component values, association graph, sparse binary interactions, electrostatic choices, reaction correlations, and known non-adopted diagnostics in one reviewable view. Section 9 replays the exact active vector after the audited R1–R3 standard-state correction using its pinned Engine wheel; Section 8 identifies the retained pre-campaign smooth-grid visual. The reported results use the parameter set shown in Tables 1–4.

The liquid species are CO_2 , MEA, H_2O , MEA^+H , MEACOO^- , HCO_3^- , CO_3^{2-} , H_3O^+ , and OH^- . The neutral vapor species and standard-state details remain governed by the future immutable packet.

1.1. How to read this live notebook

verified statements are checked against the retained input or result files, the current repository, an official software source, or a located primary source. **inference** statements interpret those checked facts. **assumption** identifies a proposed test condition, and **unknown** marks a question that the retained evidence does not answer. Unrun experiments are listed only in Section 12; they are not results.

Evidence	Present state	Meaning
verified	Parameter set	Tables 1–4 contain the active nine-species vector retained here.
verified	Current replay	All 131 observed speciation targets and 130/161 canonical pressure states evaluate with the selected bundle; current full-data metrics are reported in Section 9.
verified	Upstream work	ePC-SAFT Issues #80, #119, #126, and #149 remain open. The native Regression and repository-owner cutovers have merged through upstream PR #169, so Issue #119’s old pre-build blocker text is stale even though the scientific campaign has not run.

2. Model conditions

Choice	Current foundation
Residual Helmholtz terms	PC-SAFT hard-chain and dispersion; general-site association; Debye–Hückel and Born electrostatics.
Association topology	Reciprocal induced CO ₂ -H ₂ O cross association plus ordinary MEA and water association. No CO ₂ self-association and no separate CO ₂ -MEA chemical-association edge.
Permittivity and Born formulation	Uyan solvent-only mixing: normalized MEA–water mass fractions determine bulk ϵ_r ; dissolved CO ₂ and ions do not enter the mixing rule. Original Born remains active with $c_{\text{shell}} = 0$ and $c_{\text{dielectric}} = 0$.
Parameter identity	Active Uyan solvent-only bundle. Parameter SHA-256: 2666914f0f9cfefbdf230e96565de843f9aad9424035c940883147ff66af035c.
Refinement boundary	The campaign changes only R4 and ϵ_{CO_2}/k from the prior bundle. R1–R3, R5, the ion block, association topology, and all binary interactions remain fixed.

3. Complete component parameter bundle

Here m is segment number, σ is segment diameter, ϵ/k is dispersion energy, ϵ^{AB}/k is self-association energy, and κ^{AB} is the stored site-volume coordinate. For CO₂, zero self-association energy disables self-association and the listed site volume is used only by the induced cross-association set. Here z is charge and d_B is Born diameter, ϵ_r is relative permittivity, and f_{solv} is the solvation factor. Lengths are in Å and energies in K. References identify the literature lineage; the underlying exact vector is `results/selected-current-best-parameters.json`. The Debye–Hückel diameter is derived as $d_{\text{DH},i} = 0.88\sigma_i$ and is therefore not tabulated as an independent parameter. The hash in Section 2 identifies the unrounded authority behind the four-decimal-place display.

Table 1. Neutral-component parameters. Only parameters applicable to the neutral molecular rows are shown.

Species	Scheme	m_i	σ_i	ϵ_i/k	ϵ_i^{AB}/k	κ_i^{AB}	$\epsilon_{r,i}$	Ref.
CO ₂	2B ind.	2.0729	2.7852	173.4403	0.0000	0.0451	— ^a	[6, 12, 14]
MEA	2B	3.0353	3.0435	277.1740	2586.3000	0.0375	32.0000	[1–3]
H ₂ O	2B	1.2047	$\sigma(T)$	353.9500	2425.7000	0.0451	$\epsilon_r(T)$	[5, 8, 13, 14]

^a Go to Section 4.2 for the source, exact derivation, temperature dependence, and replacement decision.**Table 2. Ionic-species parameters.** Only parameters applicable to the ionic rows are shown.

Species	m_i	σ_i	ϵ_i/k	z_i	Ref.
MEAH ⁺	1.0000	3.4851	232.6872	+1	[5, 15, 16]
MEACOO [−]	1.0000	3.5354	453.2652	−1	[15, 16]
HCO ₃ [−]	1.0000	2.9296	70.0000	−1	[9, 14]
CO ₃ ^{2−}	1.0000	2.4422	249.2600	−2	[9, 14]
H ₃ O ⁺	1.0000	3.4654	500.0000	+1	[5, 9]
OH [−]	1.0000	2.0177	650.0000	−1	[5, 9]

Table 3. Active Born and solvation parameters. Ion-specific dielectric-suppression coordinates are not tabulated because the selected solvent-only formulation does not use them.

Species	$d_{B,i}$	$f_{solv,i}$	Ref.
CO ₂	—	1.0000	[5]
MEA	—	1.0000	[5]
H ₂ O	—	1.5000	[5]
MEAH ⁺	3.5332	1.0000	[5, 15, 16]
MEACOO [−]	3.5411	1.0000	[5, 15, 16]
HCO ₃ [−]	3.0000	1.0000	[5, 9]
CO ₃ ^{2−}	3.0000	1.0000	[5, 9]
H ₃ O ⁺	1.2180	1.0000	[5, 9]
OH [−]	3.0811	1.0000	[5, 9]

Table 4. Binary interaction matrix, k_{ij} . Upper triangle shown; the diagonal is structural and unclassified.

Species	CO ₂	MEA	H ₂ O	MEAH ⁺	MEACOO [−]	HCO ₃ [−]	CO ₃ ^{2−}	H ₃ O ⁺	OH [−]
CO ₂	—	0.0000	0.0133	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
MEA		—	−0.0735	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
H ₂ O			—	0.0000	0.0000	0.0000	−0.2500	0.2500	−0.2500
MEAH ⁺				—	−0.0020	0.0000	0.0000	0.0000	0.0000
MEACOO [−]					—	0.0000	0.0000	0.0000	0.0000
HCO ₃ [−]						—	0.0000	0.0000	0.0000
CO ₃ ^{2−}							—	0.0000	0.0000
H ₃ O ⁺								—	0.0000
OH [−]									—

Class 1: confirmed current foundation**Class 3:** local fit or derivation awaiting promotion**Class 2:** literature-backed or evidence-informed candidate**Class 4:** missing value, default, or indirect analog

4. Evidence-class justification

The class is assigned to each independent displayed value, not to an entire species. Class 1 is fixed for the current foundation and reopens only when contradictory evidence is admitted. Class 2 is a direct literature value or an explicit, evidence-informed working assumption worth testing for this MEA application. Class 3 is a local fit or derivation with numerical evidence that has not passed promotion checks. Class 4 is a missing independent value represented only by a default or indirect analog. This fourth class was previously collapsed into Class 3 only to follow the earlier request for three classifications; no scientific argument required that collapse. Separating local numerical evidence from truly missing values is more useful for the present campaign.

4.1. Class 1: confirmed current foundation

CO₂. The 2B induced scheme, $m = 2.0729$, and $\sigma = 2.7852 \text{ \AA}$ are the Gross–Sadowski molecular row carried through the induced-association literature. Zero self-association energy and $\kappa^{AB} = 0.0451$ implement the Pabsch–Schick induced topology rather than a fitted CO₂ self-association term [6, 12, 14]. The dispersion energy is no longer Class 1; its promoted local value is documented under Class 3.

MEA. The displayed m , σ , ϵ/k , ϵ^{AB}/k , and κ^{AB} are the Baygi 2B row [3]. The 2B topology is the current selection recorded by upstream PR 132; the older 3B diagnostic is not the retained foundation.

H₂O. The Held 2B family fixes m , $\sigma(T)$, ϵ/k , ϵ^{AB}/k , and κ^{AB} [8, 14]. Figiel et al. directly fit $f_{\text{solv},\text{H}_2\text{O}} = 1.5$ to NaBr mean-ionic-activity-coefficient data [5, Eqns. 8–10 and Tables 2–3].

Neutral-solvent permittivities. The retained water correlation is the Archer–Wang approximation transferred through Uyan’s implementation and gives $\epsilon_{r,\text{H}_2\text{O}}(298.15 \text{ K}) = 78.4104$. This is close to, but not identical with, the high-accuracy 0.1-MPa value 78.3752 from Pátek et al.; the difference 0.0352 exceeds that reference’s stated uncertainty of about 0.01 [13, Eq. 11 and Tables 7–8]. Pátek’s correlation ends at 383.15 K, whereas this diagnostic reaches 393.15 K, so it cannot validate the upper endpoint. The retained $\epsilon_{r,\text{MEA}} = 32.0000$ is also a confirmed current foundation: Afify and Sweatman report an experimental literature average of 32.8 ± 2.1 at 298 K, and the direct dielectric-relaxation measurements of Agieienko and Buchner give 30.45 for neat MEA at 298.15 K [1, Table II] [2, p. 2314]. The displayed 32.0000 is the retained rounded working representative within that evidence range, not an exact measured value or temperature correlation and not a value reported by Baygi and Pahlavanzadeh. Baygi supports the other MEA PC-SAFT coordinates in Table 1 but contains no dielectric measurement or permittivity correlation.

All six ions. One segment, the signed integer charge, and $f_{\text{solv},i} = 1$ are model conventions rather than fitted single-ion properties. Figiel et al. set every pure-ion solvation factor to one because a hypothetical pure-ion system contains no solvent shell [5]. These conventions are Class 1 for this model definition.

Neutral binary interactions. The current $k_{\text{CO}_2,\text{H}_2\text{O}} = 0.0132628791769$ is the accepted constant fit to 39 qualified Kiepe rows with the induced association fixed [11]. The current $k_{\text{MEA},\text{H}_2\text{O}} = -0.0735274987499$ is the accepted 2B fit to 25 admitted Cai rows [4]. Their calculation records and acceptance limits are owned by ePC-SAFT Issue 80; Class 1 here does not make the full nine-species packet accepted.

Protocol-fixed zero corrections. Uyan et al. used the same CO₂, HCO₃[−], CO₃^{2−}, H₃O⁺, and OH[−] species and stated that all binary parameters not separately supplied for the predictive reacting ePC-SAFT calculation were set to zero [15]. Accordingly, the four CO₂–trace-ion corrections and all six trace-ion–trace-ion corrections are Class 1 zeros for the current sparse foundation. Here $k_{ij} = 0$ means that the uncorrected ePC-SAFT combining rule is used; it does not mean that the physical interaction vanishes. These ten values remain outside the single-coordinate CO₂–MEA regression defined by upstream Issue 119.

4.2. Class 2: direct literature candidates

Non-adopted CO₂ permittivity. The source trail is resolved. Schick et al. define CO₂ as a solvent in their composition mixing rule and report

$$\epsilon_{r,\text{CO}_2}(T) = -0.0036T + 2.467, \quad T \text{ in K},$$

as a correlation developed from three source groups [14, Table 2, p. 4]: Leeke et al., the thermophysical report by Ely et al., and the gas-property measurements by Uhlig et al. The historical scalar is exactly this correlation at 293.0000 K:

$$-0.0036(293.0000) + 2.467 = 1.4122.$$

The earliest retained code copy confirms the same lineage with the adjacent comment “Schick 2022”; the paper’s accepted publication year is 2023.

This resolves the numerical origin but does not validate 1.4122 as a temperature-independent property. The Schick correlation gives 1.4117, 1.3937, 1.3397, 1.2677, 1.1957, and 1.1237 at 293.15, 298.15, 313.15, 333.15, 353.15, and 373.15 K, respectively. Leeke et al. measured CO₂ permittivity at 353 and 383 K over 100–300 bar, while independent measurements show that CO₂ permittivity depends on density and pressure as well as temperature. The Schick equation is therefore an ePC-SAFT modeling correlation, not a universal pure-fluid property.

Decision. The full-domain comparison in Section 11.1 selects Uyan solvent-only mixing, which excludes dissolved CO₂ from the dielectric mixing rule. Therefore neither 1.4122 nor the Schick temperature correlation is an active coordinate in Table 1. Both remain retained Class 2 comparison evidence. This decision does not alter the Class 1 molecular Gross–Sadowski row.

SSM+DS neutral solvation factors. Figiel et al. define f_{mix} using ions and solvents only, set pure-ion factors to one, and fit solvent-specific factors of 1.5, 1.4, and 1.6 for water, methanol, and ethanol, respectively [5, Eqns. 8–10 and Table 2]. They also report that 1.5 represents all three studied solvents at low ion concentration. On that evidence, $f_{\text{solv,MEA}} = 1.5$ is the primary proposed value for the first MEA SSM+DS screen: MEA contributes to the solvent environment, is polar and associating, and 1.5 is central to the observed solvent range. The exact bundle evaluated in this notebook still has $f_{\text{solv,MEA}} = 1.0$, so Table 3 now reports 1.0 rather than silently presenting the untested 1.5 proposal as a live value. Both are Class 2 assumptions because no MEA activity data were used to fit either one.

The published mixing rule includes ions and solvents; no other component type was present in Figiel’s systems. Accordingly, $f_{\text{solv,CO}_2} = 1$ is retained as a Class 2 role-based assumption meaning “no neutral-solute shell increment,” not as a published CO₂ property. Its influence should be checked cheaply, but a dedicated fit is low priority unless that sensitivity is material. A current Engine-source inspection finds that the implemented composition average sums all supplied components without a solvent-role mask. Thus the intended solvent-only interpretation and the current implementation must be compared before fitting a CO₂ factor.

Trace-ion molecular rows. The σ_i and ϵ_i/k values for HCO₃[−], CO₃^{2−}, H₃O⁺, and OH[−] are the direct Held/Uyan aqueous-ion rows carried into later ePC-SAFT work [9, 15, 16]. They are credible literature values but have not been independently ranked for reactive aqueous MEA.

Non-adopted ionic dielectric transfers. The former common ionic-region $\epsilon_{r,i} = 8$ and c_i values are direct SSM+DS literature coordinates retained in the comparison provenance, not active MEA-system parameters under the selected solvent-only mixing rule [5, 16]. The H₃O⁺ Born diameter $d_B = 1.2180 \text{ \AA}$ is also directly supported by Figiel et al. [5].

Water–ion interactions. The literature mapping retains $k_{\text{H}_2\text{O,HCO}_3^-} = 0$, $k_{\text{H}_2\text{O,CO}_3^{2-}} = -0.25$, $k_{\text{H}_2\text{O,H}_3\text{O}^+} = 0.25$, and $k_{\text{H}_2\text{O,OH}^-} = -0.25$ [15, 16]. The bicarbonate zero is a literature assumption, not a confirmed physical exclusion; Wangler later fitted nonzero bicarbonate interactions in another electrolyte application. All four therefore remain Class 2 for MEA.

CO₂–MEA. Baygi and Pahlavanzadeh used $k_{\text{CO}_2,\text{MEA}} = 0$ and explicitly interpreted zero as use of the uncorrected PC-SAFT combining rule, not absence of molecular interaction [3]. That exact-species result makes zero the direct literature candidate. It remains Class 2 because their non-electrolyte association model is not the current induced,

nine-species ePC-SAFT foundation. The bounded screen in Section 7 retains zero as the calibration baseline and -0.1957268589 only as a second optimizer start. Neither value is promoted as a fitted result.

Other zero binary corrections. Uyan et al. state that binary parameters not separately supplied for the predictive reacting ePC-SAFT calculation were set to zero [15]. The 18 additional zero entries in Table 4 therefore use the uncorrected combining rule as a literature-backed model convention. They are Class 2 rather than Class 4: the numerical choice is supported, but none is confirmed as an MEA-specific optimum or evidence that the physical interaction vanishes.

4.3. Class 3: project-fitted or derived values

The current $\epsilon_{\text{CO}_2}/k = 173.440\,25\text{ K}$ is a 2.5% increase from the Gross–Sadowski start. It was selected jointly with R4 by the bounded campaign in Section 7. Both coordinates are Class 3 because they are project-fit values rather than values copied from a paper.

The May 2026 local-speciation fit varied seven coordinates simultaneously: σ_i , ϵ_i/k , and $d_{\text{B},i}$ for both MEA^+ and MEACOO^- , plus $k_{\text{MEA}^+, \text{MEACOO}^-}$. It used 8 states and 22 residuals from Böttinger, Jakobsen, and Matin data. Overall log-residual RMSE changed only from 0.2714 to 0.2677, and two final evaluations failed. Accordingly, the six displayed ion parameters and the -0.0020 interaction are Class 3 because they are project-fit outputs. Git commit 897bb6e preserves the exact fit values and diagnostics.

The OH^- Born diameter $d_{\text{B}} = 3.0811\text{ Å}$ is a local Born hydration-energy inversion retained under `analyses/reactive_epcsaft_parameter_evidence/results/oh_born_derivation/`. It remains Class 3 until its source assumptions and transfer checks are accepted. Every Debye–Hückel diameter is excluded from the tables and this classification inventory because it is deterministically calculated as $0.88\sigma_i$.

4.4. Class 4: missing values represented by defaults or analogs

The HCO_3^- and CO_3^{2-} Born diameters of 3.0000 Å are Class 4 seeds. The retained trace-carbonate check left both values at their starts with no residual improvement, so it did not convert the defaults into fitted parameters. The former $c_{\text{MEA}^+} = 2.6000$ and $c_{\text{MEACOO}^-} = 7.0100$ defaults are retained only as non-adopted SSM+DS provenance; the active bundle does not use them.

The matrix diagonal is gray and unclassified because self-pairs do not use a binary interaction coefficient.

5. Induced-association lineage and correlations

The CO_2 molecular row is not a new induced-association fit. Gross and Sadowski report $m = 2.0729$, $\sigma = 2.7852\text{ Å}$, and $\epsilon/k = 169.21\text{ K}$ in their original PC-SAFT table [6]. Pabsch et al. retain those values, add a 1/1 induced 2B site topology with zero CO_2 self-association energy and $\kappa^{AB} = 0.0450$, and fit the CO_2 –water binary term [12, Tables 2 and 6]. Schick et al. carry the same induced row forward [14, Tables 1 and 2]. The retired $m = 2.079$ value is the unsupported value; 2.0729 is the source-correct Gross–Sadowski value.

The retained water correlations are

$$\sigma_{\text{H}_2\text{O}}(T) = 2.7927 + 10.11 \exp(-0.01775T) - 1.417 \exp(-0.01146T), \quad (1)$$

$$\epsilon_{r,\text{H}_2\text{O}}(T) = 78.41043478578715 - 0.36133766233691506(T - 298.15) \quad (2)$$

$$+ 0.00076555618295(T - 298.15)^2. \quad (3)$$

The retained CO_2 scalar is a 293 K snapshot of the Schick correlation $\epsilon_{r,\text{CO}_2}(T) = -0.0036T + 2.467$, not a constant supported across the calculation range [14, Table 2, p. 4]; see Section 4.2.

The current bundle uses the selected **reciprocal induced CO_2 – H_2O association set**. CO_2 , MEA, and H_2O each carry one donor site a and one acceptor site b . Upstream PR #132 independently selected MEA 2B over 3B and 4C using the

admitted Cai MEA–water rows. The CO₂–water interaction is physical; reactive carbamate formation remains in the reaction network.

Pair	Active edges	ϵ^{AB}/k (K)	κ^{AB}	Meaning
MEA–MEA	$a:b, b:a$	2586.30	0.03747	Ordinary MEA self-association.
H ₂ O–H ₂ O	$a:b, b:a$	2425.70	0.04509	Ordinary water self-association.
CO ₂ –H ₂ O	reciprocal a – b cross edges	1212.85	0.04509	Selected induced association: CO ₂ (a)–H ₂ O(b) and CO ₂ (b)–H ₂ O(a).
MEA–H ₂ O	reciprocal a – b	2506.00	0.0411037990944876	Combined cross-association rule.
CO ₂ –CO ₂	none	—	—	Self-association disabled.

6. Reaction correlations

For R1–R3, the Austgen source form is $\ln K_{r,\text{source}} = A + B/T + C \ln T + DT$. Before constructing an Engine problem, the replay evaluates

$$\ln K_{r,\text{common}}(T) = \ln K_{r,\text{source}}(T) + \Delta_r, \quad r \in \{\text{R1}, \text{R2}, \text{R3}\}, \quad (4)$$

with $(\Delta_1, \Delta_2, \Delta_3) = (8.0330699846, 4.0165349923, 4.0165349923)$. These are the audited shifts from the Austgen mixed mole-fraction reference to aqueous–molality–infinite–dilution–water–v1. R4 and R5 already retain their declared common–molality forms. All five reactions remain fixed while physical parameter blocks are qualified.

ID	A	B	C	D	Domain (K)	Source definition
R1	132.899	–13445.9	–22.4773	0	273.15–498.15	Water autoprotolysis.
R2	231.465	–12092.1	–36.7816	0	273.15–498.15	CO ₂ hydration to bicarbonate.
R3	216.049	–12431.7	–35.4819	0	273.15–498.15	Bicarbonate dissociation.
R4	$\ln K = 3.3515778 - 1895.3/T$			293.15–393.15 ^a		Carbamate hydrolysis; joint fit.
R5	$\ln K = -\ln(10)[2677.91/T + 0.3869 + 0.0004277T]$			273.15–323.15		MEA ⁺ acid dissociation.

^aThe source correlation was qualified over 293.15–323.15 K. The displayed fit evaluates it through 393.15 K. R2 uses Austgen’s printed $A = 231.465$, which is also the value encoded in the validated state packet; the later 231.456 transcription is retained only as a documented source discrepancy.

7. Corrected replay and parameter-start decision

7.1. Verified packet-construction defect

The upstream reaction audit records the shifts in Eq. (4), but the retained exporter evaluates the raw Austgen R1–R3 correlations and writes those values into records already labeled with the common aqueous–molality standard state. The immutable state packet is preserved unchanged; the notebook replay applies the three shifts before the Engine mapping is constructed. The affected file and Engine commit are identified in Section 12. This distinguishes a packet-export error from a reaction-model or Engine-transfer error.

7.2. Prior ion-specific baseline and reaction screen

The corrected former ion-specific foundation evaluates every one of the 44 speciation states and 76 of the 79 pressure states. Material balances, reaction affinities, and pressure residuals remain below 3.0×10^{-10} in their reported infinity norms. The model therefore solves the stated equilibrium problem accurately where it evaluates; the comparison with observations tests model validity, not nonlinear-solver convergence.

Candidate	N_p	median $p_{\text{model}}/p_{\text{data}}$	pressure RMS factor	N_x	species MAE	species RMS factor
Current R4/R5, corrected R1–R3	76/79	13.89	16.71	44/44	0.00257	1.94
Böttinger R4, corrected R1–R3	70/79	10.25	13.27	44/44	—	2.96

The former ion-specific foundation reproduces the measured species pattern within a median multiplicative factor of 1.065, but its pressure is high by a median factor of 13.89. Böttinger’s common-molality R4 lowers pressure by only 15% on the 70 paired evaluable rows, worsens species agreement, and adds six pressure non-evaluations. Its R5 is numerically indistinguishable from Bates–Pinching over the tested range (less than 1.8×10^{-5} difference in $\ln K$). Neither reaction substitution resolves the pressure discrepancy.

The pressure bias is also temperature structured:

T (K)	313.15	333.15	353.15	373.15	393.15
median $p_{\text{model}}/p_{\text{data}}$	26.75	23.71	7.27	11.03	6.82

7.3. Bounded physical-interaction screens

The constant binary screens used 18 pressure states common to every candidate and six representative speciation states spanning the available temperatures.

Coordinate and value	median pressure factor versus current	species scaled RMSE	Interpretation
$k_{\text{CO}_2, \text{MEA}} = -0.1957268589$	0.9978	0.2536	Retain only as Issue #119’s second optimizer start.
$k_{\text{CO}_2, \text{MEA}} = +0.1015997042$	1.0010	0.2542	Old fit used the unconverted reaction packet; not transferable.
$k_{\text{CO}_2, \text{MEA}} = -1$	0.9841	0.2554	Diagnostic bound remains too weak and adds a species non-evaluation.
$k_{\text{CO}_2, \text{H}_2\text{O}} = 0$	0.7838	unchanged	Still leaves a 7.76 median pressure ratio and conflicts with the independent CO_2 –water qualification.
$k_{\text{CO}_2, \text{H}_2\text{O}} = 0.016724$	1.0661	unchanged	Pabsch’s 313 K value worsens reactive pressure relative to the current constant fit.

The verified conclusion from this earlier bounded screen was that no retained or directly transferable literature bundle presently gives both a reasonable nine-species speciation replay and a reasonable CO_2 pressure replay under the selected explicit-ion structure. That screen therefore defined the then-defensible calibration start as the corrected R1–R3 basis, Tong–Aroua R4, Bates–Pinching R5, and $k_{\text{CO}_2, \text{MEA}} = 0$; the legacy negative value was only a second start. Section 11.1 superseded its dielectric choice, and Section 7.4 supersedes its R4 and CO_2 dispersion coordinates.

Baygi’s reported pressure performance cannot be transferred by copying its parameters. That calculation used 3B MEA, 4C water, nonassociating CO_2 , ideal Smith–Missen chemistry, and $k_{\text{MEA}, \text{H}_2\text{O}} = -0.052$ [3]. The present foundation uses 2B MEA, Held 2B water, induced CO_2 –water association, explicit electrolyte activities, and $k_{\text{MEA}, \text{H}_2\text{O}} = -0.0735275$. The two parameter sets do not represent the same model.

7.4. Bounded joint R4– CO_2 campaign and selection

The September 2026 campaign used direct coupled Engine reactive bubble-point and homogeneous-liquid equilibrium solves throughout. A native-regression smoke test on six states and two coordinates required 129 seconds for only two iterations and did not produce a usable fit, so the retained study used parallel, bounded direct evaluations rather than an ad hoc SciPy equilibrium surrogate. The sequence comprised a centered sensitivity screen, R4 pivot–slope grids, complete-data validation, a local R4– CO_2 refinement, and a final slope-boundary check. No numerical score was treated as an accept/reject gate; candidates were compared on error, convergence coverage, temperature residual pattern, and speciation tradeoffs.

The centered screen found no numerical response to the inactive $f_{\text{solv}, \text{HCO}_3^-}$ coordinate, negligible benefit from the tested bicarbonate Born movement, and worse pressure/speciation tradeoffs from the tested R5 direction. R4 was the only consistently useful reaction

block. Reparameterizing R4 around 333.15 K separated its local pivot from its temperature slope; a small CO₂ dispersion-energy increase then reduced the remaining common pressure bias.

The selected current bundle is

$$\ln K_{R4} = 3.3515778178 - 1895.3/T, \quad \epsilon_{\text{CO}_2}/k = 173.440\,25\,\text{K}.$$

All other coordinates remain equal to the previous bundle. The final comparison is:

Bundle	pressure coverage	pressure log RMSE	balanced pressure RMSE	speciation coverage / log RMSE
Previous	125/161	0.3974	0.3569	131/131 / 0.2664
Selected	130/161	0.3540	0.3185	131/131 / 0.2658

Pressure log RMSE improves at every temperature: 0.5047 to 0.4445 at 40 °C, 0.2888 to 0.2683 at 60 °C, 0.2491 to 0.2314 at 80 °C, 0.2315 to 0.1971 at 100 °C, and 0.4280 to 0.3809 at 120 °C. The sign pattern remains diagnostic: mean log residuals are +0.4014 and +0.1698 at 40 and 60 °C, then −0.1052, −0.0552, and −0.3661 at 80, 100, and 120 °C. Thus the current model still overpredicts low-temperature pressure and underpredicts high-temperature pressure. Extending the R4 slope to its present bound did not improve the screen and began losing speciation evaluations, so the bounded direction is closed rather than extrapolated farther.

A separate SciPy reaction-extent spot check at v1e_obs_0128, using Engine composition derivatives at the Engine bubble pressure, returned to the Engine liquid composition within 3.31×10^{-13} in mole fraction and a maximum reaction-affinity residual of 3.31×10^{-8} . This verifies that spot-check root only; no SciPy-derived parameter score is retained.

8. Nine-species equilibrium replay

Figure 1 retains the pre-campaign smooth-grid visual baseline generated with the corrected replay and retained PR #134 state packet. The bounded campaign did not rerun the additional smooth grid because the 131 observed targets were sufficient for candidate selection; exact selected-bundle statistics follow in Section 9. The single 20 °C, 30 wt% MEA panel is the densest retained same-condition comparison: 149 positive values for the five principal species and the MEA+MEA⁺ observation operator from 39 Böttinger, Jakobsen, and Matin states. These include direct, aggregate, inferred, and correlated values. The 17 exact reported-zero records remain in the retained observation snapshot but are omitted from the logarithmic plot. The broken logarithmic scale gives most of the panel to the principal-species range from 5.6248×10^{-5} , 30% below the lowest positive observed or modeled HCO₃[−] value, to 0.1134, only 3% above the displayed maximum. A compact 10^{-10} to that cutoff trace panel retains the low CO₂ curve. No retained positive CO₂ or HCO₃[−] observation is below 10^{-10} ; the values shown there are calculated molecular CO₂, CO₃^{2−}, H₃O⁺, and OH[−] remain in every equilibrium calculation; the eight reported carbonate values also remain in the figure source snapshot, but trace species are omitted from the visual. Dashed curves connect direct pinned-Engine evaluations from loading 0.075 to 1.0; they are not fitted or interpolated curves.




Figure 1. Pre-campaign smooth-grid diagnostic for the full nine-species liquid-equilibrium replay at 20 °C and 30 wt% MEA. Symbols show all positive retained Böttinger, Jakobsen, and Matin values for the displayed series; symbol shape identifies source and color identifies species. The broken logarithmic axis expands the 5.6248×10^{-5} – 0.1134 principal-species region and compresses the 10^{-10} – 5.6248×10^{-5} range. Dashed lines connect direct Engine evaluations of the immediately preceding bundle through loading 1.0; they are retained for qualitative context and are not the authority for the selected- bundle statistics.

These states are homogeneous reactive-liquid solves, not ideal-reaction or postprocessed SciPy curves. Each state couples the five reaction constraints to ePC-SAFT chemical potentials for all nine species at fixed temperature and pressure. A 2026-09-01 disposable timing check on this machine, excluding process startup and model construction, measured 0.935 s/state for a warm five-state second pass with the retained wheel. The same five feeds took 2.260 s/state with a disposable wheel built from Engine main after only the required provider-to-EOS request-schema renames. This is a route-specific diagnostic, not a general Engine benchmark: recent GREPE and regression speedups do not establish a speedup for this homogeneous-liquid owner.

9. Carbon-dioxide pressure replay

Figure 2 shows all 161 active-v1 30 wt% MEA pressure observations from Aronu, Hilliard, Idris, Jou, Mamun, and Xu at 40–120 °C. The selected replay evaluated 130 states and returned 31 explicit numerical convergence failures. Its full-data $\log_{10}$ RMSE is 0.3540 and the equal-temperature value is 0.3185. Lines interpolate the 130 evaluated selected-bundle observation states; no additional smooth-grid Engine campaign was required, and every experimental point remains shown.




Figure 2. Carbon-dioxide partial pressure for 30 wt% MEA. Colors identify temperature, symbols identify experimental source, and lines are shape-preserving log-response interpolations of newly calculated active-bundle states within each evaluated loading range; every retained observation remains visible even when its corresponding model state did not evaluate.

9.1. Current-bundle fit and coverage statistics

These are replay/agreement statistics for the exact bundle in Tables 1–4. Log residuals are $\log_{10}(y_{\text{model}}/y_{\text{obs}})$; the RMSE factor is $10^{\text{RMSE}_{\log_{10}}}$. Native pressure errors are in kPa and native speciation errors are mole fractions.

Family	Coverage	log RMSE	RMSE factor	MAPE (%)	MAE	Native RMSE
Pressure	130/161	0.3540	2.259	105.04	18.6330	99.9354
Speciation targets	131/131	0.2658	1.844	16.88	0.002643	0.003426

Pressure performance and, especially, evaluability vary strongly by temperature:

T (°C)	Coverage	log RMSE	RMSE factor	MAPE (%)	Median APE (%)
40	56/59	0.4445	2.783	176.41	178.56
60	23/24	0.2683	1.855	75.68	38.99
80	21/21	0.2314	1.704	35.43	38.26
100	18/18	0.1971	1.574	34.71	29.15
120	12/39	0.3809	2.404	55.63	59.32

The principal measured-species breakdown shows that bicarbonate, not the amine/carbamate coordinates, dominates the speciation log score:

Target	Coverage	log RMSE	RMSE factor	MAPE (%)
MEA	19/19	0.0575	1.141	10.07
MEA H^+	19/19	0.0498	1.121	11.38
MEA + MEA H^+	25/25	0.0221	1.052	3.36
MEACOO $^-$	44/44	0.0317	1.076	6.07
HCO $_3^-$	24/24	0.6154	4.125	60.53

The complete machine-readable handoff is retained in the results directory:

- row-level observed/predicted values: `current-best-fit-residuals.csv`;
- grouped overall, temperature, source, species, and temperature–source statistics: `current-best-fit-statistics.csv`;
- failed coordinates and diagnostics: `figure-calculation-failures.csv`.

The bundle can be used for absorption-column calculations. The caller should retain explicit non-evaluation for states that do not solve; at 120 °C, 12/39 pressure states evaluated.

10. What pressure constrains and what remains unresolved

At phase equilibrium the molecular carbon-dioxide fugacities satisfy

$$f_{CO_2}^V = f_{CO_2}^L. \quad (5)$$

An activity/Henry representation writes the same comparison schematically as

$$y_{CO_2} \phi_{CO_2}^V P = x_{CO_2} \gamma_{CO_2}^L H_{CO_2}. \quad (6)$$

inference: pressure observations therefore constrain the liquid molecular- CO_2 chemical potential together with loading, material balances, and the reaction network. They are valuable indirect speciation evidence, but they do not separately identify free x_{CO_2} , liquid nonideality, reaction constants, association, binary interactions, dielectric terms, or solvation factors. Direct species observations and pressure observations are combined because they constrain different projections of the same equilibrium state; pressure is not a substitute for direct speciation.

verified: the retained replay does not assume an ideal vapor. Its state packet declares a neutral incipient-vapor phase and reads `phase.partial_pressure` from the pinned ePC-SAFT equilibrium result. By contrast, the current public IDAES column property files declare an ideal liquid/Henry-law MEA solvent and an ideal vapor; they are not the modified-eNRTL model from a separate literature study. The public solvent file retains the three apparent neutral components CO_2 , MEA, and H_2O plus the three ionic species MEA^+ , MEACOO^- , and HCO_3^- [official IDAES solvent file]; the vapor file explicitly selects the ideal equation of state [official IDAES vapor file].

10.1. Reduced-chemistry diagnostic

The six-species reduction retains CO_2 , MEA, H_2O , MEA^+ , MEACOO^- , and HCO_3^- . To isolate species-set structure without importing IDAES empirical equilibrium constants, derive its two net constants from the current five-reaction basis:



The first relation is the exact R2–R4–R5 projection already recorded in MEA Issue #72. This makes the comparison a reaction-space projection of the same model rather than a change of both chemistry and equilibrium constants.

verified: across the 44 retained nine-species calculations, the largest mole fractions are 1.0307×10^{-2} for HCO_3^- , 1.1687×10^{-4} for CO_3^{2-} , 3.3641×10^{-6} for OH^- , and 8.9480×10^{-12} for H_3O^+ . The largest omitted net-charge contribution $(2x_{\text{CO}_3^{2-}} + x_{\text{OH}^-} - x_{\text{H}_3\text{O}^+})/x_{\text{MEA}^+}$ is 0.01949 at Böttinger state 030. These values are calculated directly from `figures/speciation/output/speciation-model.csv`.

The complete matched comparison is retained under `../enrtl_six_species_ideal_comparison/`; its Quarto source and rendered PDF are the detailed calculation record. In its ideal-liquid lane, all 232 unique canonical coordinates and 464 model rows converged with a maximum equation residual of 1.9895×10^{-13} . The largest difference among the six shared species was 6.5100×10^{-3} mol per mol initial MEA, and carbonate dominated the omitted inventory. In its fixed- T , P Engine/SciPy lane at 100 kPa, the carbonate amounts at 293.15, 313.15, and 323.15 K were 2.99×10^{-4} , 1.56×10^{-4} , and 1.10×10^{-4} mol per mol initial MEA; the largest shared-species changes were approximately 2.9×10^{-4} mol per mol initial MEA. That lane is a liquid-composition calculation, not a bubble-pressure calculation.

verified structural result: when the six- and nine-species states are both projected to the same conserved apparent totals, their neutral apparent pressure inputs agree to 1.321165×10^{-14} . The neutral `flashTQ` route converged for only 1 of 161 pressure observations and predicted 2038.38 kPa for an observed 0.002 02 kPa. A neutral apparent flash therefore cannot reveal the internal reaction or ionic parameters; the pressure calculation must keep the reactive species allocation coupled to the EOS.

The companion sensitivity survey evaluated 143 parameter identities grouped into 113 continuous directions with 128 Latin-hypercube samples, of which 123 were usable. Its strongest recurring directions were R5 a_K , R4 b_K , the water segment-diameter temperature terms, MEA^+ dispersion and ionic terms, and selected CO_2 –water, carbonate–water, MEA– MEA^+ , MEA^+ –water, carbamate–hydronium, and bicarbonate– CO_2 binary interactions. Those numerical ranks belong to parameter packet `0e7c5ac63c078be9d32c15f316ecdb2b0de67de6f495358ab5955450f0271c6a` and Engine-wheel hash `05cca73463b9b44dcb3a77dfd239b767ec63aafce7d18b85b2a66b80773847a5`; the separate fixed- T , P lane used packet `12e380a27eb615e25bae580f8c1ec852020879917b08fa16a5821ae76311550d`. Neither is the active bundle in Tables 1–4, so the ranking must be replayed with the active parameter file before its numerical ordering is used in the manuscript.

inference: the three omitted trace ions have small direct abundance over this replay, which makes a six-species ablation useful as a cheap conditioning test. It does not prove that they are dispensable: the z_i^2 weighting gives carbonate four times the per-mole ionic-strength contribution of a monovalent ion, and the omitted species preserve carbonate, autoprotolysis, pH, charge, and extrapolation structure. Removing them also removes weakly identified ionic parameters, so an apparent fit improvement may come from reduced flexibility rather than better physics.

Retained finding	Present use	Remaining action
Exact R2–R4–R5 and R2–R5 projection	Methods description for a controlled six-species ablation	None for the algebra; cite the retained reaction table.
Ideal and fixed- T , P species comparison	Evidence that carbonate is the dominant omitted species and that the reduction is not simple deletion	Repeat the coupled bubble calculation with the active bundle before using pressure-error conclusions.
Neutral apparent-pressure invariance	Methods/diagnostic explanation of why total absorbed carbon cannot be passed directly to a neutral CO_2 flash	Keep as a route-rejection result; do not present the 1/161 flash as a model fit.
Packet-specific sensitivity survey	Prioritizes the first active-bundle sensitivity directions	Replay the same screen with the active parameter mapping and wheel before reporting ranks or intervals.

11. Interpretation

The R4– CO_2 update reduces pressure error while preserving complete observed- speciation coverage, but the residual changes sign with temperature and 120 °C pressure coverage remains only 12/39. This identifies a remaining temperature-dependent mismatch; it does not justify opening R1–R3, R5, or the ion block as compensating coordinates.

The Class 2 $k_{\text{CO}_2, \text{MEA}} = 0$ is a direct literature candidate for the uncorrected combining rule, not a fitted result or a confirmed absence of interaction. The bounded screen shows that one constant correction is too weak and too temperature uneven to explain the observations. Likewise, loaded-MEA pressure alone cannot identify $f_{\text{solv}, \text{CO}_2}$ or $f_{\text{solv}, \text{MEA}}$ because those working assumptions are confounded with cross association, dielectric response, reaction, and other physical interactions.

Historical diagnostics remain outside the bundle:

- legacy $k_{\text{CO}_2, \text{MEA}} = -0.1957268589$;
- alternate MEAH^+ $\sigma = 2.753332853$ and MEACOO^- $\sigma = 3.017079292$; and
- trace-only bicarbonate Born-diameter movement toward 6.802 94 Å, with CO_3^{2-} left near 2.997 44 Å.

These values identify sensitivity or lineage; none is adopted. The trace-carbonate result does not establish independent identifiability.

11.1. Full-domain permittivity selection

verified: a full retained-state Engine comparison evaluated the former ion-specific baseline, Schick component mixing, and Uyan solvent-only mixing without fitting any parameter. Each candidate attempted all 79 pressure states from 313.15 to 393.15 K and all 44 speciation states from 293.15 to 353.15 K. All variants used the same base vector, corrected R1–R3 constants, reciprocal induced association, state packet, and pinned Engine wheel.

The Schick transfer uses the component-mixing relation

$$\epsilon_r = \left(\sum_{j \in \text{neutral}} \epsilon_{r,j}(T) w_j^{\text{neutral}} \right) x_{\text{neutral}} + \sum_{j \in \text{ion}} 8x_j, \quad (9)$$

with CO_2 included and $\epsilon_{r, \text{CO}_2}(T) = -0.0036T + 2.467$ [14, Eq. 18 and Table 2], while retaining the current bundle’s water correlation rather than replacing it with Schick’s water input. The Uyan transfer uses the earlier CO_2 - and ion-independent solvent relation

$$\epsilon_r = \epsilon_{r, \text{H}_2\text{O}}(T) w_{\text{H}_2\text{O}}^* + 32 w_{\text{MEA}}^*, \quad (10)$$

where the starred mass fractions are renormalized over MEA and water only, following Uyan et al.’s Eq. 5 with MEA and its retained constant $\epsilon_r = 32$ substituted for Uyan’s temperature-dependent MDEA input [15, Eq. 5]. Neither transfer

reproduces the source paper’s full parameterization.

Figure 3 is the primary comparison view. Every marker is a successfully evaluated observed–predicted target and the dashed diagonal is exact agreement. Legend RMSE values are first calculated within each temperature and then averaged equally across temperature so a densely sampled isotherm cannot dominate selection.




Figure 3: Full-domain parity view for the former ion-specific baseline, Schick component mixing, and Uyan solvent-only mixing. Legend values are temperature-balanced positive-target $\log_{10}$ RMSE and evaluated-state coverage. Uyan has the lowest pressure error; Schick and Uyan have nearly identical speciation error. No candidate was refitted.

The exact aggregate values underlying the visual comparison are:

Quantity	Ion-specific	Schick mixing	Uyan solvent-only
Pressure states	76/79	69/79	69/79
Pressure balanced $\log_{10}$ RMSE	1.0879	0.5053	0.3487
Speciation states	44/44	44/44	44/44
Speciation balanced $\log_{10}$ RMSE	0.1803	0.1670	0.1668

At every retained pressure temperature, Uyan improves the $\log_{10}$ RMSE relative to the ion-specific baseline: 0.5331, 0.3579, 0.1947, 0.2594, and 0.3007 from 313.15 through 393.15 K, compared with 1.4383, 1.2618, 0.8917, 0.9625, and 0.7313. Its 44-state speciation score is within 0.2% of Schick’s and better than the former baseline. It therefore passes the preregistered per-temperature pressure guardrail and the 10% speciation guardrail.

decision: retain Uyan solvent-only mixing in the active bundle. The subsequent bounded campaign fits R_4 and ϵ_{CO_2}/k under this formulation; Schick and the former ion-specific form remain retained comparison evidence rather than live coordinates in Tables 1–4.

12. Current conclusions and next experiments

The selected physical structure includes reciprocal induced $\text{CO}_2\text{-H}_2\text{O}$ association and excludes CO_2 self-association. The active foundation uses corrected R1–R3, Uyan solvent-only dielectric mixing, and $k_{\text{CO}_2,\text{MEA}} = 0$. The following steps provide the shortest discriminating path. Items below are proposed work, not retained results.

1. Diagnose the remaining temperature sign change using the retained row-level residuals. Test a held-out-source or held-out-temperature replay before changing another coordinate. Do not extend the R4 slope beyond the tested bound: that direction ceased improving the screen and began losing speciation evaluations.
2. Complete the remaining coupled bubble-pressure ablation with the active parameter file: nine versus six liquid species, crossed with ePC-SAFT versus ideal vapor. The retained comparison has already established the exact six-species reaction projection, ideal-liquid behavior, fixed- T, P liquid behavior, and the failure of the neutral apparent-flash shortcut. Report state evaluability, pressure and species metrics, charge/balance closure, and changes by pressure range and temperature.
3. Admit independent physical-solubility evidence before expanding the model. Jiru et al. measure physical CO_2 solubility in MEA–water by the N_2O analogy over 298.15–323.15 K [10]; Hartono et al. report N_2O solubility in unloaded and loaded MEA [7]. Wong et al. directly measure dissolved molecular CO_2 and Henry behavior [17], but every retained row contains $0.517 \text{ mol kg}^{-1} \text{ NaClO}_4$, which the nine-species model omits; use those rows as an external challenge unless the salt is represented explicitly.
4. Keep Uyan solvent-only mixing as the baseline while testing an SSM+DS extension against the retained Schick component-mixing challenge. Compare MEA $f_{\text{solv}} = 1.0, 1.4, 1.5, 1.6$ with 1.5 as the primary start. Only if the CO_2 treatment remains material should a separate CO_2 factor be considered. Evaluate both speciation and pressure so a compensating pressure improvement cannot be mistaken for physical validation.
5. With R1–R5, the ion block, and independently accepted $k_{\text{CO}_2,\text{H}_2\text{O}}$ fixed, compare only physical dissolved- CO_2 alternatives that the new data can distinguish: a $\text{CO}_2\text{-MEA}$ association edge, a physically identified solvation/activity contribution, and any richer binary correction justified by the temperature and composition coverage. Do not combine coordinates that the admitted observations cannot identify.
6. Require simultaneous comparative improvement in pressure, speciation, and convergence coverage, followed by held-out temperatures and sources before issuing an immutable packet. No unapproved numerical threshold is an acceptance gate.

unknown: whether ideal vapor, reduced chemistry, or neither materially changes the coupled bubble-pressure residual has not been tested with the active bundle.

Replace the current numerical foundation in place when a better MEA-owned regression or validation result passes the stated comparison gates. Every replacement must bind the Engine wheel and hash, parameter identities and evidence status, admitted rows and hashes, optimizer diagnostics, validation summaries, and the transfer decision. This PDF is the live calculation and interpretation record for the active bundle, including the reported fit statistics and state counts.

13. Provenance

The base numerical reconstruction uses the PR #134 file:

`validation/campaigns/2026-mea-parameter-estimation/analysis/joint-fit/parameters.json`

at commit 38e91823b6d4f26c1d549f07aaef24a089d8e16d. The selected live mapping is:


`results/selected-current-best-parameters.json`

Its SHA-256 is the parameter identity reported in Section 2. The comparison result records both the base and selected fingerprints.


Source	Role
ePC-SAFT Issue #80	Upstream full regression and immutable-packet campaign.
ePC-SAFT Issue #119	Bounded refinement scope and current foundation reconciliation.
ePC-SAFT PR #134	Closed, unmerged blocker-evidence change. It produced no parameter bundle; this notebook reconstructs only its frozen foundation vector.
Local component source audit	docs/ePC-SAFT/full-component-parameter-source-audit.md; species-level lineage and evidence gaps.
Local model-hierarchy review	docs/ePC-SAFT/amine-epcsaft-model-hierarchy-literature-review.md; selected nine-species structure and induced-association decision.
Retained parameter evidence	analyses/reactive_epcsaft_parameter_evidence/; supported-negative and diagnostic evidence.
Retained local-speciation fit	Git commit 897bb6e; the May 2026 8-state/22-residual principal-ion fit retained as Class 3 evidence awaiting promotion checks.
Reaction exporter defect	validation/campaigns/2026-mea-parameter-estimation/scripts/build_fit_input.py at Engine commit 38e91823b6d4f26c1d549f07aaef24a089d8e16d.
Corrected start comparison	results/parameter-start-comparison.csv; full corrected-foundation and Böttinger-R4 comparison statistics.
Bounded sensitivity screen	results/parameter-sensitivity-screen.csv; common-row CO ₂ -MEA and CO ₂ -water perturbation results used in Section 7.
Quick direct endpoint screen	results/quick-endpoint-perturbation-screen.csv and its JSON receipt; full reactive bubble-point endpoints and the bounded SciPy reaction-root spot check.
Best-in-slot campaign	results/best-in-slot-campaign/; centered sensitivity, R4 pivot-slope grids, full validations, boundary diagnostics, exact receipts, and the selected candidate's row-level Engine evaluations.
Permittivity model comparison	results/permittivity-formulation-comparison.json with full state and target CSVs; Figure 3 is rendered from the target CSV without new Engine evaluations.
Selected live mapping	results/selected-current-best-parameters.json; Uyan solvent-only formulation with the selected Class 3 R4-CO ₂ fit.
Trace-species abundance	figures/speciation/output/speciation-model.csv; retained 44-state calculated values summarized in Section 10.
Current IDAES comparison	Official public MEA solvent and MEA vapor property files; used only to define the proposed structural ablation.
Matched species comparison	../enrtl_six_species_ideal_comparison/notebook.qmd and ../enrtl_six_species_ideal_comparison/notebook.pdf; exact ideal projection, fixed- T , P Engine/SciPy comparison, corrected apparent-pressure diagnostic, parameter inventory, sensitivity survey, UQ checkpoints, and figures.

A. Additional-temperature speciation replays


The following panels apply the same species selection, source encoding, reported-zero omission, split-axis rule, parameter bundle, and pinned Engine route as Figure 1. They are supplementary temperature views rather than additional fits. At 40 °C, 74 positive plotted values cover 21 states and observed loading 0.11–0.99; at 60 °C, 21 values cover eight states and observed loading 0.131–0.766; at 80 °C, 14 values cover six states and observed loading 0.064–0.51. Every model grid continues to loading 1.0, and all 46 states evaluated at each temperature. Dashed lines connect those direct calculations.



Appendix Figure A1. Principal-species replay at 40 °C and 30 wt% MEA. Symbols retain every displayed Böttinger and Jakobsen observation; dashed lines are direct Engine calculations.



Appendix Figure A2. Principal-species replay at 60 °C and 30 wt% MEA. Symbols retain every displayed Böttinger observation; dashed lines are direct Engine calculations.



Appendix Figure A3. Principal-species replay at 80 °C and 30 wt% MEA. Symbols retain every displayed Böttinger observation; dashed lines are direct Engine calculations.

References

- [1] N. D. Afify and M. B. Sweatman. "Molecular Dynamics Simulation of Microwave Heating of Liquid Monoethanolamine (MEA): An Evaluation of Existing Force Fields". In: *The Journal of Chemical Physics* 148.20 (May 2018), p. 204513. ISSN: 0021-9606. DOI: 10.1063/1.5022585. (Visited on 09/01/2026).
- [2] Vira Agieienko and Richard Buchner. "What Is behind a Gas Stream Scrubbing Liquid? Monoethanolamine/Water Mixtures as Seen by Dielectric Relaxation Spectroscopy". In: *Physical Chemistry Chemical Physics* 26.3 (Jan. 2024), pp. 2312–2323. ISSN: 1463-9076. DOI: 10.1039/d3cp05027k. (Visited on 09/01/2026).
- [3] Sadjad Baygi and Hassan Pahlavanzadeh. "Application of the PC-SAFT Equation of State for Modeling CO₂ Solubility in Aqueous Monoethanolamine Solutions". In: *Chemical Engineering Research and Design* 93 (Jan. 2015), pp. 789–799. ISSN: 02638762. DOI: 10.1016/j.cherd.2014.07.017.
- [4] Zhenyun Cai, Rongjing Xie, and Zhaoli Wu. "Binary Isobaric Vapor-Liquid Equilibria of Ethanolamines + Water". In: *Journal of Chemical & Engineering Data* 41.5 (Jan. 1996), pp. 1101–1103. ISSN: 0021-9568. DOI: 10.1021/jc960118o. (Visited on 07/23/2026).
- [5] Paul Figiel, Gangqiang Yu, and Christoph Held. "Predicting Thermodynamic Properties of Ions in Single Solvents and in Mixed Solvents Using a Modified Born Term within the ePC-SAFT Framework". In: *Industrial and Engineering Chemistry Research* 64.18 (2025), pp. 9406–9418. ISSN: 15205045. DOI: 10.1021/acs.iecr.5c00475.

- [6] Joachim Gross and Gabriele Sadowski. "PC-SAFT: An Equation of State Based on a Perturbation Theory for Chain Molecules". In: *Industrial and Engineering Chemistry Research* 40.4 (Feb. 2001), pp. 1244–1260. ISSN: 08885885. DOI: 10.1021/ie0003887.
- [7] Ardi Hartono, Emmanuel Orji Mba, and Hallvard F. Svendsen. "Physical Properties of Partially CO₂ Loaded Aqueous Monoethanolamine (MEA)". In: *Journal of Chemical & Engineering Data* 59.6 (June 2014), pp. 1808–1816. ISSN: 0021-9568. DOI: 10.1021/je401081e. (Visited on 05/15/2026).
- [8] Christoph Held, Luca F. Cameretti, and Gabriele Sadowski. "Modeling Aqueous Electrolyte Solutions. Part 1. Fully Dissociated Electrolytes". In: *Fluid Phase Equilibria* 270.1-2 (2008), pp. 87–96. ISSN: 03783812. DOI: 10.1016/j.fluid.2008.06.010.
- [9] Christoph Held, Thomas Reschke, Sultan Mohammad, Armando Luza, and Gabriele Sadowski. "EPC-SAFT Revised". In: *Chemical Engineering Research and Design* 92.12 (Dec. 2014), pp. 2884–2897. ISSN: 02638762. DOI: 10.1016/j.cherd.2014.05.017.
- [10] Ying Jiru, Dag A. Eimer, and Yi Wenjuan. "Measurements and Correlation of Physical Solubility of Carbon Dioxide in (Monoethanolamine + Water) by a Modified Technique". In: *Industrial and Engineering Chemistry Research* 51.19 (May 2012), pp. 6958–6966. ISSN: 08885885. DOI: 10.1021/ie3002588.
- [11] Jörn Kiepe, Sven Horstmann, Kai Fischer, and Jürgen Gmehling. "Experimental Determination and Prediction of Gas Solubility Data for CO₂ + H₂O Mixtures Containing NaCl or KCl at Temperatures between 313 and 393 K and Pressures up to 10 MPa". In: *Industrial & Engineering Chemistry Research* 41.17 (July 2002), pp. 4393–4398. ISSN: 0888-5885. DOI: 10.1021/ie020154i. (Visited on 08/19/2026).
- [12] Daniel Pabsch, Christoph Held, and Gabriele Sadowski. "Modeling the CO₂ Solubility in Aqueous Electrolyte Solutions Using ePC-SAFT". In: *Journal of Chemical and Engineering Data* 65.12 (Dec. 2020), pp. 5768–5777. ISSN: 15205134. DOI: 10.1021/acs.jced.0c00704.
- [13] Jaroslav Pátek, Jan Hrubý, Jaroslav Klomfar, Monika Součková, and Allan H. Harvey. "Reference Correlations for Thermophysical Properties of Liquid Water at 0.1MPa". In: *Journal of Physical and Chemical Reference Data* 38.1 (Jan. 2009), pp. 21–29. ISSN: 0047-2689. DOI: 10.1063/1.3043575. (Visited on 09/01/2026).
- [14] Daniel Schick, Lena Bierhaus, Alexander Strangmann, Paul Figiel, Gabriele Sadowski, and Christoph Held. "Predicting CO₂ Solubility in Aqueous and Organic Electrolyte Solutions with ePC-SAFT Advanced". In: *Fluid Phase Equilibria* 567 (Apr. 2023), p. 113714. ISSN: 0378-3812. DOI: 10.1016/j.fluid.2022.113714. (Visited on 03/27/2026).
- [15] Mustafa Uyan, Georg Sieder, Thomas Ingram, and Christoph Held. "Predicting CO₂ Solubility in Aqueous N-methyldiethanolamine Solutions with ePC-SAFT". In: *Fluid Phase Equilibria* 393 (May 2015), pp. 91–100. ISSN: 03783812. DOI: 10.1016/j.fluid.2015.02.026.
- [16] Anton Wangler, Georg Sieder, Thomas Ingram, Manfred Heilig, and Christoph Held. "Prediction of CO₂ and H₂S Solubility and Enthalpy of Absorption in Reacting N-methyldiethanolamine /Water Systems with ePC-SAFT". In: *Fluid Phase Equilibria* 461 (2018), pp. 15–27. ISSN: 03783812. DOI: 10.1016/j.fluid.2017.12.033.
- [17] M. K. Wong, M. A. Bustam, and A. M. Shariff. "In Situ Measurement of Physical Solubility of Carbon Dioxide in Loaded Aqueous Monoethanolamine by Raman Spectroscopy". In: *Journal of Natural Gas Science and Engineering* 36 (Nov. 2016), pp. 305–313. ISSN: 1875-5100. DOI: 10.1016/j.jngse.2016.10.029. (Visited on 08/24/2026).