

Medications

None

Medical History

None

Musculoskeletal History

None

Musculoskeletal Family History

None

Musculoskeletal Surgery

None

Surgical History

None

Social History

Smoking status - Never smoker

Chief Complaints:

1. Right elbow pain, worse with gripping, 5 weeks.

HPI: This is a 43 year old male who:

Patient reports five weeks of right lateral elbow pain that began after increased racquet sport activity. Pain is worse with gripping, lifting objects with the palm down, and shaking hands. No numbness or tingling into the hand. No prior elbow injury.

Vitals:

Date	Taken By	B.P.	Pulse	Resp.	O2 Sat.	Temp.	Ht.	Wt.	BMI BSA
07/08/25	T. Nguyen, MA	124/78	68	14	99	98.3	72.0 in	195.0 lbs	26.4 2.05

Exam:

Comprehensive, Upper Extremity Neurovascular

Appearance: well nourished

Orientation: Alert and oriented to person, place, time.

Mood: mood and affect well-adjusted, pleasant and cooperative,

Right UE Pulses: normal radial pulse and good capillary refill

Left UE Pulses: normal radial pulse and good capillary refill

RUE Peripheral Sensation: intact to light touch throughout peripheral nerve distributions

LUE Peripheral Sensation: intact to light touch throughout peripheral nerve distributions

Elbow

Right Elbow Active ROM:

Flexion: 145 degrees.
Extension: 0 degrees.
Pronation: 80 degrees.
Supination: 80 degrees.

Left Elbow Active ROM:

Flexion: 145 degrees.
Extension: 0 degrees.
Pronation: 80 degrees.
Supination: 80 degrees.

Right Elbow Passive ROM:

Flexion: 145 degrees.
Extension: 0 degrees.
Pronation: 80 degrees.
Supination: 80 degrees.

Left Elbow Passive ROM:

Flexion: 145 degrees.
Extension: 0 degrees.
Pronation: 80 degrees.
Supination: 80 degrees.

Skin:

Right Elbow: skin intact, no rashes or lesions.

Skin:

Left Elbow: skin intact, no rashes or lesions.

Inspection:

Right Elbow: point tenderness over the lateral epicondyle, no effusion

Inspection:

Left Elbow: Normal alignment, no deformity, no tenderness, no warmth, no masses, no muscle atrophy, no crepitus

Strength:

Right Elbow Flexion: Strength: 5/5, normal muscle tone.
Right Elbow Extension: Strength: 5/5, normal muscle tone.

Strength:

Left Elbow Flexion: Strength: 5/5, normal muscle tone.
Left Elbow Extension: Strength: 5/5, normal muscle tone.

Stability:

Right Elbow: Stable

Special:

Right Elbow: Resisted wrist extension: positive, resisted middle finger extension: positive

Impression/Plan:

1. Elbow Pain, Right

Lateral epicondylitis, right elbow (M77.11)

Plan: Prescription.

ibuprofen 800 mg tablet PO

Sig: Take 1 po tid x 2 weeks with food

Quantity: 42 Tablet Refills: 1

Follow up in 4 weeks

Note:

Findings consistent with lateral epicondylitis. Recommended a counterforce brace, activity modification avoiding racquet sports, and scheduled NSAIDs. Discussed an eccentric wrist extensor strengthening program. No imaging indicated given the classic clinical presentation.

Staff:

Dorian Vance, MD (Primary Provider) (Bill Under)

Electronically Signed By: Dorian Vance, MD, 07/08/2025 09:15 AM CDT

Medications

ibuprofen Oral

Medical History

None

Musculoskeletal History

None

Musculoskeletal Family History

None

Musculoskeletal Surgery

None

Surgical History

None

Social History

Smoking status - Never smoker

Chief Complaints:

1. Follow-up right lateral elbow pain, partial improvement.

HPI: This is a 43 year old male who:

Patient returns reporting about thirty percent improvement in right elbow pain with brace use and activity modification. Still has pain with heavy gripping tasks at work. Has been compliant with home eccentric strengthening exercises.

Vitals:

Date	Taken By	B.P.	Pulse	Resp.	O2 Sat.	Temp.	Ht.	Wt.	BMI BSA
08/05/25	T. Nguyen, MA	122/76	66	14	99	98.1	72.0 in	194.0 lbs	26.3 2.04

Exam:

Comprehensive, Upper Extremity Neurovascular

Appearance: well nourished

Orientation: Alert and oriented to person, place, time.

Mood: mood and affect well-adjusted, pleasant and cooperative,

Right UE Pulses: normal radial pulse and good capillary refill

Left UE Pulses: normal radial pulse and good capillary refill

RUE Peripheral Sensation: intact to light touch throughout peripheral nerve distributions

LUE Peripheral Sensation: intact to light touch throughout peripheral nerve distributions

Elbow

Right Elbow Active ROM:

Flexion: 145 degrees.
Extension: 0 degrees.
Pronation: 80 degrees.
Supination: 80 degrees.

Left Elbow Active ROM:

Flexion: 145 degrees.
Extension: 0 degrees.
Pronation: 80 degrees.
Supination: 80 degrees.

Right Elbow Passive ROM:

Flexion: 145 degrees.
Extension: 0 degrees.
Pronation: 80 degrees.
Supination: 80 degrees.

Left Elbow Passive ROM:

Flexion: 145 degrees.
Extension: 0 degrees.
Pronation: 80 degrees.
Supination: 80 degrees.

Skin:

Right Elbow: skin intact, no rashes or lesions.

Skin:

Left Elbow: skin intact, no rashes or lesions.

Inspection:

Right Elbow: point tenderness over the lateral epicondyle, no effusion

Inspection:

Left Elbow: Normal alignment, no deformity, no tenderness, no warmth, no masses, no muscle atrophy, no crepitus

Strength:

Right Elbow Flexion: Strength: 5/5, normal muscle tone.
Right Elbow Extension: Strength: 5/5, normal muscle tone.

Strength:

Left Elbow Flexion: Strength: 5/5, normal muscle tone.
Left Elbow Extension: Strength: 5/5, normal muscle tone.

Stability:

Right Elbow: Stable

Special:

Right Elbow: Resisted wrist extension: positive, resisted middle finger extension: positive

Impression/Plan:

1. Elbow Pain, Right

Lateral epicondylitis, right elbow, improving (M77.11)

Plan: Prescription.

ibuprofen 600 mg tablet PO

Sig: Take 1 po tid PRN with food

Quantity: 30 Tablet Refills: 1

Follow up in 8 weeks

Note:

Continued improvement in lateral epicondylitis with conservative management. Adjusting ibuprofen to an as-needed dosing schedule and continuing eccentric strengthening. Will consider a corticosteroid injection if there is a plateau in improvement.

Staff:

Dorian Vance, MD (Primary Provider) (Bill Under)

Electronically Signed By: Dorian Vance, MD, 08/05/2025 09:30 AM CDT