

NCERT Maths 10.5.3 Q14

EE23BTECH11201 - ABBURI TANUSHA*

Question: Find the sum of odd numbers between 0 and 50.

Solution:

Symbol	Value	Description
$x(0)$	1	first term of AP
d	2	common difference
$x(n)$	$(1 + 2n)u(n)$	n -th term of AP

TABLE I
GIVEN PARAMETERS

$$x(n) = (1 + 2n)u(n) \quad (1)$$

Last term of the given sequence is 49.

$$(2n + 1) = 49 \quad (2)$$

$$2n = 48 \quad (3)$$

$$n = 24 \quad (4)$$

Applying Z transform:

$$x(z) = \frac{1}{1 - z^{-1}} + \frac{2z^{-1}}{(1 - z^{-1})^2} \quad (5)$$

$$= \frac{1 + z^{-1}}{(1 - z^{-1})^2} \quad (6)$$

Region of Convergence or R.O.C :

$$|z| > 1 \quad (7)$$

For AP, the sum of first $n+1$ terms can be written as :

$$y(n) = x(n) * u(n) \quad (8)$$

Applying Z transform on both sides

$$Y(z) = X(z)U(z) \quad (9)$$

$$= \frac{1}{(1 - z^{-1})^2} + \frac{2z^{-1}}{(1 - z^{-1})^3} \quad (10)$$

Using contour integration to find inverse Z transform:

$$y(n) = \frac{1}{2\pi j} \oint_C Y(z)z^{n-1} dz \quad (11)$$

$$= \frac{1}{2\pi j} \oint_C \left[\frac{1}{(1 - z^{-1})^2} + \frac{2z^{-1}}{(1 - z^{-1})^3} \right] z^{n-1} dz \quad (12)$$

The sum of the terms of the sequence is computed using the residue theorem, expressed as R_i , which represents the residue of the Z-transform at $z = 1$ for the expression $Y(z)$.

$$R_i = R_1 + R_2 \quad (13)$$

R_1 and R_2 are residues calculated at the poles of the Z-transform.

$$R_1 = \frac{1}{(2 - 1)!} \left. \frac{d(z^{25})}{dz} \right|_{z=1} \quad (14)$$

$$= 25 \quad (15)$$

$$R_2 = \frac{1}{(3 - 1)!} \left. \frac{d^2(2z^{25})}{dz^2} \right|_{z=1} \quad (16)$$

$$= 25 \times 24 \quad (17)$$

$$= 600 \quad (18)$$

$$R_i = 600 + 25 \quad (19)$$

$$(20)$$

Sum of the numbers is $R_i = 625$.

