










LARGE INTESTINE
REDUCED IN BOTH LENGTH AND
FUNCTION

OVARIES

REDUCED EGG RESERVE,
HORMONAL ADJUSTMENT TO
LOWER ESTROGEN LEVELS,
OVULATION OCCURS ONCE EVERY
2-3 MONTHS

FEMORAL ARTERIES

CIRCULATION DECREASES
SIGNIFICANTLY TO LESS, WHICH
PRIMARYLY SERVE AS PASSIVE
SUPPORT FOR STANDING OR
BALANCE

BLADDER

REDUCED CAPACITY DUE TO
REDUCED URINATION FREQUENCY

VAGINA


REDUCED ELASTICITY AND
MUSCULATURE

COLON

SLIMMER TO MATCH A SIMPLER
DIGESTIVE WORKLOAD

FEMORAL ARTERIES

CIRCULATION DECREASES
SIGNIFICANTLY TO LESS, WHICH
PRIMARYLY SERVE AS PASSIVE
SUPPORT FOR STANDING OR BALANCE




POPLITEAL ARTERIES
EXPANDED TO ACCOMMODATE
FREQUENT KNEE FLEXION AND
TRANSITIONS BETWEEN MODULAR
SPACES

BODY HAIR
ARMS, LEGS, AND TORSO HAS
DISAPPEARED ENTIRELY

TIBIA AND FIBULA
ATROPHIED SIGNIFICANTLY LOWER
LEGS ARE LESS MUSCULAR AND
RELY ON MINIMAL BONE STRENGTH
FOR BALANCE

PHALANGES
SHORTER AND STUBBER



TIBIAL ARTERIES
STRENGTHENED TO ENSURE
ROBUST BLOOD DELIVERY TO
COMPACT, WEIGHT-BEARING
LOWER LIMBS

TIBIA AND FIBULA
ATROPHIED SIGNIFICANTLY, LOWER
LEGS ARE LESS MUSCULAR AND
HAVE MINIMAL BONE STRENGTH
FOR BALANCE

PLANTAR ARTERIES
ENLARGED AND REORGANIZED TO
SUPPORT BROADER, FLATTER
FEET FOR IMPROVED STABILITY

TOUCH
SENSITIVITY REDUCED,
WALKING ON SMOOTHER,
PREDICTABLE SURFACES
ELIMINATES THE NEED FOR
DETAILED TEXTURE
DETECTION

METATARSALS
FOOT ARCH COLLAPSES SLIGHTLY