

Properties of Frank-Wolfe's Method and Linear Minimization Oracle

Tao Hu

October 1, 2025

Abstract Keywords Frank-Wolfe · Robust Algorithm · Projection · Linear Minimization Oracle

My final project mainly discusses the performance of Frank-Wolfe's method when a disturbed gradient is used. Also, I proved that an accurate linear minimization oracle is no slower than an inexact projection.

1 Introduction

We consider the maximization problems:

$$\begin{aligned} \max_{\lambda} \quad & h(\lambda) \\ \text{s.t.} \quad & \lambda \in Q, \end{aligned} \tag{1}$$

where $Q \subset E$ is convex and compact, and $h(\cdot) : Q \rightarrow \mathbb{R}$ is differentiable on Q .

In order to prove the convergence of the Frank-Wolfe algorithm, the following two auxiliary sequences are frequently used, where α_k and β_k are determined by the first k step-sizes, $\bar{\alpha}_1, \dots, \bar{\alpha}_k$ in Algorithm 1:

$$\beta_k = \frac{1}{\prod_{j=1}^{k-1} (1 - \bar{\alpha}_j)}, \quad \alpha_k = \frac{\beta_k \bar{\alpha}_k}{1 - \bar{\alpha}_k}, \quad k \geq 1. \tag{2}$$

(We follow the conventions: $\prod_{j=1}^0 \cdot = 1$ and $\sum_{i=1}^0 \cdot = 0$.)

Tao Hu
Xi'an Jiaotong University
Tel.: +1 (510)-384-9593
E-mail: tao.hu@berkeley.edu

Algorithm 1 Frank-Wolfe Method for maximizing $h(\lambda)$

Pick initial point $\lambda_0 \in Q$, initial upper bound U_{init} (usually determined by prior knowledge, if no prior knowledge is assumed, then $U_{init} = +\infty$), $k \leftarrow 0$.

At iteration k :

1. Evaluate $\nabla h(\lambda_k)$.
 2. Evaluate $\tilde{\lambda}_k \leftarrow \arg \max_{\lambda \in Q} \{h(\lambda_k) + \nabla h(\lambda_k)^T(\lambda - \lambda_k)\}$.
 $U_k^w \leftarrow h(\lambda_k) + \nabla h(\lambda_k)^T(\tilde{\lambda}_k - \lambda_k)$.
 $G_k \leftarrow \nabla h(\lambda_k)^T(\tilde{\lambda}_k - \lambda_k)$.
 $U_k \leftarrow \min\{U_{k-1}, U_k^w\}$, if $k \geq 1$.
 $U_0 \leftarrow \min\{U_{init}, U_0^w\}$, if $k = 0$.
 3. Set $\lambda_{k+1} \leftarrow \lambda_k + \bar{\alpha}_k(\tilde{\lambda}_k - \lambda_k)$, where $\bar{\alpha}_0 = 1$ and $\bar{\alpha}_k \in [0, 1], \forall k \geq 1$.
-

2 Frank-Wolfe Method for smooth convex functions

Assume that h is concave and its gradient is $L_{h,Q}$ -Lipschitz, that is,

$$\|\nabla h(\lambda) - \nabla h(\bar{\lambda})\|_* \leq L_{h,Q} \|\lambda - \bar{\lambda}\|, \forall \lambda, \bar{\lambda} \in Q.$$

We denote $C_{h,Q} = L_{h,Q}(\text{Diam}_Q)^2$.

In this scenario, several results have been discussed by Freund and Grigas [1].

Theorem 1 *If the step-size sequence $\{\bar{\alpha}_k\}$ is used in the iterate sequences of the Frank-Wolfe method (Method 1), then the following inequality holds for all $k \geq 0$,*

$$U_k - h(\lambda_{k+1}) \leq \frac{U_0 - h(\lambda_1)}{\beta_{k+1}} + \frac{\frac{1}{2}C_{h,Q} \sum_{i=1}^k \frac{\alpha_i^2}{\beta_{i+1}}}{\beta_{k+1}}. \quad (3)$$

Proof

$$\begin{aligned} h(\lambda_{k+1}) &= h(\lambda_k) + \nabla h(\lambda_k)^T(\lambda_{k+1} - \lambda_k) - \frac{1}{2}L_{h,Q}\|\lambda_{k+1} - \lambda_k\|^2 \\ &= h(\lambda_k) + \bar{\alpha}_k \nabla h(\lambda_k)^T(\tilde{\lambda}_k - \lambda_k) - \frac{1}{2}\bar{\alpha}_k^2 L_{h,Q}\|\tilde{\lambda}_k - \lambda_k\|^2 \\ &\geq h(\lambda_k) + \bar{\alpha}_k \nabla h(\lambda_k)^T(\tilde{\lambda}_k - \lambda_k) - \frac{1}{2}\bar{\alpha}_k^2 C_{h,Q}. \end{aligned}$$

We have that

$$U_k^w = h(\lambda_k) + \nabla h(\lambda_k)^T(\tilde{\lambda}_k - \lambda_k).$$

Therefore,

$$\begin{aligned} h(\lambda_{k+1}) &\geq (1 - \bar{\alpha}_k)h(\lambda_k) + \bar{\alpha}_k(h(\lambda_k) + \nabla h(\lambda_k)^T(\tilde{\lambda}_k - \lambda_k)) - \frac{1}{2}\bar{\alpha}_k^2 C_{h,Q} \\ &= (1 - \bar{\alpha}_k)h(\lambda_k) + \bar{\alpha}_k U_k^w - \frac{1}{2}\bar{\alpha}_k^2 C_{h,Q}. \end{aligned}$$

Hence,

$$\begin{aligned}
\beta_{k+1}h(\lambda_{k+1}) &\geq \beta_{k+1}(1 - \bar{\alpha}_k)h(\lambda_k) + \bar{\alpha}_k\beta_{k+1}U_k^w - \frac{1}{2}\bar{\alpha}_k^2\beta_{k+1}C_{h,Q} \\
&= \beta_k h(\lambda_k) + (\beta_{k+1} - \beta_k)U_k^w - \frac{1}{2}\frac{\alpha_k^2}{\beta_{k+1}}C_{h,Q} \\
&\geq \beta_k h(\lambda_k) + (\beta_{k+1} - \beta_k)U_k - \frac{1}{2}\frac{\alpha_k^2}{\beta_{k+1}}C_{h,Q}.
\end{aligned}$$

Therefore,

$$\begin{aligned}
\beta_{k+1}(U_k - h(\lambda_{k+1})) &\leq \beta_k(U_k - h(\lambda_k)) + \frac{1}{2}\frac{\alpha_k^2}{\beta_{k+1}}C_{h,Q} \\
&\leq \beta_k(U_{k-1} - h(\lambda_k)) + \frac{1}{2}\frac{\alpha_k^2}{\beta_{k+1}}C_{h,Q}.
\end{aligned}$$

Taking summation, we get that

$$\beta_{k+1}(U_k - h(\lambda_{k+1})) \leq \beta_1(U_0 - h(\lambda_1)) + \frac{1}{2}\sum_{j=1}^k \frac{\alpha_j^2}{\beta_{j+1}}C_{h,Q}.$$

We conclude that, since $\beta_{k+1} > 0$,

$$U_k - h(\lambda_{k+1}) \leq \frac{U_0 - h(\lambda_1)}{\beta_{k+1}} + \frac{C_{h,Q} \sum_{j=1}^k \frac{\alpha_j^2}{\beta_{j+1}}}{2\beta_{k+1}}.$$

□

2.1 Inexact Gradient

Assuming that we are using a δ -oracle, that is, every time we are doing a gradient evaluation, we don't get an exact gradient; instead, we get a "gradient" $g_\delta(\lambda)$ with error $\delta/(DiamQ)$, that is,

$$\|g_\delta(\lambda) - \nabla h(\lambda)\| \leq \frac{\delta}{DiamQ}, \forall \lambda \in Q.$$

We conclude that, $\forall \lambda_1, \lambda_2 \in Q$,

$$\begin{aligned}
h(\lambda_2) &\leq h(\lambda_1) + \nabla h(\lambda_1)^T(\lambda_2 - \lambda_1) \\
&\leq h(\lambda_1) + g_\delta(\lambda_1)^T(\lambda_2 - \lambda_1) + \|g_\delta(\lambda_1) - \nabla h(\lambda_1)\| \|\lambda_2 - \lambda_1\| \\
&\leq h(\lambda_1) + g_\delta(\lambda_1)^T(\lambda_2 - \lambda_1) + \delta;
\end{aligned}$$

$$\begin{aligned}
h(\lambda_2) &\geq h(\lambda_1) + \nabla h(\lambda_1)^T(\lambda_2 - \lambda_1) - \frac{1}{2}L_{h,Q}\|\lambda_2 - \lambda_1\|^2 \\
&\geq h(\lambda_1) + g_\delta(\lambda_1)^T(\lambda_2 - \lambda_1) - \|g_\delta(\lambda_1) - \nabla h(\lambda_1)\| \|\lambda_2 - \lambda_1\| - \frac{1}{2}L_{h,Q}\|\lambda_2 - \lambda_1\|^2 \\
&\geq h(\lambda_1) + g_\delta(\lambda_1)^T(\lambda_2 - \lambda_1) - \delta - \frac{1}{2}L_{h,Q}\|\lambda_2 - \lambda_1\|^2;
\end{aligned}$$

Algorithm 2 Frank-Wolfe Method Using δ -Oracle

Pick initial point $\lambda_0 \in Q$, initial upper bound U_{init} (usually determined by prior knowledge, if know prior knowledge is assumed, then $U_{init} = +\infty$), $k \leftarrow 0$.

At iteration k :

1. Evaluate $g_\delta(\lambda_k)$.
 2. Evaluate $\lambda_k \leftarrow \arg \max_{\lambda \in Q} \{h(\lambda_k) + g_\delta(\lambda_k)^T(\lambda - \lambda_k)\}$.
 $U_k^w \leftarrow h(\lambda_k) + \delta + g_\delta(\lambda_k)^T(\tilde{\lambda}_k - \lambda_k)$.
 $U_k \leftarrow \min\{U_{k-1}, U_k^w\}$, if $k \geq 1$.
 $U_0 \leftarrow \min\{U_{init}, U_0^w\}$, if $k = 0$.
 3. Set $\lambda_{k+1} \leftarrow \lambda_k + \bar{\alpha}_k(\tilde{\lambda}_k - \lambda_k)$, where $\bar{\alpha}_0 = 1$ and $\bar{\alpha}_k \in [0, 1], \forall k \geq 1$.
-

Under inexact gradient evaluation, the bound is given as follows [1],

Theorem 2 *If the step-size sequence $\{\bar{\alpha}_k\}$ is used in the iterate sequences of the Frank-Wolfe method with the δ -oracle (Algorithm 2),*

$$U_k - h(\lambda_{k+1}) \leq \frac{U_0 - h(\lambda_1)}{\beta_{k+1}} + \frac{\frac{1}{2}C_{h,Q} \sum_{i=1}^k \frac{\alpha_i^2}{\beta_{i+1}}}{\beta_{k+1}} + \frac{2\delta \sum_{i=1}^k \beta_{i+1}}{\beta_{k+1}}, \forall k \geq 0. \quad (4)$$

Proof

$$\begin{aligned} h(\lambda_{k+1}) &\geq h(\lambda_k) + g_\delta(\lambda_k)^T(\lambda_{k+1} - \lambda_k) - \delta - \frac{1}{2}L_{h,Q}\|\lambda_{k+1} - \lambda_k\|^2 \\ &= h(\lambda_k) + \bar{\alpha}_k g_\delta(\lambda_k)^T(\tilde{\lambda}_k - \lambda_k) - \delta - \frac{1}{2}\bar{\alpha}_k^2 L_{h,Q}\|\tilde{\lambda}_k - \lambda_k\|^2 \\ &\geq (1 - \bar{\alpha}_k)h(\lambda_k) + \bar{\alpha}_k(h(\lambda_k) + g_\delta(\lambda_k)^T(\tilde{\lambda}_k - \lambda_k) + \delta) - (1 + \bar{\alpha}_k)\delta - \frac{1}{2}C_{h,Q}\bar{\alpha}_k^2 \\ &\geq (1 - \bar{\alpha}_k)h(\lambda_k) + \bar{\alpha}_k U_k^w - 2\delta - \frac{1}{2}C_{h,Q}\bar{\alpha}_k^2 \\ &\geq (1 - \bar{\alpha}_k)h(\lambda_k) + \bar{\alpha}_k U_k - 2\delta - \frac{1}{2}C_{h,Q}\bar{\alpha}_k^2 \end{aligned}$$

Hence,

$$\begin{aligned} U_k - h(\lambda_{k+1}) &\leq U_k - (1 - \bar{\alpha}_k)h(\lambda_k) - \bar{\alpha}_k U_k + 2\delta + \frac{1}{2}C_{h,Q}\bar{\alpha}_k^2 \\ &= (1 - \bar{\alpha}_k)(U_{k-1} - h(\lambda_k)) + 2\delta + \frac{1}{2}C_{h,Q}\bar{\alpha}_k^2. \end{aligned}$$

Therefore,

$$\beta_k(U_k - h(\lambda_{k+1})) \leq \beta_{k-1}(U_{k-1} - h(\lambda_k)) + 2\delta\beta_k + \frac{1}{2}C_{h,Q}\bar{\alpha}_k^2\beta_k.$$

By taking summation, we get that

$$\beta_k(U_k - h(\lambda_{k+1})) \leq (U_0 - h(\lambda_1)) + 2\delta \sum_{j=1}^k \beta_j + \frac{1}{2}C_{h,Q} \sum_{j=1}^k \bar{\alpha}_j^2 \beta_j.$$

To conclude,

$$U_k - h(\lambda_{k+1}) \leq \frac{U_0 - h(\lambda_1)}{\beta_k} + 2\delta \frac{\sum_{j=1}^k \beta_j}{\beta_k} + \frac{1}{2} C_{h,Q} \sum_{j=1}^k \bar{\alpha}_j^2 \beta_j.$$

Dividing both side by β_k yields the result. \square

Consider the right hand side of Equation (4), we have that

$$\begin{aligned} & \frac{U_0 - h(\lambda_1)}{\beta_{k+1}} + \frac{\frac{1}{2} C_{h,Q} \sum_{i=1}^k \frac{\alpha_i^2}{\beta_{i+1}}}{\beta_{k+1}} + \frac{\sum_{i=1}^k \beta_{i+1} \delta}{\beta_{k+1}} \\ &= \frac{U_0 - h(\lambda_1)}{\beta_{k+1}} + \frac{\sum_{i=1}^k (\frac{1}{2} C_{h,Q} \frac{\alpha_i^2}{\beta_{i+1}} + \beta_{i+1} \delta)}{\beta_{k+1}} \\ &\geq \frac{U_0 - h(\lambda_1)}{\beta_{k+1}} + \frac{\sum_{i=1}^k \alpha_i \sqrt{2C_{h,Q} \delta}}{\beta_{k+1}} \\ &= \frac{U_0 - h(\lambda_1)}{\beta_{k+1}} + \sqrt{2C_{h,Q} \delta} \frac{\beta_{k+1} - 1}{\beta_{k+1}}. \end{aligned}$$

For k sufficiently large, we have that

$$0 < \frac{\beta_2 - 1}{\beta_2} \leq \frac{\beta_{k+1} - 1}{\beta_{k+1}} \leq 1.$$

Therefore, in order to achieve a first-order accurate solution, we need a second-order accurate gradient.

3 Frank-Wolfe Method with constant step sizes

We now take a special example of the results above.

3.1 Concave Functions with Lipschitz Gradient

Given $\bar{\alpha} \in (0, 1)$, consider using the following constant step-size rule:

$$\bar{\alpha}_i = \bar{\alpha} \quad \text{for } i \geq 1. \quad (5)$$

.

Proposition 1 *Under the step-size sequence (5), the following inequality holds for all $k \geq 1$:*

$$U_k - h(\lambda_{k+1}) \leq (U_k - h(\lambda_1)) (1 - \bar{\alpha})^k + \frac{1}{2} C_{h,Q} [\bar{\alpha} - \bar{\alpha}(1 - \bar{\alpha})^k]. \quad (6)$$

As a consequence:

$$U_k - h(\lambda_{k+1}) \leq \frac{1}{2} C_{h,Q} [(1 - \bar{\alpha})^{k+1} + \bar{\alpha}]. \quad (7)$$

It can be easily observed that the right hand side is a convex function. By taking derivatives and finding the stationary point, we can get that the optimal $\bar{\alpha}$ to choose is

$$\bar{\alpha} = 1 - \frac{1}{\sqrt[k]{k+1}}, \quad (8)$$

if we fix the iteration number before the algorithm starts.

The following proposition and the idea of prestart is provided by Freund and Grigas [1].

Proposition 2 $U_0 - h(\lambda_1) \leq \frac{1}{2}C_{h,Q}$.

Proof We have $\lambda_1 = \tilde{\lambda}_0$ and $U_0 \leq U_0^w$, whereby from the definition of $C_{h,Q}$ using $\alpha = 1$ we have:

$$h(\lambda_1) = h(\tilde{\lambda}_0) \geq h(\lambda_0) + \nabla h(\lambda_0)^T (\tilde{\lambda}_0 - \lambda_0) - \frac{1}{2}C_{h,Q} = U_0^w - \frac{1}{2}C_{h,Q} \geq U_0 - \frac{1}{2}C_{h,Q},$$

and the result follows by rearranging terms.

When we are using a constant step size $\bar{\alpha}_k = \bar{\alpha}$, and we take $\delta_i = \delta$, then we have that

$$\beta_k = (1 - \bar{\alpha})^{-k+1}, \alpha_k = \bar{\alpha}(1 - \bar{\alpha})^{-k}.$$

By substitute them in, we get that

$$\begin{aligned} & U_k - h(\lambda_{k+1}) \\ & \leq (1 - \bar{\alpha})^{k-1}(U_k - h(\lambda_1)) + \frac{1}{2}\text{Diam}_Q^2(1 - \bar{\alpha})^k \sum_{i=1}^k L_i \bar{\alpha}^2 (1 - \bar{\alpha})^{-i} + (1 - \bar{\alpha})^k \sum_{i=1}^k (1 - \bar{\alpha})^{-i} \delta \\ & \leq (1 - \bar{\alpha})^{k-1}(U_k - h(\lambda_1)) + \frac{1}{2}\text{Diam}_Q^2(1 - \bar{\alpha})^k \sum_{i=1}^k L \bar{\alpha}^2 (1 - \bar{\alpha})^{-i} + (1 - \bar{\alpha})^k \sum_{i=1}^k (1 - \bar{\alpha})^{-i} \delta \\ & = (1 - \bar{\alpha})^{k-1}(U_k - h(\lambda_1)) + \frac{1}{2}\text{Diam}_Q^2(1 - \bar{\alpha})^k \sum_{i=1}^k L \bar{\alpha}^2 (1 - \bar{\alpha})^{-i} + (1 - \bar{\alpha})^k \sum_{i=1}^k (1 - \bar{\alpha})^{-i} \delta \\ & = (1 - \bar{\alpha})^{k-1}(U_k - h(\lambda_1)) + \left(\frac{1}{2}\text{Diam}_Q^2 L \bar{\alpha}^2 + \delta \right) (1 - \bar{\alpha})^k \sum_{i=1}^k (1 - \bar{\alpha})^{-i} \\ & = (1 - \bar{\alpha})^{k-1}(U_k - h(\lambda_1)) + \left(\frac{1}{2}\text{Diam}_Q^2 L \bar{\alpha}^2 + \delta \right) \frac{1 - (1 - \bar{\alpha})^k}{\bar{\alpha}} \\ & = (1 - \bar{\alpha})^{k-1}(U_k - h(\lambda_1)) + \left(\frac{1}{2}\text{Diam}_Q^2 L \bar{\alpha} + \frac{\delta}{\bar{\alpha}} \right) (1 - (1 - \bar{\alpha})^k). \end{aligned}$$

By Proposition 2, we can proceed that,

$$\begin{aligned}
& U_k - h(\lambda_{k+1}) \\
& \leq \frac{1}{2}(1 - \bar{\alpha})^{k-1} C_{h,Q} + \left(\frac{1}{2} C_{h,Q} \bar{\alpha} + \frac{\delta}{\bar{\alpha}} \right) (1 - (1 - \bar{\alpha})^k) \\
& \leq \frac{1}{2} C_{h,Q} (1 - \bar{\alpha})^{k-1} + \left(\frac{1}{2} C_{h,Q} \bar{\alpha} + \frac{\delta}{\bar{\alpha}} \right) \\
& = \frac{1}{2} C_{h,Q} ((1 - \bar{\alpha})^{k-1} + \bar{\alpha}) + \frac{\delta}{\bar{\alpha}}.
\end{aligned}$$

If we take $\bar{\alpha} = 1 - (k - 1)^{-1/(k-2)}$, then

$$\bar{\alpha} = 1 - (k - 1)^{-1/(k-2)} = 1 - e^{-\frac{\log(k-1)}{k-2}} \in \left[\frac{1}{e} \frac{\log(k-1)}{k-2}, \frac{\log(k-1)}{k-2} \right]$$

$$\begin{aligned}
& U_k - h(\lambda_{k+1}) \\
& \leq \frac{1}{2} C_{h,Q} ((1 - \bar{\alpha})^{k-1} + \bar{\alpha}) + \frac{\delta}{\bar{\alpha}} \\
& \leq \frac{1}{2} C_{h,Q} ((1 - \bar{\alpha})^{k-2} + \bar{\alpha}) + \frac{\delta}{\bar{\alpha}} \\
& = \frac{1}{2} C_{h,Q} \left(\frac{1}{k-1} + 1 - (k-1)^{-\frac{1}{k-2}} \right) + \frac{\delta}{\bar{\alpha}} \\
& = \frac{1}{2} C_{h,Q} \left(\frac{1}{k-1} + 1 - e^{-\frac{\log(k-1)}{k-2}} \right) + \frac{\delta}{\bar{\alpha}} \\
& \leq \frac{1}{2} C_{h,Q} \left(\frac{1}{k-1} + \frac{\log(k-1)}{k-2} \right) + \frac{\delta}{\bar{\alpha}}
\end{aligned}$$

Therefore, in order to achieve a convergence rate of $\tilde{O}(\frac{1}{k})$, it suffices to let

$$\delta \sim O\left(\frac{\bar{\alpha}}{k}\right) = O\left(\frac{\log(k)}{k^2}\right).$$

4 A Comparison between the Projection Problem and the Linear Minimization Problem

4.1 Coarse Projection is More Expensive than High-accuracy Linear Minimization

Definition 1 Suppose that \mathcal{H} is a real Hilbert space equipped with inner product $\langle \cdot | \cdot \rangle$ which induces the norm $\|\cdot\|$. Assume that $C \subset \mathcal{H}$ is a nonempty compact convex set. From the compactness of C , we are allowed to define the projection operator and the set of linear minimization oracle points separately as follows:

$$\text{Proj}_C: \mathcal{H} \rightarrow \mathcal{H}: x \mapsto \text{Argmin}_{c \in C} \|c - x\|,$$

$$\text{LMO}_C : \mathcal{H} \rightarrow 2^{\mathcal{H}} : x \mapsto \text{Argmin}_{c \in C} \langle c \mid x \rangle.$$

In the studies of robust algorithms, the errors in both the projection and the minimization should be considered. For $\varepsilon \geq 0$, we define the set of ε -approximate projection oracle points and ε -approximate linear minimization oracle as follows:

$$\begin{aligned} \varepsilon\text{-Proj}_C : \mathcal{H} \rightarrow 2^{\mathcal{H}} : x \mapsto \{v \in \mathcal{H} \mid 0 \leq \frac{1}{2}\|v - x\|^2 - \min_{c \in C} \frac{1}{2}\|c - x\|^2 \leq \varepsilon\}, \\ \varepsilon\text{-LMO}_C : \mathcal{H} \rightarrow 2^{\mathcal{H}} : x \mapsto \{v \in \mathcal{H} \mid 0 \leq \langle v \mid x \rangle - \min_{c \in C} \langle c \mid x \rangle \leq \varepsilon\}. \end{aligned}$$

The definition above actually provides four common minimization problems. An interesting question is which one of them is the hardest and which one of them is the easiest. Zev Woodstock [2] proved that an exact projection evaluation would be no easier than finding a point in $\varepsilon\text{-LMO}_C$, no matter how small ε is. The main contribution of this chapter is to give a small modification of Woodstock's proof and prove that finding a point in $K\text{-Proj}_C$ would be no easier than finding a point in $\varepsilon\text{-LMO}_C$.

Proposition 3 *Suppose that $C \subset \mathcal{H}$ is a nonempty compact convex set and $x \in \mathcal{H}$, and that $p \in \varepsilon\text{-Proj}(x)$. Then*

$$\langle c - p \mid x - p \rangle \leq \varepsilon + \frac{1}{2}\|c - p\|^2, \forall c \in C.$$

Proof Since $p \in \varepsilon\text{-Proj}(x)$,

$$\frac{1}{2}\|p - x\|^2 \leq \frac{1}{2}\|c - x\|^2 + \varepsilon, \forall c \in C.$$

This could be written in the inner product form,

$$\frac{1}{2}\langle c - p \mid 2x - p - c \rangle \leq \varepsilon, \forall c \in C.$$

We thus get that

$$\langle c - p \mid x - p \rangle \leq \varepsilon + \frac{1}{2}\|c - p\|^2, \forall c \in C.$$

□

Theorem 3 *Let $x \in \mathcal{H}$ and $C \subset \mathcal{H}$ be nonempty, convex and compact. We denote $\delta_C := \sup_{(c_1, c_2) \in C^2} \|c_1 - c_2\| \geq 0$ and $\mu_C := \sup_{c \in C} \|c\| \geq 0$. Take any positive real number K and take $v \in \text{LMO}_C(x)$. For any $\lambda > 0$, take*

$$p' \in K\text{-Proj}_C(-\lambda x).$$

Then we have that,

$$0 \leq \langle p' \mid x \rangle - \langle v \mid x \rangle \leq \frac{K + \delta_C^2/2 + \min\{\mu_C \delta_C, \mu_C^2\}}{\lambda}.$$

In consequence, we have that for any $\varepsilon > 0$, if $\lambda \geq \frac{K + \delta_C^2/2 + \min\{\mu_C \delta_C, \mu_C^2\}}{\varepsilon}$, then

$$0 \leq \langle p' \mid x \rangle - \min_{c \in C} \langle c \mid x \rangle \leq \varepsilon.$$

We conclude that

$$K\text{-Proj}(-\lambda x) \subset \varepsilon\text{-LMO}(x).$$

Proof By proposition 3, we have that

$$\langle c - p' \mid -\lambda x - p' \rangle \leq K + \frac{1}{2} \|c - p'\|^2, \forall c \in C.$$

Then, and take $c = v$,

$$\lambda \langle p' \mid x \rangle - \lambda \langle v \mid x \rangle \leq K + \frac{1}{2} \|v - p'\|^2 + \langle p' \mid v - p' \rangle.$$

Next,

$$\begin{aligned} \lambda \langle p' - v \mid x \rangle &\leq K + \frac{1}{2} \|v - p'\|^2 + \langle p' \mid v - p' \rangle \\ &= K + \frac{1}{2} \|v - p'\|^2 + (\langle v \mid p' \rangle - \langle p' \mid p' \rangle) \\ &\leq K + \frac{1}{2} \|v - p'\|^2 + \|p'\|(\|v\| - \|p'\|) \\ &\leq K + \frac{1}{2} \|v - p'\|^2 + \|p'\| \|v - p'\|. \end{aligned}$$

Therefore,

$$\lambda \langle p' - v \mid x \rangle \leq K + \frac{1}{2} \delta_C^2 + \min\{\mu_C \delta_C, \mu_C^2\}.$$

Hence,

$$0 \leq \langle p' \mid x \rangle - \langle v \mid x \rangle \leq \frac{K + \frac{1}{2} \delta_C^2 + \min\{\mu_C \delta_C, \mu_C^2\}}{\lambda},$$

where the first inequality is from the fact that $v \in \text{LMO}_C(x)$. \square

4.2 Comparison Theory for exact problem solvers

Definition 2 Assume that $C \subset \mathcal{H}$ is a nonempty compact convex set, $0 \in C$, and let

$$f_C : \mathcal{H} \rightarrow 2^C, \quad g_C : \mathcal{H} \rightarrow 2^C.$$

Suppose that $I_C : \mathcal{H} \rightarrow \{0, +\infty\}$ is the indicator function of C , that is

$$I_C(x) = \begin{cases} 0 & \text{if } x \in C \\ \infty & \text{if } x \notin C \end{cases}$$

We say that evaluating f_c is not significantly harder than evaluating g_C , denoted as $f_C \preceq g_C$, if there is a function $F : 2^{\mathcal{H}} \rightarrow 2^{\mathcal{H}}$, which does not depend on C , and a sequence of points $\{x_n : \mathcal{H} \rightarrow \mathcal{H}\}_{n=1}^{+\infty}$, such that $\forall x \in \mathcal{H}$, $\exists N(x) \in \mathbb{Z}_{\geq 1}$, such that

$$f_C(x) \supset F(g_C(x_n(x))), \forall n \geq N(x).$$

Under this notation we can write proposition 3 of [2] in the following precise way,

Proposition 4 *Take $\mathcal{H} = \mathbb{R}^n$, then for any $C \in \mathcal{P} := \{\{x \in \mathcal{H} | Ax \leq b\} | A \in \mathbb{R}^{n \times m}, b \in \mathbb{R}^m\}$, we have that $\text{LMO}_C \preceq \text{Proj}_C$.*

References

1. R. M. FREUND AND P. GRIGAS, *New analysis and results for the frank-wolfe method*, Mathematical Programming, 155 (2016), p. 199–230.
2. Z. WOODSTOCK, *High-precision linear minimization is no slower than projection*, (2025). arXiv:2501.18454 [math].