

Lecture 3: Analysis of Variance

Lecturer: Prof. Jingyi Jessica Li

Subscribers: Dong Wang

Disclaimer: These notes have not been subjected to the usual scrutiny reserved for formal publications. They may be distributed outside this class only with the permission of the Instructor.

3.1 Recap

3.1.1 Testing Beta ($\hat{\beta}$): t-test

Model: $Y_{(n \times 1)} = X_{(n \times p)}\beta_{(p \times 1)} + \epsilon_{(n \times 1)}$ and $\beta \sim \begin{pmatrix} \beta_1 \\ \beta_2 \\ \dots \\ \beta_p \end{pmatrix}$

1. Null Hypothesis:

$$H_0 : \beta_j = 0; \quad H_A : \beta_j \neq 0$$

2. The t-statistics:

$$t = \frac{\hat{\beta}_j}{\sqrt{\widehat{Var}(\hat{\beta}_j)}} \sim T - distribution$$

where:

$$\widehat{Var}(\hat{\beta}) = \hat{\sigma}^2 (X^T X)^{-1}$$

3. Estimator for σ^2 :

$$\widehat{\sigma_{MLE}^2} = \frac{1}{n} (y - X\hat{\beta})^T (y - X\hat{\beta}) = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

Since RSS(Residual Sum of Squares) equals to

$$\sum_{i=1}^n (y_i - \hat{y}_i)^2 = \sum_{i=1}^n (y_i - x_i^T \hat{\beta})^2 = \text{SSE (Sum of Squared Errors)}$$

$\widehat{\sigma_{MLE}^2}$ can be expressed as

$$\widehat{\sigma_{MLE}^2} = \frac{\text{RSS}}{n} \text{ (or } \frac{\text{SSE}}{n} \text{)}$$

As $\frac{\text{RSS}}{\sigma^2} \sim \chi^2_{(n-p)}$, we know that $E[\frac{\text{RSS}}{\sigma^2}] = n - p$, so $E[\widehat{\sigma_{MLE}^2}] = \frac{n-p}{n} \sigma^2$. To make it unbiased, we can use $\hat{\sigma}_{unbiased}^2 = \frac{\text{RSS}}{n-p}$. Under the null hypothesis(H_0), we can calculate a t-statistic which will follow a t-distribution with $(n - p)$ degree of freedom:

$$t = \frac{\hat{\beta}_j}{\sqrt{\widehat{Var}(\hat{\beta}_j)}} = \frac{\hat{\beta}_j}{\sqrt{\hat{\sigma}_{unbiased}^2 \times j^{th} \text{ diagonal element of } (X^T X)^{-1}}}$$

and this t-distribution converges to normal distribution as $n \rightarrow \infty$:

$$t_{n-p} \xrightarrow{n \rightarrow \infty} N(0, 1)$$

The t-distribution is heavy-tailed relative to the normal distribution, and therefore, it is a more conservative test than the z test.

3.1.2 Wald Test

Assume: $\beta = \begin{pmatrix} \beta_1 \\ (p_1 \times 1) \\ \beta_2 \\ (p_2 \times 1) \end{pmatrix} \quad X = \begin{bmatrix} X_1 & X_2 \\ (p_1 \times 1) & (p_2 \times 1) \end{bmatrix}$

1. Null Hypothesis:

$$H_0 : \beta_2 = 0 \quad H_a : \beta_2 \neq 0$$

2. Wald test statistic:

$$W = \hat{\beta}_2^T \widehat{\text{var}(\hat{\beta}_2)}^{-1} \hat{\beta}_2, \quad \text{if } p_2 = 1, W = t^2$$

Under H_0 : $W/p_2 \sim F(p_2, n - p)$ (exact, under the normality assumption) or $W \overset{\text{approx}}{\sim} \chi_{p_2}^2$ (when n is large)

3. the smaller model: $Y = X_1\beta_1 + \epsilon$, which has p_1 predictors
the larger model: $Y = X\beta + \epsilon$, which has p predictors
4. In practice, we use the larger model to calculate $\hat{\sigma}^2 = \frac{1}{n-p} \text{RSS}_{\text{large}}$, and $\widehat{\text{var}(\hat{\beta}_2)} = \hat{\sigma}^2 \times$ the lower block diagonal $p_2 \times p_2$ matrix of $(X^T X)^{-1}$

3.1.3 Likelihood Ratio Test

1. Null Hypothesis:

$$H_0 : \beta = 0 \quad H_a : \beta \neq 0$$

2. Likelihood Ratio Statistic:

$$\begin{aligned} \Lambda &= \frac{\max L(\beta)_{H_0}}{\max L(\beta)} \\ \log \Lambda &= \max l(\beta)_{H_0} - \max l(\beta) \\ \Rightarrow \log \Lambda &= -\frac{1}{2\sigma^2} [\text{RSS}_{H_0} - \text{RSS}] \\ -2 \log \Lambda &= \frac{\text{RSS}_{H_0} - \text{RSS}}{\sigma^2} \end{aligned}$$

plug in the $\hat{\sigma}_{unbiased}^2$, we have that: $-2 \log \hat{\Lambda} = \frac{\text{RSS}_{H_0} - \text{RSS}}{\text{RSS}/(n-p)} \overset{\text{approx}}{\sim} \chi_{p_2}^2$

3.2 ANOVA

3.2.1 Extreme Linear Models

1. Null Model: (no predictors, just intercept)

$$Y_i = \mu + \epsilon_i$$

If $\epsilon_i \stackrel{\text{iid}}{\sim} N(0, \sigma^2)$, then $Y_i \stackrel{\text{iid}}{\sim} N(\mu, \sigma^2)$

2. Saturate Model:

$$Y_i = \mu_i + \epsilon_i \quad i = 1, 2, \dots, n$$

where $\mu = (\mu_1, \dots, \mu_n)^T$ are the parameters and $\hat{\mu}_i = Y_i$

3.2.2 Basic ANOVA

Model: $Y = \beta_0 + \beta_1 X_1 + \epsilon$

$$\sum_{i=1}^n (Y_i - \bar{Y})^2 = \sum_{i=1}^n (Y_i - \hat{Y}_i)^2 + \sum_{i=1}^n (\hat{Y}_i - \bar{Y})^2$$

$$SST = SSE + SSR$$

Source of Variation (SV)	Sum of Squares (SS)	Degrees of Freedom (DF)	Mean Square (MS)	F Ratios
Regression(Model)	$SSR = \sum_{i=1}^n (\hat{Y}_i - \bar{Y})^2$	$p - 1$	$MSR = \frac{SSR}{p - 1}$	$F = \frac{MSR}{MSE}$ $\sim F(p - 1, n - p)$
Residuals	$SSE = \sum_{i=1}^n (Y_i - \hat{Y}_i)^2$	$n - p$	$MSE = \frac{SSE}{n - p}$	$H_0: \beta_1 = 0$
Total	$SST = SSR + SSE$ $= \sum_{i=1}^n (Y_i - \bar{Y})^2$	$n - 1$		

Note: $MSR \sim \chi_{p-1}^2$ as $n \rightarrow \infty$, $MSE \sim \chi_{n-p}^2$, so $F \sim F_{p-1, n-p}$, so we can do hypothesis testing using F distribution.

3.2.3 Hierarchical ANOVA

Original Model: $Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \epsilon$

Small Model: $Y = \beta_0 + \beta_1 X_1 + \epsilon$

SV	SS	DF	MS	F
X_1	SSR_1	1	$MSR_1 = \frac{SSR_1}{1}$	
$X_2 X_1$	$SSR_{12} - SSR_1 = SSE_1 - SSE_{12}$	1	$MSR_{2 1} = \frac{SSR_{2 1}}{1}$	$\frac{MSR_{2 1}}{MSE} \sim F(1, n-3)$
Residual	SSE_{12}	$n-3$	$MSE_{12} = \frac{SSE_{12}}{(n-3)}$	$H_0: \beta_2 = 0$
Total		$n-1$		

Remark: In $Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \epsilon$, it's possible that $H_0: \beta_1 = \beta_2 = 0$ (ANOVA F test) is rejected at $\alpha = 0.05$, but neither $H_0: \beta_1 = 0$ nor $H_0: \beta_2 = 0$ (t test) is rejected.

	Estimate	Std. error	t
In R, <code>summary(lm(Y ~ X₁ + X₂))</code> shows the following:	$\hat{\beta}_1$	$\sqrt{\widehat{Var}(\hat{\beta}_1)}$	$t_1 = \frac{\hat{\beta}_1}{\sqrt{\widehat{Var}(\hat{\beta}_1)}} \sim t_{n-3}$
	$\hat{\beta}_2$	$\sqrt{\widehat{Var}(\hat{\beta}_2)}$	$t_2 = \frac{\hat{\beta}_2}{\sqrt{\widehat{Var}(\hat{\beta}_2)}} \sim t_{n-3}$

Note:

1. $\frac{MSR_{X_2|X_1}}{MSE} \sim F(1, n-3) \iff \hat{\beta}_2 \sim t_{n-3}$.
2. If using `summary(lm(Y ~ X2 + X1))`, then $\frac{MSR_{X_1|X_2}}{MSE} \sim F(1, n-3) \iff \hat{\beta}_1 \sim t_{n-3}$.

3.3 Coefficient of determination: R^2

1. In a simple linear model $Y = \beta_0 + \beta_1 X_1 + \epsilon$, the coefficient of determination $R^2 = 1 - \frac{SSE}{SST} = \frac{SSR}{SST} = \frac{\sum_{i=1}^n (\hat{Y}_i - \bar{\hat{Y}})^2}{\sum_{i=1}^n (Y_i - \bar{Y})^2} = r^2$, where

$$r = \frac{\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y})}{\sqrt{\sum_{i=1}^n (X_i - \bar{X})^2 \sum_{i=1}^n (Y_i - \bar{Y})^2}} = \text{cor}^2(Y, X)$$

is the Pearson correlation coefficient and is symmetric between X_1 and Y .

2. In a multiple linear model $Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \epsilon$, the coefficient of determination $R^2 = 1 - \frac{SSE}{SST}$.

3.3.1 Partial correlation

For the linear model $Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \epsilon$ with two variables, partial correlation between Y and X_2 conditional on X_1 ($r_{YX_2|X_1}$) can be calculated as follows:

1. Regress Y onto X_1 to obtain residuals $Y \sim X_1 \xrightarrow{\text{linear regression}} e_{Y|X_1}$
2. Regress X_2 onto X_1 to obtain residuals $X_2 \sim X_1 \xrightarrow{\text{linear regression}} e_{X_2|X_1}$
3. Compute Pearson correlation of the residuals $r_{YX_2|X_1} = \text{Cor}(e_{Y|X_1}, e_{X_2|X_1})$




Figure 3.1: $\cos \psi$ gives the partial correlation between Y and X_2 conditional on X_1 .

Partial correlation can be interpreted as the cosine of the angle between the projection of Y on to the plane (S_{X_1}) orthogonal to X_1 and the projection of X_2 on to S_{X_1} . More specifically, let $e_{Y|X_1} = \text{proj}_{S_{X_1}} Y$, $e_{X_2|X_1} = \text{proj}_{S_{X_1}} X_2$, and $\psi = \angle(e_{Y|X_1}, e_{X_2|X_1})$, then $r_{YX_2|X_1} = \cos \psi$.

The sample partial correlation can be calculated as

$$r_{YX_2|X_1} = \frac{r_{YX_2} - r_{YX_1}r_{X_1X_2}}{\sqrt{(1 - r_{YX_1}^2)(1 - r_{X_1X_2}^2)}}.$$

As a recap, for the linear model $Y = \beta_0 + \beta X + \epsilon$, the estimate of β is $\hat{\beta} = r_{XY} \frac{\text{sd}(Y)}{\text{sd}(X)}$. The estimate of β_0 is $\hat{\beta}_0 = \bar{Y} - \hat{\beta}\bar{X}$. And the residual is $e_{Y|X} = Y - \hat{\beta}_0 - \hat{\beta}X$.

Partial correlation is widely used in network analysis. In a network with n nodes $X = \{X_1, \dots, X_n\}$, the partial correlation between two nodes conditional on the rest of the nodes, $r_{X_i X_j | X \setminus \{X_i, X_j\}}$, can be used to detect the association between two nodes.