## Homework 4

## MATH 541: Abstract Algebra 1 Spring 2023

## Hongtao Zhang

Sec. 3.1: 2, 14, 22, 24, 36, 40, 41

 $\mathbf{2}$ 

Let  $\varphi:G\to H$  be a homomorphism of groups with kernel K and let  $a,b\in\varphi(G)$ . Let  $X\in G/K$  be the fiber above a and let Y be the fiber above b, i.e.,  $X=\varphi^{-1}(a),Y=\varphi^{-1}(b)$ . Fix an element u of X (so  $\varphi(u)=a$ ). Prove that if XY=Z in the quotient group G/K and w is any member of Z, then there is some  $v\in Y$  such that uv=w. [Show  $u^{-1}w\in Y$ ]

Proof.

$$Z = XY = \phi^{-1}(a)\phi^{-1}(b) = \phi^{-1}(ab)$$

We know that  $\phi$  is a group homomorphism, so  $\phi(xy) = \phi(x)\phi(y)$ . Therefore,  $\phi(u^{-1}w) = \phi(u^{-1})\phi(w) = a^{-1}ab = b$ , which implies that  $u^{-1}w \in Y$ .

- 14: Consider the additive quotient group  $\mathbb{Q}/\mathbb{Z}$ 
  - a Show that every coset of  $\mathbb{Z}$  in  $\mathbb{Q}$  contains exactly one representative  $q \in \mathbb{Q}$  in the range  $0 \leqslant q < 1$ .
  - b Show that every element of  $\mathbb{Q}/\mathbb{Z}$  has finite order but that there are elements of arbitrarily large order.
  - c Show that  $\mathbb{Q}/\mathbb{Z}$  is the torsion subgroup of  $\mathbb{R}/\mathbb{Z}$  (cf. Exercise 6, Section 2.1).
  - d Show that  $\mathbb{Q}/\mathbb{Z}$  is isomorphic to the multiplicative group of root of unity in  $\mathbb{C}^{\times}$ .

22

Proof. a

$$\forall q' \in \mathbb{Q} : q'\mathbb{Z} = \{q' + z : z \in \mathbb{Z}\}, \exists z' \in \mathbb{Z} : q' + z' \in [0, 1), q = q' + z' \implies q\mathbb{Z} = (q' + z')\mathbb{Z}$$

It is easy to see that  $(q' + z')\mathbb{Z} = (q')\mathbb{Z}$ , which means  $q\mathbb{Z} = q'\mathbb{Z}$ 

b Note  $\mathbb{Q}/\mathbb{Z}$  is an equivalence classes based on all  $q \in [0,1)$ .

$$\forall q \in \mathbb{Q}/\mathbb{Z}, \exists z_1, z_2 \in \mathbb{Z} : q = \frac{z_1}{z_2}$$

Therefore, the order is  $lcm(z_1, z_2)$ , which can be arbitrarily large but finite.

c It suffices to show  $\mathbb{Q}/\mathbb{Z}$  is a subgroup  $\mathbb{R}/\mathbb{Z}$  with b. We know that the sum of two rational number is a rational number. We also know that the inverse of a rational number is -q which is equivalent to 1-q. 0 is clearly a rational number.

d We can just write out the isomorphism.

For all order  $z \in \mathbb{Z}$ , the root of unity contains exactly z elements such that z is the order of the element.

Therefore, we can just map all element from  $\frac{\mathbb{Q}}{\mathbb{Z}}$  to the root of unity with the denominator as the order, and numerator as the index of the element, which is bijective, and vice versa.

24:

- a Prove that if H and K are normal subgroups of a group G then their intersection  $H \cap K$  is also a normal subgroup of G.
- b Prove that the intersection of an arbitrary non-empty collection of normal subgroups of a group is a normal subgroup (do not assume the collection is countable).

*Proof.* a Denote  $H \cap K = L$ 

$$\forall l_1, l_2 \in L : l_1 l_2 \in H, l_1 l_2 \in K \implies l_1 l_2 \in L$$

$$\forall l \in L : l^{-1} \in H, l^{-1} \in K \implies l^{-1} \in L$$

$$\forall l \in L : glg^{-1} \in H, glg^{-1} \in K \implies glg^{-1} \in L$$

b There're no difference between the two argument, but just suggesting the element is in all normal subgroups instead of H, K.

**36:** Prove that if  $N \triangleleft G$  and H is any subgroup of G then  $N \cap H \triangleleft H$ .

- **40:** Prove that if G/Z(G) is cyclic then G is abelian. [If G/Z(G) is cyclic with generator xZ(G), show that every element of G can be written in the form  $x^az$  for some integer  $a \in \mathbb{Z}$  and some element  $z \in Z(G)$ .]
- **41:** Let G be a group, let N be a normal subgroup of G and let  $\overline{G}=G/N$ . Prove that  $\overline{x}$  and  $\overline{y}$  commute in  $\overline{G}$  if and only if  $x^{-1}y^{-1}xy \in N$ . (The element  $x^{-1}y^{-1}xy$  is called the commutator of x and y and is denoted by [x,y].)
- : Let G be a group. Prove that  $N=\langle x^{-1}y^{-1}xy|x,y\in G\rangle$  is a normal subgroup of G and G/N is abelian (N is called the commutator subgroup of G)