

Math 542 HW7

Hongtao Zhang

1 Factorization of Cyclotomic Polynomials

Let l be a prime and let $\Phi_l(x) = \frac{x^l - 1}{x - 1} = x^{l-1} + x^{l-2} \dots + x + 1 \in \mathbb{Z}[x]$ be the l^{th} cyclotomic polynomial, which is irreducible in $\mathbb{Z}[x]$. This exercise determines the factorization of $\Phi_{l(x)}$ modulo p for any prime p . Let ζ denote any fixed primitive l^{th} root of unity.

1.1

Show that $p = l \Rightarrow \Phi_l(x) = (x - 1)^{l-1} \in \mathbb{F}_{l[x]}$

Solution 1.1.1

$$(x - 1)^{l-1} = \sum_{i=0}^{l-1} \binom{l-1}{i} x^i (-1)^{l-1-i}$$

Consider each binomial coefficient $\binom{l-1}{i}$ modulo l . Since l is prime, $(l-1)! \equiv -1 \pmod{l}$.

$$\binom{l-1}{i} = \frac{(l-1)!}{(l-1-i)!i!}$$

$$\Leftrightarrow \binom{l-1}{i} (l-1-i)!i! \equiv (l-1)! \equiv -1 \pmod{l} \quad (\text{Wilson Theorem})$$

$$\Leftrightarrow \binom{l-1}{i} \equiv -\frac{1}{(l-1-i)!i!} \pmod{l}$$

1.2

Suppose $p \neq l$ and let f denote the order of $p \pmod{l}$, i.e. f is the smallest power of p with $p^f \equiv 1 \pmod{l}$. Use the fact that $\mathbb{F}_{p^n}^\times$ is a cyclic group to show that $n = f$ is the smallest power p^n of p with $\zeta \in \mathbb{F}_{p^n}$. Conclude that the minimal polynomial of ζ over \mathbb{F}_p has degree f .

Solution 1.2.1

Since $\mathbb{F}_{p^n}^\times$ is a cyclic group, and ζ is a l -th primitive root of unity, for ζ to be in \mathbb{F}_{p^n} , we must have some element that has order l . Therefore $n = f$ is the smallest power of p^n of p with $\zeta \in \mathbb{F}_{p^n}$ by construction.

Solution 1.2.2

Because we have the minimum extension of ζ to be in \mathbb{F}_p^n , which is a degree n extension, the minimal polynomial of ζ over \mathbb{F}_p has degree $n = f$.

1.3

Show that $\mathbb{F}_p(\zeta) = \mathbb{F}_p(\zeta^a)$ for any integer a not divisible by l . [Hint:]

Solution 1.3.1

One direction, it suffices to check that ζ^a can be generated by ζ , which is obvious.

The other direction suffices to check that ζ can be generated by ζ^a , which follows from the hint that $\zeta = (\zeta^a)^b$ where b is the multiplicative inverse of $a \bmod l$.

Conclude using (Section 1.2) that, in $\mathbb{F}_p[x]$, $\Phi_l(x)$ is the product of $\frac{l-1}{f}$ distinct irreducible polynomials of degree f .

Solution 1.3.2

Since all primitive roots of unity have f -degree minimal polynomial, and all other roots of unity are generated by primitive roots of unity, we have that $\Phi_{l(x)}$ is the product of $\frac{l-1}{f}$ distinct irreducible polynomials of degree f .

1.4

In particular, prove that, viewed in $\mathbb{F}_p[x]$, $\Phi_7(x) = x^6 + x^5 + \dots + x + 1$ is $(x-1)^6$ for $p = 7$, a product of distinct linear factor for $p \equiv 1 \bmod 7$, a product of 3 irreducible quadratics for $p \equiv 6 \bmod 7$, a product of 2 irreducible cubics for $p \equiv 2, 4 \bmod 7$, and is irreducible for $p \equiv 3, 5 \bmod 7$.

Solution 1.4.1

2

2.1

Let φ denote the Frobenius map $x \mapsto x^p$ on the finite field \mathbb{F}_p^n as in the previous exercise. Determine the rational canonical form over \mathbb{F}_p for φ considered as an \mathbb{F}_p -linear transformation of the n -dimensional \mathbb{F}_p -vector space \mathbb{F}_p^n .

Solution 2.1.1

2.2

Let φ denote the Frobenius map $x \mapsto x^p$ on the finite field \mathbb{F}_p^n as in the previous exercise. Determine the Jordan canonical form (over a field containing all the eigenvalues) for φ considered as an \mathbb{F}_p -linear transformation of the n -dimensional \mathbb{F}_p -vector space \mathbb{F}_p^n .

3 Wedderburn's Theorem on Finite Division Rings

The exercise outline a proof of Wedderburn's Theorem that a finite division ring D is a field.

3.1

Let Z denote the center of D . Prove that Z is a field containing \mathbb{F}_p for some prime p . If $Z = \mathbb{F}_q$ prove that D has order q^n for some integer n .

Solution 3.1.1

Because we know that the center of D is finite and commutative, and thus is a finite field. Further, we know that any finite field containing some \mathbb{F}_p for some prime p .

We can view D as additive group, and thus Z is the center of D , so the order of D divides the order of Z .

3.2

The nonzero elements D^\times of D form a multiplicative group. For any $x \in D^\times$ shows that the elements of D which commute with x form a division ring which contains Z . Show that this division ring is of order q^m for some integer m and that $m < n$ if x is not an element of Z .

Solution 3.2.1

Since Z is the center, so all elements of D^\times commutex with Z and thus form a division ring contains Z .

3.3

Show that the class equation for the group D^\times is

$$q^n - 1 = (q - 1) + \sum_{i=1}^r \frac{q^n - 1}{|C_D^\times(x_i)|}$$

where x_i are representatives of the distinct conjugacy classes in D^\times not contained in the center of D^\times . Conclude that for each i , $|C_D^\times(x_i)| = q^{m_i} - 1$ for some $m_i < n$.

3.4

Prove that since $\frac{q^n-1}{q^m-1} = |D^\times : C_D^\times(x_i)|$ is an integer then m_i divides n . Conclude that $\Phi_n(x)$ divides $\frac{x^n-1}{x^{m_i}-1}$ and hence that the integer $\Phi_n(q)$ divides $\frac{q^n-1}{q^{m_i}-1}$ for $i = 1, 2, \dots, r$.

3.5

Prove that $\Phi_n(q) = \prod_{\zeta \text{ primitive}} (q - \zeta)$ divides $q - 1$. Prove that $|q - \zeta| > q - 1$ (complex absolute value) for any root of unity $\zeta \neq 1$. [note that 1 is the closest point on the unit circle in \mathbb{C} to the point q on the real line]

Conclude that $n = 1 \Leftrightarrow D = \mathbb{Z}$.

4 Dirichlet's Theorem

4.1

Given any monic polynomial $P(x) \in \mathbb{Z}[x]$ of degree at least one show that there are infinitely many distinct prime divisors of the integers

$$P(1), P(2), P(3), \dots, P(n), \dots$$

[Suppose p_1, p_2, \dots, p_k are the only primes dividing the values $P(n)$, $n = 1, 2, \dots$. Let N be an integer with $P(N) = a \neq 0$. Show that $Q(x) = a^{-1}P(N + ap_1p_2 \dots p_k x)$ is an element of $\mathbb{Z}[x]$ and that $Q(n) \equiv 1 \pmod{p_1p_2 \dots p_k}$ for $n = 1, 2, \dots$. Conclude that there is some integer M such that $Q(M)$ has a prime factor different from p_1, p_2, \dots, p_k and hence that $P(N + ap_1p_2 \dots p_k M)$ has a prime factor different from p_1, p_2, \dots, p_k .]

Suppose p_1, p_2, \dots, p_k are the only primes the dividing values $P(n)$.

Consider a integer N such that $P(N) = a \neq 0$. Consider the polynomial $Q(x) = a^{-1}P(N + ap_1p_2 \dots p_k x)$.

Lemma 4.1.1

$$Q(x) \in \mathbb{Z}[x]$$

Proof: Since P is a polynomial, we can write $P = b_1x^n + b_2x^{n-1} + \dots b_{n+1}$. Then consider $P(N + ap_1p_2 \dots p_k x)$, by binomial theorem we have each terms being written as some product of N and $ap_1p_2 \dots p_k x$. Any term involving the second part is certainly divisible by a , and the grouping of term that only contains N is equal to $P(N)$, and by assumption, is divisible by a since $P(N) = a$. Therefore $Q(x) \in \mathbb{Z}[x]$. \square

Lemma 4.1.2

$$Q(n) \equiv 1$$

Proof: We can show the following by a similar construction as above:

$$Q(n) = \frac{P(N + nap_1p_2\cdots p_k)}{a} \equiv \frac{P(N)}{a} \equiv 1 \pmod{p_1p_2\cdots p_k}$$

□

4.2

Let p be an odd prime not dividing m and let $\Phi_m(x)$ be the m^{th} cyclotomic polynomial. Suppose $a \in \mathbb{Z}$ satisfies $\Phi_m(a) \equiv 0 \pmod{p}$. Prove that a is relatively prime to p and that the order of a in $(\mathbb{Z}/p\mathbb{Z})^\times$ is precisely m . [Since

$$x^m - 1 = \prod_{d|m} \Phi_d(x) = \Phi_m(x) \prod_{\substack{d|m \\ d < m}} \Phi_d(x)$$

we see first that $a^m - 1 \equiv 0 \pmod{p}$ i.e., $a^m \equiv 1 \pmod{p}$. If the order of $a \pmod{p}$ were less than m , then $a^d \equiv 1 \pmod{p}$ for some d dividing m , so then $\Phi_d(a) \equiv 0 \pmod{p}$ for some $d < m$. But then $x^m - 1$ would have a as a multiple root mod p , a contradiction.]

4.3

Let $a \in \mathbb{Z}$. Show that if p is an odd prime dividing $\Phi_m(a)$ then either p divides m or $p \equiv 1 \pmod{m}$.

4.4

Prove there are infinitely many primes p with $p \equiv 1 \pmod{m}$.