DGMS Examination 2019 — Question 47 Detailed Solution

Junior Scientific Officer (Geology) | Marine Geology, Geodynamics & Plate Tectonics

QUESTION 47

The mean age of the oceanic floor is approximately:

(a) 20 Ma	**(b) 60 Ma** CORRECT ANSWER
(c) 80 Ma	(d) 230 Ma

1. Executive Summary & Answer Verification

The correct option is **(b) 60 Ma** (approximately 60–65 million years).

The Earth's oceanic floor is dynamically recycled through seafloor spreading at mid-ocean ridges and subduction at convergent margins. Consequently, oceanic lithosphere has a remarkably youthful age profile compared to continental crust:

- **Maximum in situ Age:** The oldest existing in situ oceanic crust is only ~180–200 Ma (Early Jurassic, located in the western Pacific Pigafetta basin and the northwest Atlantic margins).
- **Area-Weighted Mean Age:** Global digital seafloor age compilations (e.g., Müller et al., 2008; Sclater et al., 1980; Cogné & Humler, 2004) establish that the global area-weighted mean age of the ocean basins is **64.2 Ma**, matching the standard textbook value of **~60 Ma**.
- **Theoretical Derivation:** Because seafloor is continually subducted, its age-area distribution is roughly triangular between 0 Ma (ridge crest) and $T_{\max} \approx 180\text{ Ma}$. For a triangular distribution, the centroid / mean age is $\bar{T} = \frac{1}{3} T_{\max} = \frac{180}{3} = \mathbf{60\text{ Ma}}$.




FIGURE 16.26 Morphology of a midocean ridge axis. (a) Simplified profiles of a fast ridge versus a slow ridge. Notice that fast ridges are wider. That's because the depth to which the seafloor sinks depends on its age. The slow plate at point *A* is the same age as the fast plate at point *B*, but both are at the same depth. (b) Profiles contrasting fast and slow ridge axes. The top profile is across the East Pacific Rise (3° S latitude), while the lower profile is across the Mid-Atlantic Ridge (37° N latitude). Note that an axial trough only occurs at slow ridges, not at fast ridges. "V" marks the position of volcanic vents. (c) Three-dimensional block

Figure 47.1: *(e) Thermal subsidence and oceanic floor age-depth curve: Ocean depth increases systematically with crustal age as $D = 2500 + 350\sqrt{T}$ (where T is age in m.y.). Note the standard age timeline from 0 to 80+ Ma, illustrating that typical oceanic floor is young (~60 Ma average). (Van der Pluijm & Marshak, Earth Structure: An Introduction to Structural Geology and Tectonics, 2nd ed., Figure 16.26, p. 404).*

2. Geodynamic Derivation: Why is the Oceanic Floor So Young?

2.1 The Oceanic "Conveyor Belt" and Subduction Recycling

Unlike continental crust—which has an average age of ~2.0 to 2.5 Ga and contains rocks exceeding 4.0 Ga (e.g., the Acasta Gneiss)—oceanic crust rarely survives past 180–200 Ma. The underlying physical reasons are:

1. **High Density & Negative Buoyancy:** As oceanic lithosphere moves away from mid-ocean ridges, it conducts heat upward into seawater, cools, and thickens by underplating lithospheric mantle. Because cold lithospheric mantle ($\rho \approx 3.30\text{ g/cm}^3$) is denser than the underlying asthenosphere ($\rho \approx 3.20\text{ g/cm}^3$), oceanic lithosphere develops *negative buoyancy* once it exceeds ~20–30 Ma, driving slab pull and subduction.
2. **Continuous Creation & Destruction:** New oceanic crust forms at mid-ocean ridges at a global rate of ~3.0 to $3.4\text{ km}^2/\text{yr}$ (generating ~20 km^3/yr of new basaltic crust). An equal area must be subducted into the deep mantle to conserve global surface area.

2.2 Mathematical Age-Area Distribution

If we model the global ocean basin area $A(t)$ as a function of age t:

At the ridge crest ($t = 0$), new crust is continually generated. As crust ages toward $T_{\max} \approx 180\text{ Ma}$, older slabs are progressively consumed by subduction. If the probability of subduction per unit time is roughly uniform, the area of seafloor of age t decreases approximately linearly:

$$\frac{dA}{dt} = C \cdot \left(1 - \frac{t}{T_{\max}}\right) \quad \text{for } 0 \le t \le T_{\max}$$

The area-weighted mean age $\bar{t}$ is obtained by integrating:

$$\bar{t} = \frac{\int_0^{T_{\max}} t \cdot \left(1 - \frac{t}{T_{\max}}\right) dt}{\int_0^{T_{\max}} \left(1 - \frac{t}{T_{\max}}\right) dt} = \frac{\left[\frac{t^2}{2} - \frac{t^3}{3T_{\max}}\right]_0^{T_{\max}}}{\left[t - \frac{t^2}{2T_{\max}}\right]_0^{T_{\max}}} = \frac{\frac{T_{\max}^2}{6}}{\frac{T_{\max}}{2}} = \frac{T_{\max}}{3}$$

Substituting $T_{\max} \approx 180\text{ Ma}$:

$$\bar{t} = \frac{180\text{ Ma}}{3} = \mathbf{60\text{ Ma}}$$

3. Quantitative Ocean Basin Age Statistics (Global Data)

Ocean Basin	Maximum In Situ Age	Mean Crustal Age	Dominant Spreading Rates & Characteristics
Pacific Ocean	~170–180 Ma (Pigafetta Basin)	**~50–55 Ma**	Fast spreading (East Pacific Rise: 80–150 mm/yr); rapid recycling
Atlantic Ocean	~180–190 Ma (Blake Bahama Basin)	**~65–70 Ma**	Slow spreading (Mid-Atlantic Ridge: 20–40 mm/yr); passive margins
Indian Ocean	~150–160 Ma (Argo Abyssal Plain)	**~55–60 Ma**	Intermediate spreading; complex fracture zones
Global Total	**~180–200 Ma**	**~60–64 Ma (~60 Ma)**	Average age of all oceanic crust currently covering ~60% of Earth

4. Evaluation of Multiple Choice Options

- **(a) 20 Ma — INCORRECT:** 20 Ma is the age of young flank lithosphere near ridge crests. While a substantial fraction of seafloor is young, the overall mean including abyssal plains is three times higher (~60 Ma).
- **(b) 60 Ma — CORRECT:** Exactly corresponds to the accepted mean age of oceanic floor (empirical global average is 64.2 Ma, standard approximation 60 Ma).

- **(c) 80 Ma — INCORRECT:** 80 Ma corresponds to the age where thermal subsidence begins to flatten toward the thermal plate boundary asymptote, but it exceeds the global area-weighted mean.
- **(d) 230 Ma — INCORRECT:** 230 Ma is Triassic. Virtually no oceanic crust of this age exists in situ in the global ocean (except rare, isolated trapped relics in the Eastern Mediterranean Herodotus basin, ~280 Ma, which represent trapped Tethyan crust rather than typical open ocean floor).

Standard Textbook Reference

Ben A. van der Pluijm & Stephen Marshak, *Earth Structure* (2nd ed., Chapter 14: The Crust and Moho, Table 14.2, p. 344):
"Age: Most oceanic crust gets carried back into the mantle during subduction, so there is no oceanic crust on Earth older than about 200 Ma... The mean age of oceanic crust is about 60 Ma, whereas continental crust has an average age of >2 Ga."
C. M. R. Fowler, *The Solid Earth: An Introduction to Global Geophysics* (Cambridge University Press):
"The age of the ocean floor ranges from 0 Ma at the crest of the ridges to a maximum of about 180 Ma. The mean age of the oceanic crust is approximately 60 Ma."