

Convex sets		Comments
Set	Comments	
Convex hull: <ul style="list-style-type: none"><li>conv <math>C = \left\{\sum_{i=1}^k \theta_i \mathbf{x}_i \mid \mathbf{x}_i \in C, 0 \leq \theta \leq 1, \mathbf{1}^\top \boldsymbol{\theta} = 1\right\}</math></li></ul>	<ul style="list-style-type: none"><li>conv <math>C</math> will be the smallest convex set that contains <math>C</math>.</li><li>conv <math>C</math> will be a finite set as long as <math>C</math> is also finite.</li></ul>	
Affine hull: <ul style="list-style-type: none"><li>aff <math>C = \left\{\sum_{i=1}^k \theta_i \mathbf{x}_i \mid \mathbf{x}_i \in C \text{ for } i = 1, \cdots, k, \mathbf{1}^\top \boldsymbol{\theta} = 1\right\}</math></li></ul>	<ul style="list-style-type: none"><li><math>A</math> will be the smallest affine set that contains <math>C</math>.</li><li>Different from the convex set, <math>\theta_i</math> is not restricted between 0 and 1</li><li>aff <math>C</math> will always be an infinite set. If aff <math>C</math> contains the origin, it is also a subspace.</li></ul>	
Conic hull: <ul style="list-style-type: none"><li><math>A = \left\{\sum_{i=1}^k \theta_i \mathbf{x}_i \mid \mathbf{x}_i \in C, \theta_i &gt; 0 \text{ for } i = 1, \cdots, k\right\}</math></li></ul>	<ul style="list-style-type: none"><li><math>A</math> will be the smallest convex conic that contains <math>C</math>.</li><li>Different from the convex and affine sets, <math>\theta_i</math> does not need to sum up 1.</li></ul>	
Ray: <ul style="list-style-type: none"><li><math>\mathcal{R} = \{\mathbf{x}_0 + \theta \mathbf{v} \mid \theta \geq 0\}</math></li></ul>	<ul style="list-style-type: none"><li>The ray is an infinite set that begins in <math>\mathbf{x}_0</math> and extends infinitely in direction of <math>\mathbf{v}</math>. In other words, it has a beginning, but it has no end.</li></ul>	
Hyperplane: <ul style="list-style-type: none"><li><math>\mathcal{H} = \{\mathbf{x} \mid \mathbf{a}^\top \mathbf{x} = b\}</math></li><li><math>\mathcal{H} = \{\mathbf{x} \mid \mathbf{a}^\top (\mathbf{x} - \mathbf{x}_0) = \mathbf{0}\}</math></li><li><math>\mathcal{H} = \mathbf{x}_0 + a^\perp</math></li></ul>	<ul style="list-style-type: none"><li>It is an infinite set <math>\mathbb{R}^{n-1} \subset \mathbb{R}^n</math> that divides the space into two halfspaces.</li><li><math>a^\perp = \{\mathbf{v} \mid \mathbf{a}^\top \mathbf{v} = 0\}</math> is the set of vectors perpendicular to <math>\mathbf{a}</math>. It passes through the origin.</li><li><math>a^\perp</math> is offset from the origin by <math>\mathbf{x}_0</math>, which is any vector in <math>\mathcal{H}</math>.</li></ul>	
Halfspaces: <ul style="list-style-type: none"><li><math>\mathcal{H}_- = \{\mathbf{x} \mid \mathbf{a}^\top \mathbf{x} \leq b\}</math></li><li><math>\mathcal{H}_+ = \{\mathbf{x} \mid \mathbf{a}^\top \mathbf{x} \geq b\}</math></li></ul>	<ul style="list-style-type: none"><li>They are infinite sets of the parts divided by <math>\mathcal{H}</math>.</li></ul>	
Euclidean ball: <ul style="list-style-type: none"><li><math>B(\mathbf{x}_c, r) = \{\mathbf{x} \mid \ \mathbf{x} - \mathbf{x}_c\ _2 \leq r\}</math></li><li><math>B(\mathbf{x}_c, r) = \{\mathbf{x} \mid (\mathbf{x} - \mathbf{x}_c)^\top (\mathbf{x} - \mathbf{x}_c) \leq r\}</math></li><li><math>B(\mathbf{x}_c, r) = \{\mathbf{x}_c + r \ \mathbf{u}\  \mid \ \mathbf{u}\  \leq 1\}</math></li></ul>	<ul style="list-style-type: none"><li><math>B(\mathbf{x}_c, r)</math> is a finite set as long as <math>r &lt; \infty</math>.</li><li><math>\mathbf{x}_c</math> is the center of the ball.</li><li><math>r</math> is its radius.</li></ul>	
Ellipsoid: <ul style="list-style-type: none"><li><math>\mathcal{E} = \{\mathbf{x} \mid (\mathbf{x} - \mathbf{x}_c)^\top \mathbf{P}^{-1} (\mathbf{x} - \mathbf{x}_c) \leq 1\}</math></li><li><math>\mathcal{E} = \{\mathbf{x}_c + \mathbf{A} \mathbf{u} \mid \ \mathbf{u}\  \leq 1\}</math>, where <math>\mathbf{A} = \mathbf{P}^{1/2}</math>.</li></ul>	<ul style="list-style-type: none"><li><math>\mathcal{E}</math> is a finite set as long as <math>\mathbf{P}</math> is a finite matrix.</li><li><math>\mathbf{P}</math> is symmetric and positive definite, that is, <math>\mathbf{P} = \mathbf{P}^\top \succ \mathbf{0}</math>.</li><li><math>\mathbf{x}_c</math> is the center of the ellipsoid.</li><li>The lengths of the semi-axes are given by <math>\sqrt{\lambda_i}</math>.</li><li><math>\mathbf{A}</math> is invertible. When it is not, we say that <math>\mathcal{E}</math> is a degenerated ellipsoid (degenerated ellipsoids are also convex).</li></ul>	
Norm cone: <ul style="list-style-type: none"><li><math>C = \{(x_1, x_2, \cdots, x_n, t) \in \mathbb{R}^{n+1} \mid \mathbf{x} \in \mathbb{R}^n, \ \mathbf{x}\ _p \leq t\} \subseteq \mathbb{R}^{n+1}</math></li></ul>	<ul style="list-style-type: none"><li>Although it is named "Norm cone", it is a set, not a scalar.</li><li>The cone norm increases the dimension of <math>\mathbf{x}</math> in 1.</li><li>For <math>p = 2</math>, it is called the second-order cone, quadratic cone, Lorentz cone or ice-cream cone.</li></ul>	
Proper cone: $K \subset \mathbb{R}^n$ is a proper cone when it has the following properties <ul style="list-style-type: none"><li><math>K</math> is a convex cone, i.e., <math>\alpha K \equiv K, \alpha &gt; 0</math>.</li><li><math>K</math> is closed.</li><li><math>K</math> is solid.</li><li><math>K</math> is pointed, i.e., <math>-K \cap K = \{\mathbf{0}\}</math>.</li></ul>	<ul style="list-style-type: none"><li>The proper cone <math>K</math> is used to define the <i>generalized inequality</i> (or <i>partial ordering</i>) in some set <math>S</math>. For the generalized inequality, one must define both the proper cone <math>K</math> and the set <math>S</math>.</li><li><math>\mathbf{x} \leq \mathbf{y} \iff \mathbf{y} - \mathbf{x} \in K</math> for <math>\mathbf{x}, \mathbf{y} \in S</math> (generalized inequality)</li><li><math>\mathbf{x} &lt; \mathbf{y} \iff \mathbf{y} - \mathbf{x} \in \text{int } K</math> for <math>\mathbf{x}, \mathbf{y} \in S</math> (strict generalized inequality).</li><li>There are two cases where <math>K</math> and <math>S</math> are understood from context and the subscript <math>K</math> is dropped out:<ul style="list-style-type: none"><li>When <math>S = \mathbb{R}^n</math> and <math>K = \mathbb{R}_+^n</math> (the nonnegative orthant). In this case, <math>\mathbf{x} \leq \mathbf{y}</math> means that <math>x_i \leq y_i</math>.</li><li>When <math>S = S^n</math> and <math>K = S_+^n</math> or <math>K = S_{++}^n</math>, where <math>S^n</math> denotes the set of symmetric <math>n \times n</math> matrices, <math>S_+^n</math> is the space of the positive semidefinite matrices, and <math>S_{++}^n</math> is the space of the positive definite matrices. <math>S_+^n</math> is a proper cone in <math>S^n</math> (??). In this case, the generalized inequality <math>\mathbf{Y} \succeq \mathbf{X}</math> means that <math>\mathbf{Y} - \mathbf{X}</math> is a positive semidefinite matrix belonging to the positive semidefinite cone <math>S_+^n</math> in the subspace of symmetric matrices <math>S^n</math>. It is usual to denote <math>\mathbf{X} &gt; \mathbf{0}</math> and <math>\mathbf{X} \geq \mathbf{0}</math> to mean that <math>\mathbf{X}</math> is a positive definite and semidefinite matrix, respectively, where <math>\mathbf{0} \in \mathbb{R}^{n \times n}</math> is a zero matrix.</li></ul></li><li>Another common usage is when <math>S = \mathbb{R}^n</math> and <math>K = \{\mathbf{e} \in \mathbb{R}^n \mid c_1 + c_2 t + \cdots + c_n t^{n-1} \geq 0, \text{ for } 0 \leq t \leq 1\}</math>. In this case, <math>\mathbf{x} \leq_K \mathbf{y}</math> means that <math>x_1 + x_2 t + \cdots + x_n t^{n-1} \leq y_1 + y_2 t + \cdots + y_n t^{n-1}</math>.</li><li>The generalized inequality has the following properties:<ul style="list-style-type: none"><li>If <math>\mathbf{x} \leq_K \mathbf{y}</math> and <math>\mathbf{u} \leq_K \mathbf{v}</math>, then <math>\mathbf{x} + \mathbf{u} \leq_K \mathbf{y} + \mathbf{v}</math> (preserve under addition).</li><li>If <math>\mathbf{x} \leq_K \mathbf{y}</math> and <math>\mathbf{y} \leq_K \mathbf{z}</math>, then <math>\mathbf{x} \leq_K \mathbf{z}</math> (transitivity).</li><li>If <math>\mathbf{x} \leq_K \mathbf{y}</math>, then <math>\alpha \mathbf{x} \leq_K \mathbf{y}</math> for <math>\alpha \geq 0</math> (preserve under nonnegative scaling).</li><li><math>\mathbf{x} \leq_K \mathbf{x}</math> (reflexivity).</li><li>If <math>\mathbf{x} \leq_K \mathbf{y}</math> and <math>\mathbf{y} \leq_K \mathbf{x}</math>, then <math>\mathbf{x} = \mathbf{y}</math> (antisymmetric).</li><li>If <math>\mathbf{x}_i \leq_K \mathbf{y}_i</math>, for <math>i = 1, 2, \dots</math>, and <math>\mathbf{x}_i \rightarrow \mathbf{x}</math> and <math>\mathbf{y}_i \rightarrow \mathbf{y}</math> as <math>i \rightarrow \infty</math>, then <math>\mathbf{x} \leq_K \mathbf{y}</math>.</li></ul></li><li>It is called partial ordering because <math>\mathbf{x} \not\leq_K \mathbf{y}</math> and <math>\mathbf{y} \not\leq_K \mathbf{x}</math> for many <math>\mathbf{x}, \mathbf{y} \in S</math>. When it happens, we say that <math>\mathbf{x}</math> and <math>\mathbf{y}</math> are not comparable (this case does not happen in ordinary inequality, <math>&lt;</math> and <math>&gt;</math>).</li><li><math>\mathbf{x} \in S</math> is the <i>minimum</i> element of <math>S</math> with respect to the proper cone <math>K</math> if <math>\mathbf{x} \leq_K \mathbf{y}, \forall \mathbf{y} \in S</math> (for <i>maximum</i>, <math>\mathbf{x} \succeq_K \mathbf{y}, \forall \mathbf{y} \in S</math>). It means that <math>S \subseteq \mathbf{x} + K</math> (for the maximum, <math>S \subseteq \mathbf{x} - K</math>), where <math>\mathbf{x} + K</math> denotes the set <math>K</math> shifted from the origin by <math>\mathbf{x}</math>. Note that any point in <math>K + \mathbf{x}</math> is comparable with <math>\mathbf{x}</math> and is greater or equal to <math>\mathbf{x}</math> in the generalized inequality mean. The set <math>S</math> does not necessarily have a minimum (maximum), but the minimum (maximum) is unique if it does.</li><li><math>\mathbf{x} \in S</math> is the <i>minimal</i> element of <math>S</math> with respect to the proper cone <math>K</math> if <math>\mathbf{y} \leq_K \mathbf{x}</math> only when <math>\mathbf{y} = \mathbf{x}</math> (for the <i>maximal</i>, <math>\mathbf{y} \geq_K \mathbf{x}</math> only when <math>\mathbf{y} = \mathbf{x}</math>). It means that <math>(\mathbf{x} - K) \cap S = \{\mathbf{x}\}</math> for minimal (for the maximal <math>(\mathbf{x} + K) \cap S = \{\mathbf{x}\}</math>), where <math>\mathbf{x} - K</math> denotes the reflected set <math>K</math> shift by <math>\mathbf{x}</math>. Note that any point in <math>\mathbf{x} - K</math> is comparable with <math>\mathbf{x}</math> and is less than or equal to <math>\mathbf{x}</math> in the generalized inequality mean. The set <math>S</math> can have many different minimal (maximal) elements.</li><li>When <math>K = \mathbb{R}_+</math> and <math>S = \mathbb{R}</math> (ordinary inequality), the minimal is equal to the minimum and the maximal is equal to the maximum.</li><li>When we say that a scalar-valued function <math>f: \mathbb{R}^n \rightarrow \mathbb{R}</math> is nondecreasing, it means that whenever <math>\mathbf{u} \leq \mathbf{v}</math>, we have <math>\hat{h}(\mathbf{u}) \leq \hat{h}(\mathbf{v})</math>. Similar results hold for decreasing, increasing, and nonincreasing scalar functions.</li></ul>	
Dual cone: <ul style="list-style-type: none"><li><math>K^* = \{\mathbf{y} \mid \mathbf{x}^\top \mathbf{y} \geq 0, \forall \mathbf{x} \in K\}</math></li></ul>	<ul style="list-style-type: none"><li><math>K^*</math> is a cone, and it is convex even when the original cone <math>K</math> is nonconvex.</li><li><math>K^*</math> has the following properties:<ul style="list-style-type: none"><li><math>K^*</math> is closed and convex.</li><li><math>K_1 \subseteq K_2</math> implies <math>K_1^* \supseteq K_2^*</math>.</li><li>If <math>K</math> has a nonempty interior, then <math>K^*</math> is pointed.</li><li>If the closure of <math>K</math> is pointed then <math>K^*</math> has a nonempty interior.</li><li><math>K^{**}</math> is the closure of the convex hull of <math>K</math>. Hence, if <math>K</math> is convex and closed, <math>K^{**} = K</math>.</li></ul></li></ul>	
Polyhedra: <ul style="list-style-type: none"><li><math>\mathcal{P} = \left\{\mathbf{x} \mid \mathbf{a}_j^\top \mathbf{x} \leq b_j, j = 1, \dots, m, \mathbf{a}_j^\top \mathbf{x} = d_j, j = 1, \dots, p\right\}</math></li><li><math>\mathcal{P} = \{\mathbf{x} \mid \mathbf{A} \mathbf{x} \leq \mathbf{b}, \mathbf{C} \mathbf{x} = \mathbf{d}\}</math>, where <math>\mathbf{A} = \begin{bmatrix} \mathbf{a}_1 &amp; \mathbf{a}_2 &amp; \dots &amp; \mathbf{a}_m \end{bmatrix}^\top</math> and <math>\mathbf{C} = \begin{bmatrix} \mathbf{c}_1 &amp; \mathbf{c}_2 &amp; \dots &amp; \mathbf{c}_m \end{bmatrix}^\top</math></li></ul>	<ul style="list-style-type: none"><li>The polyhedron may or may not be an infinite set.</li><li>Polyhedron is the result of the intersection of <math>m</math> halfspaces and <math>p</math> hyperplanes.</li><li>Subspaces, hyperplanes, lines, rays line segments, and halfspaces are all special cases of polyhedra.</li><li>The <i>nonnegative orthant</i>, <math>\mathbb{R}_+^n = \{\mathbf{x} \in \mathbb{R}^n \mid x_i \leq 0 \text{ for } i = 1, \dots, n\} = \{\mathbf{x} \in \mathbb{R}^n \mid \mathbf{I} \mathbf{x} \geq \mathbf{0}\}</math>, is a special polyhedron.</li></ul>	
Simplex: <ul style="list-style-type: none"><li><math>\mathcal{S} = \text{conv} \left\{\mathbf{v}_m\right\}_{m=0}^k = \left\{\sum_{i=0}^k \theta_i \mathbf{v}_i \mid \mathbf{0} \leq \theta \leq 1, \mathbf{1}^\top \boldsymbol{\theta} = 1\right\}</math></li><li><math>\mathcal{S} = \{\mathbf{x} \mid \mathbf{x} = \mathbf{v}_0 + \mathbf{V} \boldsymbol{\theta}\}</math>, where <math>\mathbf{V} = \begin{bmatrix} \mathbf{v}_1 - \mathbf{v}_0 &amp; \dots &amp; \mathbf{v}_n - \mathbf{v}_0 \end{bmatrix} \in \mathbb{R}^{n \times k}</math></li><li><math>\mathcal{S} = \{\mathbf{x} \mid \underbrace{\mathbf{A}_1 \mathbf{x} \leq \mathbf{A}_1 \mathbf{v}_0, \mathbf{1}^\top \mathbf{A}_1 \mathbf{x} \leq 1 + \mathbf{1}^\top \mathbf{A}_1 \mathbf{v}_0}_{\text{Linear inequalities in } \mathbf{x}}, \underbrace{\mathbf{A}_2 \mathbf{x} = \mathbf{A}_2 \mathbf{v}_0}_{\text{Linear equalities in } \mathbf{x}}\}</math> (Polyhedra form), where <math>\mathbf{A} = \begin{bmatrix} \mathbf{A}_1 \\ \mathbf{A}_2 \end{bmatrix}</math> and <math>\mathbf{A} \mathbf{V} = \begin{bmatrix} \mathbf{I}_{k \times k} \\ \mathbf{0}_{n-k \times n-k} \end{bmatrix}</math></li></ul>	<ul style="list-style-type: none"><li>Simplexes are a subfamily of the polyhedra set.</li><li>Also called k-dimensional Simplex in <math>\mathbb{R}^n</math>.</li><li>The set <math>\{\mathbf{v}_m\}_{m=0}^k</math> is a affinely independent, which means <math>\{\mathbf{v}_1 - \mathbf{v}_0, \dots, \mathbf{v}_k - \mathbf{v}_0\}</math> are linearly independent.</li><li><math>\mathbf{V} \in \mathbb{R}^{n \times k}</math> is a full-rank tall matrix, i.e., <math>\text{rank}(\mathbf{V}) = k</math>. All its column vectors are independent. The matrix <math>\mathbf{A}</math> is its left pseudoinverse.</li></ul>	
$\alpha$ -sublevel set: <ul style="list-style-type: none"><li><math>C_\alpha = \{\mathbf{x} \in \text{dom}(f) \mid f(\mathbf{x}) \leq \alpha\}</math> (regarding convexity), where <math>f: \mathbb{R}^n \rightarrow \mathbb{R}</math></li><li><math>C_\alpha = \{\mathbf{x} \in \text{dom}(f) \mid f(\mathbf{x}) \geq \alpha\}</math> (regarding concavity), where <math>f: \mathbb{R}^n \rightarrow \mathbb{R}</math></li></ul>	<ul style="list-style-type: none"><li>If <math>f</math> is a convex (concave) function, then sublevel sets of <math>f</math> are convexes (concaves) for any <math>\alpha \in \mathbb{R}</math>.</li><li>The converse is not true: a function can have all its sublevel set convex and not be a convex function.</li><li><math>C_\alpha \subseteq \text{dom}(f)</math></li></ul>	
Functions (or operators) and their implications regarding convexity		
Function	Convex (concave)?	Comments
Union: $C = A \cup B$	Not in most of the cases.	
Intersection: $C = A \cap B$	Yes, if $A$ and $B$ are convex sets.	
Convex function: $f: \text{dom}(f) \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(\theta \mathbf{x} + (1 - \theta) \mathbf{y}) \leq \theta f(\mathbf{x}) + (1 - \theta) f(\mathbf{y})</math>, where <math>0 \leq \theta \leq 1</math>.</li><li><math>\text{dom}(f)</math> shall be a convex set to <math>f</math> be considered a convex function.</li></ul>	Yes. <ul style="list-style-type: none"><li><math>f</math> is convex iff <math>\mathbf{P} \geq \mathbf{0}</math>.</li><li><math>f</math> is strictly convex iff <math>\mathbf{P} &gt; \mathbf{0}</math>.</li><li><math>f</math> is concave iff <math>\mathbf{P} \leq \mathbf{0}</math>.</li><li><math>f</math> is strictly concave iff <math>\mathbf{P} &lt; \mathbf{0}</math>.</li></ul>	<ul style="list-style-type: none"><li>Graphically, the line segment between <math>(\mathbf{x}, f(\mathbf{x}))</math> and <math>(\mathbf{y}, f(\mathbf{y}))</math> lies always above the graph <math>f</math>.</li><li>In terms of sets, a function is convex iff a line segment within <math>\text{dom}(f)</math>, which is a convex set, gives an image set that is also convex.</li><li><math>\text{dom} f</math> is convex iff all points for any line segment within <math>\text{dom}(f)</math> belong to it.</li><li><i>First-order condition</i>: <math>f</math> is convex iff <math>\text{dom}(f)</math> is convex and <math>f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}), \forall \mathbf{x}, \mathbf{y} \in \text{dom}(f), \mathbf{x} \neq \mathbf{y}</math>, being <math>\nabla f(\mathbf{x})</math> the gradient vector. This inequation says that the first-order Taylor approximation is a <i>underestimator</i> for convex functions. The first-order condition requires that <math>f</math> is differentiable.</li><li>If <math>\nabla f(\mathbf{x}) = \mathbf{0}</math>, then <math>f(\mathbf{y}) \geq f(\mathbf{x}), \forall \mathbf{y} \in \text{dom}(f)</math> and <math>\mathbf{x}</math> is a global minimum.</li><li><i>Second-order condition</i>: <math>f</math> is convex iff <math>\text{dom}(f)</math> is convex and <math>\mathbf{H} \geq \mathbf{0}</math>, that is, the Hessian matrix <math>\mathbf{H}</math> is a positive semidefinite matrix. It means that the graphic of the curvature has a positive (upward) curvature at <math>\mathbf{x}</math>. It is important to note that, if <math>\mathbf{H} &gt; \mathbf{0}, \forall \mathbf{x} \in \text{dom}(f)</math>, then <math>f</math> is strictly convex. But is <math>f</math> is strictly convex, not necessarily that <math>\mathbf{H} &gt; \mathbf{0}, \forall \mathbf{x} \in \text{dom}(f)</math>. Therefore, strict convexity can only be partially characterized.</li></ul>
Affine function $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$ <ul style="list-style-type: none"><li><math>f(\mathbf{x}) = \mathbf{A} \mathbf{x} + \mathbf{b}</math>, where <math>\mathbf{A} \in \mathbb{R}^{m \times n}, \mathbf{b} \in \mathbb{R}^m, \mathbf{x} \in \mathbb{R}^n</math></li></ul>	Yes, if the domain $S \subseteq \mathbb{R}^n$ is a convex set, then its image $f(S) = \{f(\mathbf{x}) \mid \mathbf{x} \in S\} \subseteq \mathbb{R}^m$ is also convex.	<ul style="list-style-type: none"><li>The affine function, <math>f(\mathbf{x}) = \mathbf{A} \mathbf{x} + \mathbf{b}</math>, is a broader category that encompasses the linear function, <math>f(\mathbf{x}) = \mathbf{A} \mathbf{x}</math>. The linear function has its origin fixed at <math>\mathbf{0}</math> after the transformation, whereas the affine function does not necessarily have it (when not, this makes the affine function nonlinear). Graphically, we can think of an affine function as a linear transformation plus a shift from the origin of <math>\mathbf{b}</math>.</li><li>A special case of the linear function is when <math>\mathbf{A} = \mathbf{c}^\top</math>. In this case, we have <math>f(\mathbf{x}) = \mathbf{c}^\top \mathbf{x}</math>, which is the inner product between the vector <math>\mathbf{c}</math> and <math>\mathbf{x}</math>.</li><li>The inverse image of <math>C, f^{-1}(C) = \{\mathbf{x} \mid f(\mathbf{x}) \in C\}</math>, is also convex.</li><li>The <i>linear matrix inequality</i> (LMI), <math>\mathbf{A}(\mathbf{x}) = x_1 \mathbf{A}_1 + \cdots + x_n \mathbf{A}_n \leq \mathbf{B}</math>, is a special case of affine function. In other words, <math>f(S) = \{\mathbf{x} \mid \mathbf{A}(\mathbf{x}) \leq \mathbf{B}\}</math> is a convex set if <math>S</math> is convex. Many optimization problems can be formulated as LMI problems and solved optimally.</li></ul>
Exponential function $f: \mathbb{R} \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(x) = e^{ax} \in \mathbb{R}</math>, where <math>a \in \mathbb{R}</math></li></ul>	Yes.	
Quadratic function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(\mathbf{x}) = \mathbf{a} \mathbf{x}^\top \mathbf{P} \mathbf{x} + \mathbf{p}^\top \mathbf{x} + r \in \mathbb{R}</math>, where <math>\mathbf{x}, \mathbf{p} \in \mathbb{R}^n, \mathbf{P} \in \mathbb{R}^{n \times n}</math>, and <math>a, b \in \mathbb{R}</math></li></ul>	It depends on the matrix $\mathbf{P}$ : <ul style="list-style-type: none"><li><math>f</math> is convex iff <math>\mathbf{P} \geq \mathbf{0}</math>.</li><li><math>f</math> is strictly convex iff <math>\mathbf{P} &gt; \mathbf{0}</math>.</li><li><math>f</math> is concave iff <math>\mathbf{P} \leq \mathbf{0}</math>.</li><li><math>f</math> is strictly concave iff <math>\mathbf{P} &lt; \mathbf{0}</math>.</li></ul>	
Power function $f: \mathbb{R}_{++} \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(x) = x^a</math></li></ul>	It depends on $a$ <ul style="list-style-type: none"><li><math>f</math> is convex iff <math>a \geq 1</math> or <math>a \leq 0</math>.</li><li><math>f</math> is concave iff <math>0 \leq a \leq 1</math>.</li></ul>	
Power of absolute value: $f: \mathbb{R} \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(x) =  x ^p</math>, where <math>p \leq 1</math>.</li></ul>	Yes.	
Logarithm function: $f: \mathbb{R}_{++} \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(x) = \log x</math></li></ul>	Yes.	
Negative entropy function: $f: \mathbb{R}_+ \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(x) = x \log x</math></li></ul>	Yes	<ul style="list-style-type: none"><li>When it is defined <math>f(x) _{x=0} = 0</math>, <math>\text{dom}(f) = \mathbb{R}</math>.</li></ul>
Minkowski distance, $p$ -norm function, or $l_p$ norm function: $f: \mathbb{R}^n \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(\mathbf{x}) = \ \mathbf{x}\ _p</math>, where <math>p \in \mathbb{N}_{++}</math>.</li></ul>	Yes.	<ul style="list-style-type: none"><li>It can be proved by triangular inequality.</li></ul>
Maximum function: $f: \mathbb{R}^n \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(\mathbf{x}) = \max\{x_1, \dots, x_n\}</math>.</li></ul>	Yes.	
Pointwise maximum (maximum function): $f: \mathbb{R}^n \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(\mathbf{x}) = \max\{f_1(\mathbf{x}), \dots, f_n(\mathbf{x})\}</math>.</li></ul>	Yes, if $f_1, \dots, f_n$ are convex function.	<ul style="list-style-type: none"><li>Its domain <math>\text{dom}(f) = \bigcap_{i=1}^n \text{dom}(f_i)</math> is also convex.</li></ul>
Pointwise infimum: <ul style="list-style-type: none"><li><math>f(x) = \inf_{y \in \mathcal{A}} g(x, y)</math>.</li></ul>	<ul style="list-style-type: none"><li>.</li></ul>	<ul style="list-style-type: none"><li>For each value of <math>x</math>, we have an infinite set of points <math>g(x, y) _{y \in \mathcal{A}}</math>. The value <math>f(x)</math> will be the greatest value in the codomain of <math>f</math> that is less than or equal this set.</li><li><math>\text{dom}(f) = \left\{x \mid (x, y) \in \text{dom}(g) \ \forall y \in \mathcal{A}, \inf_{y \in \mathcal{A}} g(x, y) &gt; -\infty\right\}</math>.</li></ul>
Pointwise supremum: <ul style="list-style-type: none"><li><math>f(x) = \sup_{y \in \mathcal{A}} g(x, y)</math>.</li></ul>	<ul style="list-style-type: none"><li><math>f</math> is convex in <math>x</math> if <math>g</math> is convex in <math>x</math> for each <math>y \in \mathcal{A}</math>.</li><li><math>f</math> is concave in <math>x</math> if <math>g</math> is concave in <math>x</math> for each <math>y \in \mathcal{A}</math>.</li></ul>	<ul style="list-style-type: none"><li>For each value of <math>x</math>, we have an infinite set of points <math>g(x, y) _{y \in \mathcal{A}}</math>. The value <math>f(x)</math> will be the least value in the codomain of <math>f</math> that is greater than or equal this set.</li><li><math>\text{dom}(f) = \left\{x \mid (x, y) \in \text{dom}(g) \ \forall y \in \mathcal{A}, \sup_{y \in \mathcal{A}} g(x, y) &lt; \infty\right\}</math>.</li><li>In terms of epigraphs, the pointwise supremum of the infinite set of functions <math>g(x, y) _{y \in \mathcal{A}}</math> corresponds to the intersection of the following epigraphs: <math>\text{epi } f = \bigcap_{y \in \mathcal{A}} \text{epi } g(\cdot, y)</math></li></ul>
Minimum function: $f: \mathbb{R}^n \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(\mathbf{x}) = \min\{f_1(\mathbf{x}), \dots, f_n(\mathbf{x})\}</math>.</li></ul>	Not in most of the cases.	
Log-sum-exp function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(\mathbf{x}) = \log(e^{x_1} + \cdots + e^{x_n})</math></li></ul>	Yes.	<ul style="list-style-type: none"><li>This function is interpreted as the approximation of the maximum element function, since <math>\max\{x_1, \dots, x_n\} \leq f(\mathbf{x}) \leq \max\{x_1, \dots, x_n\} + \log n</math></li></ul>
Geometric (mean) function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(\mathbf{x}) = (\prod_{i=1}^n x_i)^{1/n}</math></li></ul>	Yes	
Log-determinant function $f: S_{++}^n \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(\mathbf{X}) = \log  \mathbf{X} </math></li></ul>	Yes	<ul style="list-style-type: none"><li><math>\mathbf{X}</math> is positive semidefinite, i.e., <math>\mathbf{X} &gt; \mathbf{0} : \mathbf{X} \in S_{++}^n</math>.</li></ul>
Composite function $f = h \circ g: \mathbb{R}^n \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f = g \circ h</math>, i.e., <math>f(\mathbf{x}) = (h \circ g)(\mathbf{x}) = h(g(\mathbf{x}))</math>, where:<ul style="list-style-type: none"><li><math>g: \mathbb{R}^n \rightarrow \mathbb{R}^{m+1}</math> is an affine function given by <math>g(\mathbf{x}) = \begin{bmatrix} \mathbf{A} \\ \mathbf{c}^\top \end{bmatrix} \mathbf{x} + \begin{bmatrix} \mathbf{b} \\ d \end{bmatrix}</math>, being <math>\mathbf{A} \in \mathbb{R}^{m \times n}, \mathbf{b} \in \mathbb{R}^m, \mathbf{c} \in \mathbb{R}^n</math>, and <math>d \in \mathbb{R}</math>.</li><li><math>\text{dom}(f) = \{\mathbf{x} \in \text{dom}(g) \mid g(\mathbf{x}) \in \text{dom}(h)\}</math>.</li></ul></li><li><math>f(\mathbf{x}) = \mathcal{P}^{-1}(\mathbf{Q} \mathcal{P}(\mathbf{x}))</math><ul style="list-style-type: none"><li><math>\mathcal{P}(\mathbf{x}) = \{(x, t) \mid t \geq 0\} \subset \mathbb{R}^{n+1}</math></li><li><math>\mathbf{Q} = \begin{bmatrix} \mathbf{A} &amp; \mathbf{b} \\ \mathbf{c}^\top &amp; d \end{bmatrix} \in \mathbb{R}^{(m+1) \times (n+1)}</math></li></ul></li></ul>	<ul style="list-style-type: none"><li>Scalar composition: the following statements hold for <math>k = 1</math> and <math>n \geq 1</math>, i.e., <math>h: \mathbb{R} \rightarrow \mathbb{R}</math> and <math>g: \mathbb{R}^n \rightarrow \mathbb{R}</math>:<ul style="list-style-type: none"><li><math>f</math> is convex if <math>h</math> is convex, <math>\tilde{h}</math> is nondecreasing, and <math>g</math> is convex. In this case, <math>\text{dom}(h)</math> is either <math>(-\infty, a]</math> or <math>(-\infty, a)</math>.</li><li><math>f</math> is convex if <math>h</math> is convex, <math>\tilde{h}</math> is nonincreasing, and <math>g</math> is concave. In this case, <math>\text{dom}(h)</math> is either <math>[a, \infty)</math> or <math>(a, \infty)</math>.</li><li><math>f</math> is concave if <math>h</math> is concave, <math>\tilde{h}</math> is nondecreasing, and <math>g</math> is convex.</li><li><math>f</math> is concave if <math>h</math> is concave, <math>\tilde{h}</math> is nonincreasing, and <math>g</math> is concave.</li></ul></li><li>Vector composition: the following statements hold for <math>k \geq 1</math> and <math>p \geq 1</math>, i.e., <math>h: \mathbb{R}^k \rightarrow \mathbb{R}</math> and <math>g: \mathbb{R}^n \rightarrow \mathbb{R}^k</math>. Hence, <math>g(\mathbf{x}) = (g_1(\mathbf{x}), g_2(\mathbf{x}), \dots, g_k(\mathbf{x}))</math> is a vector-valued function (or simply, vector function), where <math>g_i: \mathbb{R}^k \rightarrow \mathbb{R}</math> for <math>1 \leq i \leq k</math>.<ul style="list-style-type: none"><li><math>f</math> is convex if <math>h</math> is convex, <math>\tilde{h}</math> is nondecreasing in each argument of <math>\mathbf{x}</math>, and <math>\{g_i\}_{i=1}^k</math> is a set of convex functions.</li><li><math>f</math> is convex if <math>h</math> is convex, <math>\tilde{h}</math> is nonincreasing in each argument of <math>\mathbf{x}</math>, and <math>\{g_i\}_{i=1}^k</math> is a set of concave functions</li><li><math>f</math> is concave if <math>h</math> is convex, <math>\tilde{h}</math> is nondecreasing in each argument of <math>\mathbf{x}</math>, and <math>\{g_i\}_{i=1}^k</math> is a set of concave functions.</li></ul></li></ul> <p>Where <math>\tilde{h}</math> is the extended-value extension of the function <math>h</math>, which assigns the value <math>\infty</math> (<math>-\infty</math>) to the point not in <math>\text{dom}(h)</math> for <math>h</math> convex (concave).</p>	<ul style="list-style-type: none"><li>The composition function allows us to see a large class of functions as convex (or concave).</li><li>For scale composition, the remarkable ones are:<ul style="list-style-type: none"><li>If <math>g</math> is convex then <math>f(\mathbf{x}) = h(g(\mathbf{x}))</math> is convex.</li><li>If <math>g</math> is concave and <math>\text{dom}(g) \subseteq \mathbb{R}_{++}</math>, then <math>f(\mathbf{x}) = h(g(\mathbf{x})) = \log g(\mathbf{x})</math> is concave.</li><li>If <math>g</math> is concave and <math>\text{dom}(g) \subseteq \mathbb{R}_{++}</math>, then <math>f(\mathbf{x}) = h(g(\mathbf{x})) = 1/g(\mathbf{x})</math> is convex.</li><li>If <math>g</math> is convex and <math>\text{dom}(g) \subseteq \mathbb{R}_{++}</math>, then <math>f(\mathbf{x}) = h(g(\mathbf{x})) = g^p(\mathbf{x})</math> is convex, where <math>p \geq 1</math>.</li><li>If <math>g</math> is convex then <math>f(\mathbf{x}) = h(g(\mathbf{x})) = -\log(-g(\mathbf{x}))</math> is convex, where <math>\text{dom}(f) = \{\mathbf{x} \mid g(\mathbf{x}) &lt; 0\}</math>.</li></ul></li><li>For vector composition, we have the following examples:<ul style="list-style-type: none"><li>If <math>g(\mathbf{x}) = \mathbf{A} \mathbf{x} + \mathbf{b}</math> is an affine function, then <math>f = h \circ g</math> is convex (concave) if <math>h</math> is convex (concave).</li><li>Let <math>h(\mathbf{x}) = x_{[1]} + \cdots + x_{[r]}</math> be the sum of the <math>r</math> largest components of <math>\mathbf{x} \in \mathbb{R}^k</math>. If <math>g_1, g_2, \dots, g_k</math> are convex, where <math>\text{dom}(g_i) = \mathbb{R}^n</math>, then <math>f = h \circ g</math>, which is the pointwise sum of the largest <math>g_i</math>'s, is convex.</li><li><math>f = h \circ g</math> is a convex function when <math>h(\mathbf{x}) = \log\left(\sum_{i=1}^n e^{x_i}\right)</math> and <math>g_1, g_2, \dots, g_k</math> are convex functions.</li><li>For <math>0 &lt; p \leq 1</math>, the function <math>h(\mathbf{x}) = \left(\sum_{i=1}^n x_i^p\right)^{1/p}</math>, where <math>\text{dom}(h) = \mathbb{R}_+^n</math>, is concave. If <math>g_1, g_2, \dots, g_k</math> are concaves (convexes) and nonnegatives, then <math>f = h \circ g</math> is concave (convex).</li></ul></li></ul>
Nonnegative weighted sum: $f: \text{dom}(f) \rightarrow \mathbb{R}$ <ul style="list-style-type: none"><li><math>f(\mathbf{x}) = \sum_{i=1}^n w_i f_i(\mathbf{x})</math>, where <math>w \geq 0</math>.</li></ul>	<ul style="list-style-type: none"><li>If <math>f_1, f_2, \dots, f_n</math> are convex or concave functions, then <math>f</math> is a convex or concave function, respectively.</li><li>If <math>f_1, f_2, \dots, f_n</math></li></ul>	

