# List of Symbols

Version: December 6, 2022

### 1 Font notation

$a, b, c, \ldots, A, B, C, \ldots$	Scalars
$a, b, c, \dots$	Vectors
$A, B, C, \dots$	Matrices
$\mathcal{A},\mathcal{B},\mathcal{C},\dots$	Tensors
$A, B, C, \ldots, A, B, C, \ldots$	Sets

## 2 Common symbols

$\mathbf{\nabla} f, \mathbf{g}$	Gradient vector
$\nabla_x f, \mathbf{g}_x$	Gradient vector with respect $x$
$\mathbf{g}$ (or $\hat{\mathbf{g}}$ if the gradient vector is $\mathbf{g}$ )	Stochastic approximation of the gra-
0 ( 0 0	dient vector
$J(\cdot)$	Cost-function or objective function
$\Lambda(\cdot)$	Likelihood function
$\Lambda_l(\cdot)$	Log-likelihood function
$\mathcal{O}(\cdot), \mathcal{O}(\cdot)$	big-O notation
Q(x)	Q-function
$\mu_x, \mathbf{m}_x$	Mean vector
$\hat{\boldsymbol{\mu}}_{x}^{'},\hat{\mathbf{m}}_{x}$	Sample mean vector
$r_x( au), R_x( au)$	Autocorrelation function of the sig-
	$\operatorname{nal} x(t) \text{ or } x[n]$
$\hat{r}_{\scriptscriptstyle X}( au),\hat{R}_{\scriptscriptstyle X}( au)$	Estimated autocorrelation function
	of the signal $x(t)$ or $x[n]$
$R_x$	(Auto)correlation matrix of $\mathbf{x}$
$\hat{\mathbf{R}}_{\mathbf{x}}$	Sample (auto)correlation matrix
$r_{x,d}(\tau), R_{x,d}(\tau)$	Cross-correlation between $x[n]$ and
	d[n] or $x(t)$ and $d(t)$
$\hat{r}_{x,d}( au), \hat{R}_{x,d}( au)$	Estimated cross-correlation between
	x[n] and $d[n]$ or $x(t)$ and $d(t)$

$\mathbf{R}_{\mathbf{x}\mathbf{y}}$	Cross-correlation matrix of $\mathbf{x}$ and $\mathbf{y}$
$\hat{\mathbf{R}}_{\mathbf{x}\mathbf{y}}$	Sample cross-correlation matrix of
n ,	$R_{xy}$ Cross-correlation vector
$\mathbf{P}_{\mathbf{x}d}$ $\rho_{x,y}$	Pearson correlation coefficient be-
<i>PX</i> , <i>y</i>	tween $x$ and $y$
$\hat{ ho}_{x,y}$	Estimated Pearson correlation coefficient between $x$ and $y$
$c_X(\tau), C_X(\tau)$	Autocovariance function of the signal $x(t)$ or $x[n]$
$\hat{c}_x( au), \hat{C}_x( au)$	Estimated autocovariance function of the signal $x(t)$ or $x[n]$
$C_x, K_x, \Sigma_x$	(Auto)covariance matrix of $\mathbf{x}$
$\hat{\mathbf{C}}_{\mathbf{x}}, \hat{\mathbf{K}}_{\mathbf{x}}, \hat{\mathbf{\Sigma}}_{\mathbf{x}}$	Sample (auto)covariance matrix
$c_{xy}(\tau), C_{xy}(\tau)$	Cross-covariance function of the sig-
	$\operatorname{nal} x(t) \text{ or } x[n]$
$\hat{c}_{xy}( au), \hat{C}_{xy}( au)$	Estimated cross-covariance function
$\mathrm{C}_{\mathrm{xy}}, \mathrm{K}_{\mathrm{xy}}, \Sigma_{\mathrm{xy}}$	of the signal $x(t)$ or $x[n]$ Cross-covariance matrix of $\mathbf{x}$
$\hat{\mathbf{C}}_{\mathbf{xy}},\hat{\mathbf{K}}_{\mathbf{xy}},\mathbf{\Sigma}_{\mathbf{xy}}$ $\hat{\mathbf{C}}_{\mathbf{xy}},\hat{\mathbf{K}}_{\mathbf{xy}},\hat{\mathbf{\Sigma}}_{\mathbf{xy}}$	Sample cross-covariance matrix
$\delta(t)$	Delta function
$\delta[n]$	Kronecker function
h(t), h[n]	Impulse response (continuous and discrete time)
$\mathbf{C}$	Cofactor matrix
$\mathbf{W}, \mathbf{D}$	Diagonal matrix
$\mathbf{w}, \mathbf{\theta}$	Parameters, coefficients, or weights vector
$\mathbf{w}_{o}, \mathbf{w}^{\star}, \mathbf{\theta}_{o}, \mathbf{\theta}^{\star}$	Optimum value of the parameters,
	coefficients, or weights vector
$\mathbf{W}$	Matrix of the weights
P	Projection matrix; Permutation ma-
	trix
Λ L	Eigenvalue matrix
U	Lower matrix Upper matrix; Left singular vectors
$\mathbf{U}_r$	Left singular nondegenerated vectors
$\Sigma$	Singular value matrix
$\Sigma_r$	Singular value matrix with nonzero
	singular values in the main diagonal
$\Sigma^+$	Singular value matrix of the pseudoinverse
$\Sigma_r^+$	Singular value matrix of the pseudoinverse with nonzero singular values in the main diagonal

$egin{array}{c} {f V} \\ {f V}_r \end{array}$	Right singular vectors Right singular nondegenerated vectors
J	Jordan matrix; Jacobian matrix
$\mathbf{S}$	Symmetric matrix
Q	Orthogonal matrix
$\mathbf{I}_N$	$N \times N$ -dimensional identity matrix
$0_{M imes N}$	$M \times N$ -dimensional null matrix
$0_N$	N-dimensional null vector
0	Null matrix, vector, or tensor (di-
	mensionality understood by context)
$1_{M  imes N}$	$M \times N$ -dimensional ones matrix
$1_N$	N-dimensional ones vector
1	Ones matrix, vector, or tensor (di-
	mensionality understood by context)
j	$\sqrt{-1}$

## ${\bf 3}\quad {\bf Linear~Algebra~operations}$

. 1	
$\mathbf{A}^{-1}$	Inverse matrix
$\mathbf{A}^+,\mathbf{A}^\dagger$	Moore-Penrose pseudoinverse
$\mathbf{A}^{ op}$	Transpose
$\mathbf{A}^*$	Complex conjugate
$\mathbf{A}^H$	Hermitian
$\ \mathbf{A}\ _{\mathrm{F}}$	Frobenius norm
$\ \mathbf{A}\ $	Matrix norm
$\ \mathbf{a}\ $	$l_1$ norm, 1-norm, or Manhatan norm
$\ \mathbf{a}\ , \ \mathbf{a}\ _2$	$l_2$ norm, 2-norm, or Euclidean norm
$\ \mathbf{a}\ _p$	$l_p$ norm, $p$ -norm, or Minkowski norm
$\ \mathbf{a}\ _{\infty}^{'}$	$l_{\infty}$ norm, $\infty$ -norm, or Chebyshev
	norm
$ \mathbf{A} , \det{(\mathbf{A})}$	Determinant
$\operatorname{diag}\left(\mathbf{a}\right),\operatorname{diag}\left(\mathbf{A}\right)$	Diagonalization: a square, diagonal
	matrix with entries given by the vec-
	tor a or the elements in the diagonal
	of ${f A}$
$\text{vec}(\mathbf{A})$	Vectorization: stacks the columns of
. ,	the matrix A into a long column vec-
	tor
$\operatorname{vec}_{\operatorname{d}}\left(\mathbf{A}\right)$	Extracts the diagonal elements of a
	square matrix and returns them in a
	column vector

(4)	
$\operatorname{vec}_{\operatorname{l}}\left(\mathbf{A}\right)$	Extracts the elements strictly below
	the main diagonal of a square matrix in a column-wise manner and returns
	them into a column vector
**************************************	
$\operatorname{vec}_{\mathrm{u}}\left(\mathbf{A}\right)$	Extracts the elements strictly above
	the main diagonal of a square matrix
	in a column-wise manner and returns
(A)	them into a column vector
$\mathrm{vec_{b}}\left(\mathbf{A} ight)$	Block vectorization operator: stacks
	square block matrices of the input
(4)	into a long block column matrix
$\operatorname{unvec}\left(\mathbf{A}\right)$	Reshapes a column vector into a ma-
	trix
$cof(\mathbf{A})$	Cofactor matrix of A
$\operatorname{eig}\left(\mathbf{A}\right)$	Set of the eigenvalues of <b>A</b>
$[\![\mathbf{A},\mathbf{B},\mathbf{C},\ldots]\!]$	CANDECOMP/PARAFAC (CP) de-
	composition of the tensor ${\cal X}$ from the
	outer product of column vectors of <b>A</b> ,
	B, C,
$[\![\boldsymbol{\lambda};\mathbf{A},\mathbf{B},\mathbf{C},\ldots]\!]$	Normalized CANDE-
	COMP/PARAFAC (CP) decom-
	position of the tensor $\mathcal{X}$ from the
	outer product of column vectors of
	$\mathbf{A}, \mathbf{B}, \mathbf{C}, \dots$
$N(\mathbf{A})$ , $nullspace(\mathbf{A})$ , $kernel(\mathbf{A})$	Nullspace (or kernel)
$C(\mathbf{A})$ , columnspace( $\mathbf{A}$ ), range( $\mathbf{A}$ )	Columnspace (or range), i.e., the
	space span $(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n)$ , where $\mathbf{a}_i$ is
	the ith column vector of the matrix
	A
$\mathrm{span}\left(\mathbf{a}_{1},\mathbf{a}_{2},\ldots,\mathbf{a}_{n}\right)$	Vector space spanned by the argu-
	ment vectors
$\operatorname{span}\left(\mathbf{A}\right)$	Vector space spanned by the col-
	umn vectors of <b>A</b> , which gives the
1 (4)	columnspace of A
$\operatorname{rank}\left(\mathbf{A}\right)$	Rank, that is, $\dim(\operatorname{span}(\mathbf{A})) =$
74.	$\dim (C(\mathbf{A}))$
nullity (A)	Nullity of $\mathbf{A}$ , i.e., dim $(N(\mathbf{A}))$
$\mathrm{tr}\left(\mathbf{A}\right)$	trace
$\mathbf{a} \perp \mathbf{b}$	a is orthogonal to b
a ≠ b	a is not orthogonal to b
$\langle \mathbf{a}, \mathbf{b} \rangle$	Inner product, i.e., $\mathbf{a}^{T}\mathbf{b}$
$\mathbf{a} \circ \mathbf{b}$	Outer product, i.e., $\mathbf{a}\mathbf{b}^{T}$
⊗ -	Kronecker product
⊙	Hadamard (elementwise) product
<b>♦</b>	Khatri-Rao product
$\otimes$	Kronecker Product

<b>×</b>	<i>n</i> -mode product
$egin{array}{c} X_{(n)} \end{array}$	<i>n</i> -mode product <i>n</i> -mode matricization of the tensor $\mathcal{X}$
$\mathcal{X}^{(n)} \leq 0$	Nonnegative tensor
$\mathbf{a} \leq \mathbf{b}$	Generalized inequality meaning that
$a \supseteq_K b$	$\mathbf{b} - \mathbf{a}$ belongs to the conic subset $K$ in
	the space $\mathbb{R}^n$
$\mathbf{a} \prec_K \mathbf{b}$	Strict generalized inequality meaning
$\mathbf{a} \prec_K \mathbf{b}$	that $\mathbf{b} - \mathbf{a}$ belongs to the interior of
	that $\mathbf{b} - \mathbf{a}$ belongs to the interior of the conic subset $K$ in the space $\mathbb{R}^n$
$\mathbf{a} \leq \mathbf{b}$	
$a \leq b$	Generalized inequality meaning that $\mathbf{b} - \mathbf{a}$ belongs to the nonnegative or-
	thant conic subset, $\mathbb{R}^n_+$ , in the space $\mathbb{R}^n$
$\mathbf{a} \prec \mathbf{b}$	Strict generalized inequality meaning
a < b	that $\mathbf{b} - \mathbf{a}$ belongs to the positive or-
	that $\mathbf{b} = \mathbf{a}$ belongs to the positive of- thant conic subset, $\mathbb{R}_{++}^n$ , in the space
	$\mathbb{R}^n$
$\mathbf{A} \leq_K \mathbf{B}$	Generalized inequality meaning that
$A \supseteq_K B$	$\mathbf{B} - \mathbf{A}$ belongs to the conic subset $K$
	in the space $\mathbb{S}^n$
$\mathbf{A} \prec_K \mathbf{B}$	Strict generalized inequality meaning
A \K B	that $\mathbf{B} - \mathbf{A}$ belongs to the interior of
	the conic subset $K$ in the space $\mathbb{S}^n$
$A \leq B$	Generalized inequality meaning that
$\mathbf{A} \supseteq \mathbf{B}$	B-A belongs to the positive semidef-
	inite conic subset, $\mathbb{S}_{+}^{n}$ , in the space $\mathbb{S}^{n}$
A < B	Strict generalized inequality meaning
	that $\mathbf{B} - \mathbf{A}$ belongs to the positive or-
	that $\mathbf{b} - \mathbf{k}$ belongs to the positive of than conic subset, $\mathbb{S}_{++}^n$ , in the space
	$\mathbb{S}^n$
	$\sim$

#### 3.1 Indexing

$x_{i_1,i_2,\ldots,i_N}$	Element in the position
	$(i_1,i_2,\ldots,i_N)$ of the tensor $\boldsymbol{\mathcal{X}}$
$\mathcal{X}^{(n)}$	nth tensor in a nontemporal sequence
$[\mathcal{X}]_{i_1,i_2,,i_N}$	Element $x_{i_1,i_2,,i_N}$
$\mathbf{x}_n, \mathbf{x}_{:n}$	nth column of the matrix $X$
$\mathbf{x}_{n}$ :	nth row of the matrix $X$
$\mathbf{x}_{i_1,,i_{n-1},:,i_{n+1},,i_N}$	Mode- $n$ fiber of the tensor $\boldsymbol{\mathcal{X}}$
$\mathbf{X}_{:,i_{2},i_{3}}$	Column fiber (mode-1 fiber) of the
	thrid-order tensor $\boldsymbol{\mathcal{X}}$
$\mathbf{x}_{i_1,:,i_3}$	Row fiber (mode-2 fiber) of the thrid-
2 0	order tensor $\mathcal{X}$

$X_{i_1,i_2,:}$	Tube fiber (mode-3 fiber) of the
	thrid-order tensor $\mathcal{X}$
$\mathbf{X}_{i_1,:,:}$	Horizontal slice of the thrid-order
	tensor ${\cal X}$
$X_{:,i_2,:}$	Lateral slices slice of the thrid-order
	tensor ${\cal X}$
$\mathbf{X}_{i_3}, \mathbf{X}_{:,:,i_3}$	Frontal slices slice of the thrid-order
	tensor $\mathcal{X}$

### 4 Sets

A + B	Set addition (Minkowski sum)
A - B	Minkowski difference
$A \setminus B, A - B$	Set difference or set subtraction,
	i.e., the set containing the elements
	of $A$ that are not in $B$
$A \cup B$	Set of union
$A \cap B$	Set of intersection
$A \times B$	Cartesian product
$A^{\perp}$	Orthogonal complement of $A$ , e.g.,
	$N(\mathbf{A}) = C(\mathbf{A}^{T})^{\perp}$
$A \oplus B$	Direct sum, e.g., $C(\mathbf{A}^{\top}) \oplus C(\mathbf{A}^{\top})^{\perp} =$
	$\mathbb{R}^n$
$A^c, \bar{A}$	Complement set
#A,  A	Cardinality
$a \in A$	a is element of A
$a \notin A$	a is not element of A
$\{1,2,\ldots,n\}$	Discrete set containing the integer el-
	ements $1, 2, \ldots, n$
$\mathbb{R}$	Set of real numbers
$\mathbb{C}$	Set of complex numbers
$\mathbb Z$	Set of integer number
$\mathbb{B} = \{0, 1\}$	Boolean set
Ø	Empty set
$\mathbb{N}$	Set of natural numbers
$\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$	???
$\mathbb{K}^{I_1  imes I_2  imes \cdots  imes I_N}$	$I_1 \times I_2 \times \cdots \times I_N$ -dimensional real (or
	complex) space
$\mathbb{K}_{+}$	Nonnegative real (or complex) space
$\mathbb{K}_{++}$	Positive real (or complex) space, i.e.,
	$\mathbb{K}_{++} = \mathbb{K}_+ \setminus \{0\}$
$\mathbb{S}^n,\mathcal{S}^n$	Conic set of the symmetric matrices
	in $\mathbb{R}^{n \times n}$

 $\mathbb{S}^n_+, \mathcal{S}^n_+$ Conic set of the symmetric positive semidefinite matrices in  $\mathbb{R}^{n\times n}$ Conic set of the symmetric positive  $\mathbb{S}_{++}^n, \mathcal{S}_{++}^n$ definite matrices in  $\mathbb{R}^{n\times n}$ , i.e.,  $\mathbb{S}^n_{++}=$  $\mathbb{S}^n_+ \setminus \{\mathbf{0}\}$ Set of all hermitian matrices in  $\mathbb{C}^{n\times n}$  $\mathbb{H}^n$ [a,b]Closed interval of a real set from a to (a,b)Opened interval of a real set from a [a, b), (a, b]Half-opened intervals of a real set from a to b

#### 5 Signals and functions operations and indexing

 $f:A\to B$ A function f whose domain is A and codomain is  $\boldsymbol{B}$  $f^{(n)}$ nth derivative of the function f $f^{-1}$ Inverse function of f $f \circ g$ Composition of the functions f and  $\inf_{\mathbf{y}\in\mathcal{A}}g(\mathbf{x},\mathbf{y})$ Infimum  $\sup_{\mathbf{y}\in\mathcal{A}}g(\mathbf{x},\mathbf{y})$ Supremum Convolution (N) Circular convolution Continuous-time tx(t) $x[n],x[k],x[m],x[i],\ldots$ Discrete-time  $n, k, m, i, \ldots$ Discrete-time  $n, k, m, i, \ldots$  (it should  $x(n), x(k), x(m), x(i), \dots$ be used only if there are no continuous-time signals in the context to avoid ambiguity) Estimate of x(t) or x[n]; the Hilbert  $\hat{x}(t)$  or  $\hat{x}[n]$ transform of x(t) or x[n] $\tilde{x}[n]$ Periodic discrete-time signal  $x\left[\left((n-m)\right)_{N}\right], x\left((n-m)\right)_{N}$ Circular shift in m samples within a N-samples window  $x_I(t)$  or  $x_I[n]$ Real or in-phase part of x(t) or x[n]Imaginary or quadrature part of x(t) $x_Q(t)$  or  $x_Q[n]$ or x[n] $\mathcal{F}\left\{ \cdot \right\}$ Fourier transform  $\mathcal{L}\left\{ \cdot \right\}$ Laplace transform  $\mathcal{Z}\left\{ \cdot \right\}$ z-transform X(s)Laplace transform of x(t)

X(f)	Fourier transform (FT) (in linear fre-
	quency, Hz) of $x(t)$
$X(j\omega)$	Fourier transform (FT) (in angular
	frequency, rad/sec) of $x(t)$
$X(e^{j\omega})$	Discrete-time Fourier transform
	(DTFT) of $x[n]$
X[k], X(k)	Discrete Fourier transform (DFT) or
	fast Fourier transform (FFT) of $x[n]$ ,
	or even the Fourier series (FS) of the
	periodic signal $x(t)$
$\tilde{X}[k], \tilde{X}(k)$	Discrete Fourier series (DFS) of $\tilde{x}[n]$
X(z)	z-transform of $x[n]$
$S_x(f)$	Power spectral density of $x(t)$ in lin-
	ear frequency
$S_x(j\omega)$	Power spectral density of $x(t)$ in an-
	gular frequency

## 6 Probability and stochastic processes

$E\left[\cdot ight]$	Statistical expectation
$E_u\left[\cdot\right]$	Statistical expectation with respect
	to u
var(x)	Variance of the random variable $x$
$\operatorname{erfc}(\cdot)$	Complementary error function
P(A)	Probability of the event or set $A$
$p(\cdot)$	Probability density function
$p(x \mid A)$	Conditional probability density func-
	tion
$a \sim P$	Random variable $a$ with distribution
	P
$\mathcal{N}(\mu,\sigma^2)$ $\mathcal{C}\mathcal{N}(\mu,\sigma^2)$	Gaussian distribution of a random
	variable with mean $\mu$ and variance $\sigma^2$
$\mathcal{CN}(\mu,\sigma^2)$	Complex Gaussian distribution of a
	random variable with mean $\mu$ and
	variance $\sigma^2$
$\mathcal{N}(oldsymbol{\mu}, oldsymbol{\Sigma})$	Gaussian distribution of a vector ran-
	dom variable with mean $\mu$ and co-
	variance matrix $\Sigma$
$\mathcal{CN}(oldsymbol{\mu},oldsymbol{\Sigma})$	Complex Gaussian distribution of a
	vector random variable with mean $\mu$
	and covariance matrix $\Sigma$
$\mathcal{U}(a,b)$	Uniform distribution from $a$ to $b$

#### 7 General notations

 $a \wedge b$ Logical AND of a and b $a \lor b$ Logical OR of a and bLogical negation of a $\neg a$ Э There exists ∄ There does not exist ∃! There exist an unique Α For all Such that Therefore Logical equivalence ≜ Equal by definition # Not equal Infinity  $\infty$ Absolute value of a|a|Base-10 logarithm or decimal logalog rithm lnNatual logarithm  $\text{Re}\left\{x\right\}$ Real part of x $\operatorname{Im}\left\{ x\right\}$ Imaginary part of x $\lceil \cdot \rceil$ Ceiling operation  $\lfloor \cdot \rfloor$ Floor operation ∠. phase (complex argument)  $x \mod y$ Remainder, i.e., x - y|x/y| ${\rm frac}\,(x)$ Fractional part, i.e.,  $x \mod 1$ 

#### 8 Abbreviations

 $\begin{array}{ccc} \text{wrt.} & \text{With respect to} \\ \text{st.} & \text{Subject to} \\ \text{iff.} & \text{If and only if} \\ \text{EVD} & \text{Eigenvalue decomposition, or eigendecomposition} \\ \text{SVD} & \text{Singular value decomposition} \\ \text{CP} & \text{CANDECOMP/PARAFAC} \end{array}$