

Convex sets		Comments
Convex hull:	Set <ul style="list-style-type: none">$\text{conv } C = \left\{\sum_{i=1}^k \theta_i \mathbf{x}_i \mid \mathbf{x}_i \in C, 0 \leq \theta_i \leq 1, \mathbf{1}^\top \boldsymbol{\theta} = 1\right\}$	<ul style="list-style-type: none">conv C will be the smallest convex set that contains C.conv C will be a finite set as long as C is also finite.
Affine hull:	<ul style="list-style-type: none">aff $C = \left\{\sum_{i=1}^k \theta_i \mathbf{x}_i \mid \mathbf{x}_i \in C \text{ for } i = 1, \cdots, \mathbf{1}^\top \boldsymbol{\theta} = 1\right\}$	<ul style="list-style-type: none">A will be the smallest affine set that contains C.Different from the convex set, θ_i is not restricted between 0 and 1aff C will always be an infinite set. If aff C contains the origin, it is also a subspace.
Conic hull:	<ul style="list-style-type: none">$A = \left\{\sum_{i=1}^k \theta_i \mathbf{x}_i \mid \mathbf{x}_i \in C, \theta_i > 0 \text{ for } i = 1, \cdots, k\right\}$	<ul style="list-style-type: none">A will be the smallest convex conic that contains C.Different from the convex and affine sets, θ_i does not need to sum up 1.
Ray:	<ul style="list-style-type: none">$\mathcal{R} = \{\mathbf{x}_0 + \theta \mathbf{v} \mid \theta \geq 0\}$	<ul style="list-style-type: none">The ray is an infinite set that begins in \mathbf{x}_0 and extends infinitely in direction of \mathbf{v}. In other words, it has a beginning, but it has no end.
Hyperplane:	<ul style="list-style-type: none">$\mathcal{H} = \{\mathbf{x} \mid \mathbf{a}^\top \mathbf{x} = b\}$$\mathcal{H} = \{\mathbf{x} \mid \mathbf{a}^\top (\mathbf{x} - \mathbf{x}_0) = \mathbf{0}\}$$\mathcal{H} = \mathbf{x}_0 + a^\perp$	<ul style="list-style-type: none">It is an infinite set $\mathbb{R}^{n-1} \subset \mathbb{R}^n$ that divides the space into two halfspaces.$a^\perp = \{\mathbf{v} \mid \mathbf{a}^\top \mathbf{v} = 0\}$ is the set of vectors perpendicular to \mathbf{a}. It passes through the origin.a^\perp is offset from the origin by \mathbf{x}_0, which is any vector in \mathcal{H}.
Halfspaces:	<ul style="list-style-type: none">$\mathcal{H}_\leq = \{\mathbf{x} \mid \mathbf{a}^\top \mathbf{x} \leq b\}$$\mathcal{H}_\geq = \{\mathbf{x} \mid \mathbf{a}^\top \mathbf{x} \geq b\}$	<ul style="list-style-type: none">They are infinite sets of the parts divided by \mathcal{H}.
Euclidean ball:	<ul style="list-style-type: none">$B(\mathbf{x}_c, r) = \{\mathbf{x} \mid \ \mathbf{x} - \mathbf{x}_c\ _2 \leq r\}$$B(\mathbf{x}_c, r) = \{\mathbf{x} \mid (\mathbf{x} - \mathbf{x}_c)^\top (\mathbf{x} - \mathbf{x}_c) \leq r^2\}$$B(\mathbf{x}_c, r) = \{\mathbf{x}_c + r \ \mathbf{u}\ \mid \ \mathbf{u}\ \leq 1\}$	<ul style="list-style-type: none">$B(\mathbf{x}_c, r)$ is a finite set as long as $r < \infty$.\mathbf{x}_c is the center of the ball.r is its radius.
Ellipsoid:	<ul style="list-style-type: none">$\mathcal{E} = \{\mathbf{x} \mid (\mathbf{x} - \mathbf{x}_c)^\top \mathbf{P}^{-1} (\mathbf{x} - \mathbf{x}_c) \leq 1\}$$\mathcal{E} = \{\mathbf{x}_c + \mathbf{A} \mathbf{u} \mid \ \mathbf{u}\ \leq 1\}$, where $\mathbf{A} = \mathbf{P}^{1/2}$.	<ul style="list-style-type: none">\mathcal{E} is a finite set as long as \mathbf{P} is a finite matrix.\mathbf{P} is symmetric and positive definite, that is, $\mathbf{P} = \mathbf{P}^\top > \mathbf{0}$.$\mathbf{x}_c$ is the center of the ellipsoid.The lengths of the semi-axes are given by $\sqrt{\lambda_i}$.\mathbf{A} is invertible. When it is not, we say that \mathcal{E} is a degenerated ellipsoid (degenerated ellipsoids are also convex).
Norm cone:	<ul style="list-style-type: none">$C = \{[x_1, x_2, \cdots, x_n, t]^\top \mid \mathbf{x} \in \mathbb{R}^n, \ \mathbf{x}\ _p \leq t\} \subseteq \mathbb{R}^{n+1}$	<ul style="list-style-type: none">Although it is named “Norm cone”, it is a set, not a scalar.The cone norm increases the dimension of \mathbf{x} in 1.For $p = 2$, it is called the second-order cone, quadratic cone, Lorentz cone or ice-cream cone.
Proper cone: $K \subset \mathbb{R}^n$ is a proper cone when it has the following properties	<ul style="list-style-type: none">K is a convex cone, i.e., $\alpha K \equiv K, \alpha > 0$.$K$ is closed.K is solid.K is pointed, i.e., $-K \cap K = \{\mathbf{0}\}$.	<ul style="list-style-type: none">The proper cone K is used to define the <i>generalized inequality</i> (or <i>partial ordering</i>) in some set S. For the generalized inequality, one must define both the proper cone K and the set S.$\mathbf{x} \leq \mathbf{y} \iff \mathbf{y} - \mathbf{x} \in K$ for $\mathbf{x}, \mathbf{y} \in S$ (generalized inequality)$\mathbf{x} < \mathbf{y} \iff \mathbf{y} - \mathbf{x} \in \text{int } K$ for $\mathbf{x}, \mathbf{y} \in S$ (strict generalized inequality).There are two cases where K and S are understood from context and the subscript K is dropped out:<ul style="list-style-type: none">When $S = \mathbb{R}^n$ and $K = \mathbb{R}_+^n$ (the nonnegative orthant). In this case, $\mathbf{x} \leq \mathbf{y}$ means that $x_i \leq y_i$.When $S = S^n$ and $K = S_+^n$ or $K = S_{++}^n$, where S^n denotes the set of symmetric $n \times n$ matrices, S_+^n is the space of the positive semidefinite matrices, and S_{++}^n is the space of the positive definite matrices. S_+^n is a proper cone in S^n (?). In this case, the generalized inequality $\mathbf{Y} \succeq \mathbf{X}$ means that $\mathbf{Y} - \mathbf{X}$ is a positive semidefinite matrix belonging to the positive semidefinite cone S_+^n in the subspace of symmetric matrices S^n. It is usual to denote $\mathbf{X} > \mathbf{0}$ and $\mathbf{X} \geq \mathbf{0}$ to mean that \mathbf{X} is a positive definite and semidefinite matrix, respectively, where $\mathbf{0} \in \mathbb{R}^{n \times n}$ is a zero matrix.Another common usage is when $S = \mathbb{R}^n$ and $K = \{\mathbf{e} \in \mathbb{R}^n \mid c_1 + c_2 t + \cdots + c_n t^{m-1} \geq 0, \text{ for } 0 \leq t \leq 1\}$. In this case, $\mathbf{x} \leq_K \mathbf{y}$ means that $x_1 + x_2 t + \cdots + x_n t^{m-1} \leq y_1 + y_2 t + \cdots + y_n t^{m-1}$.The generalized inequality has the following properties:<ul style="list-style-type: none">If $\mathbf{x} \leq_K \mathbf{y}$ and $\mathbf{u} \leq_K \mathbf{v}$, then $\mathbf{x} + \mathbf{u} \leq_K \mathbf{y} + \mathbf{v}$ (preserve under addition).If $\mathbf{x} \leq_K \mathbf{y}$ and $\mathbf{y} \leq_K \mathbf{z}$, then $\mathbf{x} \leq_K \mathbf{z}$ (transitivity).If $\mathbf{x} \leq_K \mathbf{y}$, then $\alpha \mathbf{x} \leq_K \mathbf{y}$ for $\alpha \geq 0$ (preserve under nonnegative scaling).$\mathbf{x} \leq_K \mathbf{x}$ (reflexivity).If $\mathbf{x} \leq_K \mathbf{y}$ and $\mathbf{y} \leq_K \mathbf{x}$, then $\mathbf{x} = \mathbf{y}$ (antisymmetric).If $\mathbf{x}_i \leq_K \mathbf{y}_i$, for $i = 1, 2, \dots$, and $\mathbf{x}_i \rightarrow \mathbf{x}$ and $\mathbf{y}_i \rightarrow \mathbf{y}$ as $i \rightarrow \infty$, then $\mathbf{x} \leq_K \mathbf{y}$.It is called partial ordering because $\mathbf{x} \not\leq_K \mathbf{y}$ and $\mathbf{y} \not\leq_K \mathbf{x}$ for many $\mathbf{x}, \mathbf{y} \in S$. When it happens, we say that \mathbf{x} and \mathbf{y} are not comparable (this case does not happen in ordinary inequality, $<$ and $>$).$\mathbf{x} \in S$ is the <i>minimum</i> element of S with respect to the convex cone K if $\mathbf{x} \leq_K \mathbf{y}, \forall \mathbf{y} \in S$ (for <i>maximum</i>, $\mathbf{x} \succeq_K \mathbf{y}, \forall \mathbf{y} \in S$). It means that $S \subseteq \mathbf{x} + K$ (for the maximum, $S \subseteq \mathbf{x} - K$), where $\mathbf{x} + K$ denotes the set K shifted from the origin by \mathbf{x}. Note that any point in $\mathbf{x} + K$ is greater or equal to \mathbf{x} in the generalized inequality mean. The set S does not necessarily have a minimum (maximum), but the minimum (maximum) is unique if it does.$\mathbf{x} \in S$ is the <i>minimal</i> element of S if $\mathbf{y} \leq_K \mathbf{x}$ only when $\mathbf{y} = \mathbf{x}$. The same is true for <i>maximal</i>. We can have many different minimal (maximal) elements. The mathematical notation for that is $(\mathbf{x} - K) \cap S = \{\mathbf{x}\}$, where $\mathbf{x} - K$ denotes all points that are comparable to \mathbf{x} and less than or equal to \mathbf{x} (for the maximal, we have $(\mathbf{x} + K) \cap S = \{\mathbf{x}\}$).When $K = \mathbb{R}_+$ and $S = \mathbb{R}$ (ordinary inequality), the minimal is equal to the minimum and the maximal is equal to the maximum.When we say that a scalar-valued function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is nondecreasing, it means that whenever $\mathbf{u} \leq \mathbf{v}$, we have $\bar{h}(\mathbf{u}) \leq \bar{h}(\mathbf{v})$. Similar results hold for decreasing, increasing, and nonincreasing scalar functions.
Dual cone:	<ul style="list-style-type: none">$K^* = \{\mathbf{y} \mid \mathbf{x}^\top \mathbf{y} \geq 0, \forall \mathbf{x} \in K\}$	<ul style="list-style-type: none">K^* is a cone, and it is convex even when the original cone K is nonconvex.K^* has the following properties:<ul style="list-style-type: none">K^* is closed and convex.$K_1 \subseteq K_2$ implies $K_1^* \supseteq K_2^*$.If K has a nonempty interior, then K^* is pointed.If the closure of K is pointed then K^* has a nonempty interior.K^{**} is the closure of the convex hull of K. Hence, if K is convex and closed, $K^{**} = K$.
Polyhedra:	<ul style="list-style-type: none">$\mathcal{P} = \left\{\mathbf{x} \mid \mathbf{a}_j^\top \mathbf{x} \leq b_j, j = 1, \dots, m, \mathbf{a}_j^\top \mathbf{x} = d_j, j = 1, \dots, p\right\}$$\mathcal{P} = \{\mathbf{x} \mid \mathbf{A} \mathbf{x} \leq \mathbf{b}, \mathbf{C} \mathbf{x} = \mathbf{d}\}$, where $\mathbf{A} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \dots & \mathbf{a}_m \end{bmatrix}^\top$ and $\mathbf{C} = \begin{bmatrix} \mathbf{c}_1 & \mathbf{c}_2 & \dots & \mathbf{c}_m \end{bmatrix}^\top$	<ul style="list-style-type: none">The polyhedron may or may not be an infinite set.Polyhedron is the result of the intersection of m halfspaces and p hyperplanes.Subspaces, hyperplanes, lines, rays line segments, and halfspaces are all polyhedra.The <i>nonnegative orthant</i>, $\mathbb{R}_+^n = \{\mathbf{x} \in \mathbb{R}^n \mid x_i \geq 0 \text{ for } i = 1, \dots, n\} = \{\mathbf{x} \in \mathbb{R}^n \mid \mathbf{I} \mathbf{x} \geq \mathbf{0}\}$, is a special polyhedron.
Simplex:	<ul style="list-style-type: none">$\mathcal{S} = \text{conv} \{ \mathbf{v}_m \}_{m=0}^k = \left\{\sum_{i=0}^k \theta_i \mathbf{v}_i \mid \mathbf{0} \leq \theta_i \leq 1, \mathbf{1}^\top \boldsymbol{\theta} = 1\right\}$$\mathcal{S} = \{\mathbf{x} \mid \mathbf{x} = \mathbf{v}_0 + \mathbf{V} \boldsymbol{\theta}\}$, where $\mathbf{V} = \begin{bmatrix} \mathbf{v}_1 - \mathbf{v}_0 & \dots & \mathbf{v}_n - \mathbf{v}_0 \end{bmatrix} \in \mathbb{R}^{n \times k}$$\mathcal{S} = \{\mathbf{x} \mid \underbrace{\mathbf{A}_1 \mathbf{x} \leq \mathbf{A}_1 \mathbf{v}_0, \mathbf{1}^\top \mathbf{A}_1 \mathbf{x} \leq 1 + \mathbf{1}^\top \mathbf{A}_1 \mathbf{v}_0}_{\text{Linear inequalities in } \mathbf{x}}, \underbrace{\mathbf{A}_2 \mathbf{x} = \mathbf{A}_2 \mathbf{v}_0}_{\text{Linear equalities in } \mathbf{x}}\}$ (Polyhedra form), where $\mathbf{A} = \begin{bmatrix} \mathbf{A}_1 \\ \mathbf{A}_2 \end{bmatrix}$ and $\mathbf{A} \mathbf{V} = \begin{bmatrix} \mathbf{I}_{k \times k} \\ \mathbf{0}_{n-k \times n-k} \end{bmatrix}$	<ul style="list-style-type: none">Simplexes are a subfamily of the polyhedra set.Also called k-dimensional Simplex in \mathbb{R}^n.The set $\{\mathbf{v}_m\}_{m=0}^k$ is an affinely independent, which means $\{\mathbf{v}_1 - \mathbf{v}_0, \dots, \mathbf{v}_k - \mathbf{v}_0\}$ are linearly independent.$\mathbf{V} \in \mathbb{R}^{n \times k}$ is a full-rank tall matrix, i.e., $\text{rank}(\mathbf{V}) = k$. All its column vectors are independent. The matrix \mathbf{A} is its left pseudoinverse.
α -sublevel set:	<ul style="list-style-type: none">$C_\alpha = \{\mathbf{x} \in \text{dom}(f) \mid f(\mathbf{x}) \leq \alpha\}$ (regarding convexity)$C_\alpha = \{\mathbf{x} \in \text{dom}(f) \mid f(\mathbf{x}) \geq \alpha\}$ (regarding concavity)	<ul style="list-style-type: none">If f is a convex function, then sublevel sets of f are convexes for any $\alpha \in \mathbb{R}$.The converse is not true: a function can have all its sublevel set convex and not be a convex function.
Functions (or operators) and their implications regarding convexity		
Function	Convex?	Comments
Union: $C = A \cup B$	Not in most of the cases.	
Intersection: $C = A \cap B$	Yes, if A and B are convex sets.	
Convex function: $f : \text{dom}(f) \rightarrow \mathbb{R}$	Yes.	<ul style="list-style-type: none">Graphically, the line segment between $(\mathbf{x}, f(\mathbf{x}))$ and $(\mathbf{y}, f(\mathbf{y}))$ lies always above the graph f.In terms of sets, a function is convex iff a line segment within $\text{dom}(f)$, which is a convex set, gives an image set that is also convex.$\text{dom } f$ is convex iff all points for any line segment within $\text{dom}(f)$ belong to it.<i>First-order condition</i>: f is convex iff $\text{dom}(f)$ is convex and $f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}), \forall \mathbf{x}, \mathbf{y} \in \text{dom}(f), \mathbf{x} \neq \mathbf{y}$, being $\nabla f(\mathbf{x})$ the gradient vector. This inequation says that the first-order Taylor approximation is a <i>underestimator</i> for convex functions. The first-order condition requires that f is differentiable.If $\nabla f(\mathbf{x}) = \mathbf{0}$, then $f(\mathbf{y}) \geq f(\mathbf{x}), \forall \mathbf{y} \in \text{dom}(f)$ and \mathbf{x} is a global minimum.<i>Second-order condition</i>: f is convex iff $\text{dom}(f)$ is convex and $\mathbf{H} \succeq \mathbf{0}$, that is, the Hessian matrix \mathbf{H} is a positive semidefinite matrix. It means that the graphic of the curvature has a positive (upward) curvature at \mathbf{x}. It is important to note that, if $\mathbf{H} > \mathbf{0}, \forall \mathbf{x} \in \text{dom}(f)$, then f is strictly convex. But f is strictly convex, not necessarily that $\mathbf{H} > \mathbf{0}, \forall \mathbf{x} \in \text{dom}(f)$. Therefore, strict convexity can only be partially characterized.
Affine function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$	Yes, if the domain $S \subseteq \mathbb{R}^n$ is a convex set, then its image $f(S) = \{f(\mathbf{x}) \mid \mathbf{x} \in S\} \subseteq \mathbb{R}^m$ is also convex.	<ul style="list-style-type: none">The affine function, $f(\mathbf{x}) = \mathbf{A} \mathbf{x} + \mathbf{b}$, is a broader category that encompasses the linear function, $f(\mathbf{x}) = \mathbf{A} \mathbf{x}$. The linear function has its origin fixed at $\mathbf{0}$ after the transformation, whereas the affine function does not necessarily have it (when not, this makes the affine function nonlinear). Graphically, we can think of an affine function as a linear transformation plus a shift from the origin of \mathbf{b}.A special case of the linear function is when $\mathbf{A} = \mathbf{c}^\top$. In this case, we have $f(\mathbf{x}) = \mathbf{c}^\top \mathbf{x}$, which is the inner product between the vector \mathbf{c} and \mathbf{x}.The inverse image of $C, f^{-1}(C) = \{\mathbf{x} \mid f(\mathbf{x}) \in C\}$, is also convex.The <i>linear matrix inequality</i> (LMI), $\mathbf{A}(\mathbf{x}) = x_1 \mathbf{A}_1 + \cdots + x_n \mathbf{A}_n \leq \mathbf{B}$, is a special case of affine function. In other words, $f(S) = \{\mathbf{x} \mid \mathbf{A}(\mathbf{x}) \leq \mathbf{B}\}$ is a convex set if S is convex. Many optimization problems can be formulated as LMI problems and solved optimally.
Exponential function $f : \mathbb{R} \rightarrow \mathbb{R}$	Yes.	
Quadratic function $f : \mathbb{R}^n \rightarrow \mathbb{R}$	It depends on the matrix \mathbf{P} :	<ul style="list-style-type: none">f is convex iff $\mathbf{P} \succeq \mathbf{0}$.$f$ is strictly convex iff $\mathbf{P} > \mathbf{0}$.f is concave iff $\mathbf{P} \preceq \mathbf{0}$.$f$ is strictly concave iff $\mathbf{P} < \mathbf{0}$.
Power function $f : \mathbb{R}_{++} \rightarrow \mathbb{R}$	It depends on a	<ul style="list-style-type: none">f is convex iff $a \geq 1$ or $a \leq 0$.f is concave iff $0 \leq a \leq 1$.
Power of absolute value: $f : \mathbb{R} \rightarrow \mathbb{R}$	Yes.	
Logarithm function: $f : \mathbb{R}_{++} \rightarrow \mathbb{R}$	Yes.	
Negative entropy function: $f : \mathbb{R}_+ \rightarrow \mathbb{R}$	Yes	<ul style="list-style-type: none">When it is defined $f(x) _{x=0} = 0$, $\text{dom}(f) = \mathbb{R}$.
Minkowski distance, p -norm function, or l_p norm function: $f : \mathbb{R}^n \rightarrow \mathbb{R}$	Yes.	<ul style="list-style-type: none">It can be proved by triangular inequality.
Maximum element: $f : \mathbb{R}^n \rightarrow \mathbb{R}$	Yes.	
Maximum function: $f : \mathbb{R}^n \rightarrow \mathbb{R}$	Yes, if f_1, \dots, f_n are convex function.	
Minimum function: $f : \mathbb{R}^n \rightarrow \mathbb{R}$	Not in most of the cases.	
Log-sum-exp function: $f : \mathbb{R}^n \rightarrow \mathbb{R}$	Yes.	<ul style="list-style-type: none">This function is interpreted as the approximation of the maximum element function, since $\max\{x_1, \dots, x_n\} \leq f(\mathbf{x}) \leq \max\{x_1, \dots, x_n\} + \log n$
Geometric mean function $f : \mathbb{R}^n \rightarrow \mathbb{R}$	Yes	
Log-determinant function $f : \mathcal{S}_{++}^n \rightarrow \mathbb{R}$	Yes	<ul style="list-style-type: none">\mathbf{X} is positive semidefinite, i.e., $\mathbf{X} > \mathbf{0} : \mathbf{X} \in \mathcal{S}_{++}^n$.
Composite function $f = h \circ g : \mathbb{R}^n \rightarrow \mathbb{R}$	<ul style="list-style-type: none">Scalar composition: the following statements hold for $k = 1$ and $n \geq 1$, i.e., $h : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R}^n \rightarrow \mathbb{R}$:<ul style="list-style-type: none">f is convex if h is convex, \tilde{h} is nondecreasing, and g is convex. In this case, $\text{dom}(h)$ is either $(-\infty, a]$ or $(-\infty, a)$.f is convex if h is convex, \tilde{h} is nonincreasing, and g is concave. In this case, $\text{dom}(h)$ is either $[a, \infty)$ or (a, ∞).f is concave if h is concave, \tilde{h} is nondecreasing, and g is concave.f is concave if h is concave, \tilde{h} is nonincreasing, and g is convex.Vector composition: the following statements hold for $k \geq 1$ and $n \geq 1$, i.e., $h : \mathbb{R}^k \rightarrow \mathbb{R}$ and $g : \mathbb{R}^n \rightarrow \mathbb{R}^k$. Hence, $g(\mathbf{x}) = (g_1(\mathbf{x}), g_2(\mathbf{x}), \dots, g_k(\mathbf{x}))$ is a vector-valued function (or simply, vector function), where $g_i : \mathbb{R}^k \rightarrow \mathbb{R}$ for $1 \leq i \leq k$.<ul style="list-style-type: none">f is convex if h is convex, \tilde{h} is nondecreasing in each argument of \mathbf{x}, and $\{g_i\}_{i=1}^k$ is a set of convex functions.f is convex if h is convex, \tilde{h} is nonincreasing in each argument of \mathbf{x}, and $\{g_i\}_{i=1}^k$ is a set of concave functions.f is concave if h is concave, \tilde{h} is nondecreasing in each argument of \mathbf{x}, and $\{g_i\}_{i=1}^k$ is a set of convex functions.f is concave if h is concave, \tilde{h} is nonincreasing in each argument of \mathbf{x}, and $\{g_i\}_{i=1}^k$ is a set of concave functions. Where \tilde{h} is the extended-value extension of the function h , which assigns the value ∞ ($-\infty$) to the point not in $\text{dom}(h)$ for h convex (concave).	<ul style="list-style-type: none">The composition function allows us to see a large class of functions as convex (or concave).For scale composition, the remarkable ones are:<ul style="list-style-type: none">If g is convex then $f(\mathbf{x}) = h(g(\mathbf{x})) = \exp g(\mathbf{x})$ is convex.If g is concave and $\text{dom}(g) \subseteq \mathbb{R}_{++}$, then $f(\mathbf{x}) = h(g(\mathbf{x})) = \log g(\mathbf{x})$ is concave.If g is concave and $\text{dom}(g) \subseteq \mathbb{R}_{++}$, then $f(\mathbf{x}) = h(g(\mathbf{x})) = 1/g(\mathbf{x})$ is convex.If g is convex and $\text{dom}(g) \subseteq \mathbb{R}_+$, then $f(\mathbf{x}) = h(g(\mathbf{x})) = g^p(\mathbf{x})$ is convex, where $p \geq 1$.If g is convex then $f(\mathbf{x}) = -\log(-g(\mathbf{x}))$ is convex, where $\text{dom}(f) = \{\mathbf{x} \mid g(\mathbf{x}) < 0\}$.For vector composition, we have the following examples:<ul style="list-style-type: none">If $g(\mathbf{x}) = \mathbf{A} \mathbf{x} + \mathbf{b}$ is an affine function, then $f = h \circ g$ is convex (concave) if h is convex (concave).Let $h(\mathbf{x}) = x_{[1]} + \cdots + x_{[r]}$ be the sum of the r largest components of $\mathbf{x} \in \mathbb{R}^k$. If g_1, g_2, \dots, g_k are convex, where $\text{dom}(g_i) = \mathbb{R}^n$, then $f = h \circ g$, which is the pointwise sum of the largest g_i's, is convex.$f = h \circ g$ is a convex function when $h(\mathbf{x}) = \log \left(\sum_{i=1}^k x_i^p \right)$ and g_1, g_2, \dots, g_k are convex functions.For $0 < p \leq 1$, the function $h(\mathbf{x}) = \left(\sum_{i=1}^k x_i^p \right)^{1/p}$, where $\text{dom}(h) = \mathbb{R}_+^k$, is concave. If g_1, g_2, \dots, g_k are concaves (convexes) and nonnegatives, then $f = h \circ g$ is concave (convex).
Nonnegative weighted sum: $f : \text{dom}(f) \rightarrow \mathbb{R}$	<ul style="list-style-type: none">If f_1, f_2, \dots, f_n are convex or concave functions, then f is a convex or concave function, respectively.If f_1, f_2, \dots, f_n are strictly convex or concave functions, then f is a strictly convex or concave function, respectively.	<ul style="list-style-type: none">Special cases is when $f = w f$ (a nonnegative scalar factor) and $f = f_1 + f_2$ (sum).
Integral function $f : \mathbb{R}^n \rightarrow \mathbb{R}$:	<ul style="list-style-type: none">If g is convex in \mathbf{x} for each $\mathbf{y} \in \mathcal{A}$ and if $w(\mathbf{y}) \geq 0, \forall \mathbf{y} \in \mathcal{A}$, then f is convex (provided the integral exists).	
Perspective function $f : \mathbb{R}^n \times \mathbb{R}_{++} \rightarrow \mathbb{R}^n$	Yes, if $S \subseteq \text{dom}(f)$ is a convex set, then its image, $f(S) = \{f(\mathbf{x}) \mid \mathbf{x} \in S\} \subseteq \mathbb{R}^n$, is also convex.	<ul style="list-style-type: none">The perspective function decreases the dimension of the function domain since $\text{dim}(\text{dom}(f)) = n + 1$.Its effect is similar to the camera zoom.The inverse image is also convex, that is, if $C \subseteq \mathbb{R}^n$ is convex, then $f^{-1}(C) = \{(\mathbf{x}, t) \in \mathbb{R}^{n+1} \mid \mathbf{x}/t \in C, t > 0\}$ is also convex.A special case is when $n = 1$, which is called <i>quadratic-over-linear function</i>.
Projective (or linear-fractional) function, $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$	Yes, if $S \subseteq \text{dom}(f)$ is a convex set, then its image, $f(S) = \{f(\mathbf{x}) \mid \mathbf{x} \in S\} \subseteq \mathbb{R}^n$, is also convex.	<ul style="list-style-type: none">The linear and affine functions are special cases of the linear-fractional function.$\text{dom}(f) = \{\mathbf{x} \in \mathbb{R}^n \mid \mathbf{c}^\top \mathbf{x} + d > 0\}$$\mathcal{P}(\mathbf{x}) \subset \mathbb{R}^{n+1}$ is a ray set that begins at the origin and its last component takes only positive values. For each $\mathbf{x} \in \text{dom}(f)$, it is associated a ray set in \mathbb{R}^{n+1} in this form. This (projective) correspondence between all points in $\text{dom}(f)$ and their respective sets \mathcal{P} is a biunivocal mapping.The linear transformation \mathbf{Q} acts on these rays, forming another set of rays.Finally we take the inverse projective transformation to recover $f(\mathbf{x})$.
Epigraph:	<ul style="list-style-type: none">Visually, it is the graph above the $\mathbf{x}, f(\mathbf{x})$ curve.The function f is convex iff its epigraph is convex.	
Hypograph:	<ul style="list-style-type: none">Visually, it is the graph below the $\mathbf{x}, f(\mathbf{x})$ curve.The function f is concave iff its hypograph is convex.	

