


Trial Sequential Analysis of Hazard Ratios

Random-effects model | Diversity $D^2 = 0\%$ | Anticipated HR = 0.80 | $\psi = 0.500$ | $\alpha = 0.05$




Alpha boundaries Naive boundaries Non-binding futility boundaries Z scores

Methods: Random-effects (DerSimonian-Laird) model, allocation $\psi = 0.500$

Alpha spending: O'Brien-Fleming-type (asOF); Non-binding futility: RTSA retrospective inner-wedge algorithm, O'Brien-Fleming-
 $\alpha = 5\%$ (two-sided), power = 80% | Diversity $D^2 = 0\%$, Adjustment factor = 1.00

Trial Sequential Analysis of Hazard Ratios

Random-effects model | Diversity $D^2 = 0\%$ | Anticipated HR = 0.80 | $\psi = 0.500$ | $\alpha = 0.05$




Methods: Random-effects (DerSimonian-Laird) model, allocation $\psi = 0.500$

Alpha spending: O'Brien-Fleming-type (asOF); Non-binding futility: RTSA retrospective inner-wedge algorithm, O'Brien-Fleming-
 $\alpha = 5\%$ (two-sided), power = 80% | Diversity $D^2 = 0\%$, Adjustment factor = 1.00

Trial Sequential Analysis of Hazard Ratios

Random-effects model | Diversity $D^2 = 0\%$ | Anticipated HR = 0.90 | $\psi = 0.533$ | $\alpha = 5\%$




—●— Alpha boundaries —●— Naive boundaries —●— Non-binding futility boundaries —●— Z scores

Methods: Random-effects (DerSimonian-Laird) model, allocation $\psi = 0.533$

Alpha spending: O'Brien-Fleming-type (asOF); Non-binding futility: RTSA retrospective inner-wedge algorithm, O'Brien-Fleming-type
 $\alpha = 5\%$ (two-sided), power = 80% | Diversity $D^2 = 0\%$, Adjustment factor = 1.00

Trial Sequential Analysis of Hazard Ratios

Random-effects model | Diversity $D^2 = 0\%$ | Anticipated HR = 0.80 | $\psi = 0.500$ | $\alpha = 0.05$




Methods: Random-effects (DerSimonian-Laird) model, allocation $\psi = 0.500$

Alpha spending: O'Brien-Fleming-type (asOF); Non-binding futility: RTSA retrospective inner-wedge algorithm, O'Brien-Fleming-
alpha = 5% (two-sided), power = 80% | Diversity $D^2 = 0\%$, Adjustment factor = 1.00

Trial Sequential Analysis of Hazard Ratios

Random-effects model | Diversity $D^2 = 0\%$ | Anticipated HR = 0.80 | $\psi = 0.500$ | $\alpha = 0.05$




Methods: Random-effects (DerSimonian-Laird) model, allocation $\psi = 0.500$

Alpha spending: O'Brien-Fleming-type (asOF); Non-binding futility: RTSA retrospective inner-wedge algorithm, O'Brien-Fleming-
 $\alpha = 5\%$ (two-sided), power = 80% | Diversity $D^2 = 0\%$, Adjustment factor = 1.00

Trial Sequential Analysis of Hazard Ratios

Random-effects model | Diversity $D^2 = 0\%$ | Anticipated HR = 0.80 | $\psi = 0.500$ | $\alpha = 0.05$




Alpha boundaries Naive boundaries Non-binding futility boundaries Z scores

Methods: Random-effects (DerSimonian-Laird) model, allocation $\psi = 0.500$

Alpha spending: O'Brien-Fleming-type (asOF); Non-binding futility: RTSA retrospective
 $\alpha = 5\%$ (two-sided), power = 80% | Diversity $D^2 = 0\%$, Adjustment factor = 1.00

Trial Sequential Analysis of Hazard Ratios

Random-effects model | Diversity $D^2 = 0\%$ | Anticipated HR = 0.80 | $\psi = 0.500$ | $\alpha = 0.05$



Alpha boundaries Naive boundaries Non-binding futility boundaries Z scores