

# CIs and Bernoulli

Tarek El-Hajjaoui

2023-03-01

Suppose that a random sample of eight observations is taken from the normal distribution with unknown mean and unknown variance, and that the observed values are 3.1, 3.5, 2.6, 3.4, 3.8, 3.0, 2.9, and 2.2.

```
x <- c(3.1, 3.5, 2.6, 3.4, 3.8, 3.0, 2.9, 2.2)
x_mu <- mean(x)
x_sd <- sd(x)
n <- 8
```

Find the shortest confidence interval for mean with each of the following three confidence coefficients.

- a) 0.90,

```
q <- qt(1.0 - 0.1 / 2, n - 1)
ci <- x_mu + c(-1, 1) * q * x_sd / sqrt(n)
ci
```

```
## [1] 2.719195 3.405805
```

- b) 0.95,

```
q <- qt(1.0 - 0.05 / 2, n - 1)
ci <- x_mu + c(-1, 1) * q * x_sd / sqrt(n)
ci
```

```
## [1] 2.634021 3.490979
```

- c) 0.99.

```
q <- qt(1.0 - 0.01 / 2, n - 1)
ci <- x_mu + c(-1, 1) * q * x_sd / sqrt(n)
ci
```

```
## [1] 2.42838 3.69662
```

6d

```
p_hat <- 0.66
n <- 2000
ci <- p_hat + c(-1, 1) * qnorm(0.975) * sqrt(p_hat * (1 - p_hat) / n)
ci
```

```
## [1] 0.6392392 0.6807608
```