

MCAT Modeling & Analysis

Tarek El-Hajjaoui

2023-02-25

Data loading & pre-processing

Loading the dataset

```
file_path = '/Users/Tarek/Documents/UCI_MDS_Coding/Stats210P/R_Statistical_Modeling/MCAT/MedGPA.txt'
df = read.table(file_path, header=TRUE, sep=" ", dec=".")
```

Summary of dataset

```
str(df)

## 'data.frame':    55 obs. of  11 variables:
## $ Accept      : chr  "D" "A" "A" "A" ...
## $ Acceptance: int   0 1 1 1 1 1 1 0 1 1 ...
## $ Sex         : chr  "F" "M" "F" "F" ...
## $ BCPM        : num   3.59 3.75 3.24 3.74 3.53 3.59 3.85 3.26 3.74 3.86 ...
## $ GPA         : num   3.62 3.84 3.23 3.69 3.38 3.72 3.89 3.34 3.71 3.89 ...
## $ VR          : int   11 12 9 12 9 10 11 11 8 9 ...
## $ PS          : int   9 13 10 11 11 9 12 11 10 9 ...
## $ WS          : int   9 8 5 7 4 7 6 8 6 6 ...
## $ BS          : int   9 12 9 10 11 10 11 9 11 10 ...
## $ MCAT        : int   38 45 33 40 35 36 40 39 35 34 ...
## $ Apps        : int   5 3 19 5 11 5 5 7 5 11 ...
```

Transforming categorical columns to factor data type.

```
categorical_cols <- c('Accept', 'Acceptance', 'Sex')
df[categorical_cols] <- lapply(df[categorical_cols], as.factor)
```

Ensuring column data types are correct now.

```
str(df)

## 'data.frame':    55 obs. of  11 variables:
## $ Accept      : Factor w/ 2 levels "A","D": 2 1 1 1 1 1 1 2 1 1 ...
## $ Acceptance: Factor w/ 2 levels "0","1": 1 2 2 2 2 2 2 1 2 2 ...
## $ Sex        : Factor w/ 2 levels "F","M": 1 2 1 1 1 2 2 2 1 1 ...
## $ BCPM       : num   3.59 3.75 3.24 3.74 3.53 3.59 3.85 3.26 3.74 3.86 ...
## $ GPA        : num   3.62 3.84 3.23 3.69 3.38 3.72 3.89 3.34 3.71 3.89 ...
## $ VR         : int   11 12 9 12 9 10 11 11 8 9 ...
## $ PS         : int   9 13 10 11 11 9 12 11 10 9 ...
## $ WS         : int   9 8 5 7 4 7 6 8 6 6 ...
## $ BS         : int   9 12 9 10 11 10 11 9 11 10 ...
## $ MCAT       : int   38 45 33 40 35 36 40 39 35 34 ...
## $ Apps       : int   5 3 19 5 11 5 5 7 5 11 ...
```

Splitting up data to supplement analysis in 2-sample t-test

Separating the dataset into Female observations & Male observations

```
df_female <- df[df$Sex == 'F',]  
df_male <- df[df$Sex == 'M',]
```

Female dataset summary

```
str(df_female)  
  
## 'data.frame': 28 obs. of 11 variables:  
## $ Accept : Factor w/ 2 levels "A","D": 2 1 1 1 1 1 1 1 1 1 ...  
## $ Acceptance: Factor w/ 2 levels "0","1": 1 2 2 2 2 2 2 2 2 2 ...  
## $ Sex : Factor w/ 2 levels "F","M": 1 1 1 1 1 1 1 1 1 1 ...  
## $ BCPM : num 3.59 3.24 3.74 3.53 3.74 3.86 4 3.35 3.26 3.71 ...  
## $ GPA : num 3.62 3.23 3.69 3.38 3.71 3.89 3.97 3.49 3.54 3.71 ...  
## $ VR : int 11 9 12 9 8 9 11 11 12 13 ...  
## $ PS : int 9 10 11 11 10 9 9 8 8 10 ...  
## $ WS : int 9 5 7 4 6 6 8 4 8 8 ...  
## $ BS : int 9 9 10 11 11 10 11 8 10 10 ...  
## $ MCAT : int 38 33 40 35 35 34 39 31 38 41 ...  
## $ Apps : int 5 19 5 11 5 11 6 9 6 6 ...
```

Male dataset summary

```
str(df_male)  
  
## 'data.frame': 27 obs. of 11 variables:  
## $ Accept : Factor w/ 2 levels "A","D": 1 1 1 2 1 2 2 2 1 2 ...  
## $ Acceptance: Factor w/ 2 levels "0","1": 2 2 2 1 2 1 1 1 2 1 ...  
## $ Sex : Factor w/ 2 levels "F","M": 2 2 2 2 2 2 2 2 2 2 ...  
## $ BCPM : num 3.75 3.59 3.85 3.26 3.77 3.6 3.29 3.75 3.51 3.27 ...  
## $ GPA : num 3.84 3.72 3.89 3.34 3.77 3.61 3.3 3.65 3.54 3.25 ...  
## $ VR : int 12 10 11 11 8 9 11 8 9 8 ...  
## $ PS : int 13 9 12 11 10 9 8 8 10 9 ...  
## $ WS : int 8 7 6 8 7 4 6 8 9 5 ...  
## $ BS : int 12 10 11 9 10 10 7 11 11 10 ...  
## $ MCAT : int 45 36 40 39 35 32 32 35 39 32 ...  
## $ Apps : int 3 5 5 7 5 8 15 6 1 5 ...
```

2-sample t-test: Y=MCAT scores, split dataset on Sex (male or female)

```
two_sample_t_test <- t.test(df$MCAT[df$Sex=="F"],df$MCAT[df$Sex=="M"], var.equal=TRUE)  
two_sample_t_test
```

```
##  
## Two Sample t-test  
##  
## data: df$MCAT[df$Sex == "F"] and df$MCAT[df$Sex == "M"]  
## t = 0.020173, df = 53, p-value = 0.984  
## alternative hypothesis: true difference in means is not equal to 0  
## 95 percent confidence interval:  
## -2.603913 2.656823  
## sample estimates:  
## mean of x mean of y  
## 36.28571 36.25926
```

Model Creation

Model 1: Y=MCAT scores, X1=GPA, X2=Sex (male or female)

```
model <- lm(MCAT ~ GPA + Sex, data=df)
```

Summary of Model 1

```
summary(model)
```

```
##
## Call:
## lm(formula = MCAT ~ GPA + Sex, data = df)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -11.5825  -2.5260  -0.0993   2.6574   8.4228
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   3.4820     7.0736   0.492   0.625
## GPA           9.1695     1.9652   4.666 2.19e-05 ***
## SexM          0.4261     1.1158   0.382   0.704
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 4.121 on 52 degrees of freedom
## Multiple R-squared:  0.2951, Adjusted R-squared:  0.268
## F-statistic: 10.89 on 2 and 52 DF,  p-value: 0.0001125
```