

a
Where the score separates the groups
VIS
SMN
DAN
SAL
LIM
FPN
DMN
SUB
0.15
0.10
0.05
0.00
−0.05
−0.10
−0.15
contribution per connection, autistic − non-autistic (×10⁻³ logit)
b
Each system pair on its own
DMN–DMN
VIS–DMN
DMN–SUB
VIS–SAL
LIM–DMN
DAN–SAL
FPN–DMN
DAN–DAN
VIS–DAN
SAL–DMN
SAL–SUB
SMN–DMN
VIS–LIM
SMN–SMN
DAN–DMN
SAL–SAL
VIS–VIS
VIS–FPN
SMN–SUB
FPN–SUB
SAL–LIM
VIS–SUB
LIM–LIM
SAL–FPN
SMN–DAN
SMN–SAL
VIS–SMN
SMN–FPN
SUB–SUB
DAN–FPN
FPN–FPN
LIM–FPN
LIM–SUB
DAN–SUB
DAN–LIM
SMN–LIM
familywise p < 0.05
not distinguished
0.50
0.55
0.60
0.65
0.70
held-out AUROC of the pair's contribution
c
Replication across independent halves
black ticks: permutation 95th percentile
correlation
static tangent
windowed tangent
Spearman ρ between independent halves
1.0
0.8
0.6
0.4
0.2
0.0
connections
regions
system pairs
d
Do the ranked pairs matter to the trained model?
held-out AUROC
0.775
0.750
0.725
0.700
0.675
0.650
0.625
0.600
dashed and shaded: random pairs
delete top-ranked
keep only top-ranked
0
5
10
15
20
25
30
35
number of system pairs deleted or kept (of 36)