

Collinearity

- ▶ We often encounter problems in which some predictors are highly correlated, e.g., the seatpos data. In this case, the contribution of a particular predictor could be masked by other predictors, which create difficulties for statistical inference on β .
- ▶ Typical symptoms of collinearity: high pair-wise (sample) correlation between predictors; R^2 is relatively large, overall F test is significant, but none of the predictors is significant.

Collinearity

- ▶ We often encounter problems in which some predictors are highly correlated, e.g., the seatpos data. In this case, the contribution of a particular predictor could be masked by other predictors, which create difficulties for statistical inference on β .
- ▶ Typical symptoms of collinearity: high pair-wise (sample) correlation between predictors; R^2 is relatively large, overall F test is significant, but none of the predictors is significant.
- ▶ What to do with collinearity? Remove some predictors or combine collinear predictors (e.g., PCA).

Collinearity

- ▶ We often encounter problems in which some predictors are highly correlated, e.g., the seatpos data. In this case, the contribution of a particular predictor could be masked by other predictors, which create difficulties for statistical inference on β .
- ▶ Typical symptoms of collinearity: high pair-wise (sample) correlation between predictors; R^2 is relatively large, overall F test is significant, but none of the predictors is significant.
- ▶ What to do with collinearity? Remove some predictors or combine collinear predictors (e.g., PCA).
- ▶ How would collinearity affect prediction of Y ?