

# Patterns of response to 200 mg linvoseltamab in patients with relapsed/refractory multiple myeloma: Longer follow-up of the LINKER-MM1 study

Sundar Jagannath<sup>1</sup>, Joshua Richter<sup>1</sup>, Madhav V. Dhodapkar<sup>2</sup>, James E. Hoffman<sup>3</sup>, Hans C. Lee<sup>4</sup>, Attaya Suvannasankha<sup>5</sup>, Mansi R. Shah<sup>6</sup>, Suzanne Lentzsch<sup>7</sup>, Jeffrey A. Zonder<sup>8</sup>, Rachid Baz<sup>9</sup>, Joseph J. Maly<sup>10</sup>, Swathi Namburi<sup>11</sup>, Ka Lung Wu<sup>12</sup>, Matthew Panko<sup>13</sup>, Jing Christine Ye<sup>13</sup>, Rebecca Silberman<sup>14</sup>, Chang-Ki Min<sup>15</sup>, Marie-Christiane Vekemans<sup>16</sup>, Markus Munder<sup>17</sup>, Ja Min Byun<sup>18</sup>, Joaquín Martínez López<sup>19</sup>, Michelle DeVeaux<sup>20</sup>, Dhruvi Chokshi<sup>20</sup>, Anita Boyapati<sup>20</sup>, Anasuya Hazra<sup>20</sup>, Karen Rodriguez Lorenc<sup>20</sup>, Glenn S. Kroog<sup>20</sup>, Yariv Houvras<sup>20</sup>, Naresh Bumma<sup>21</sup>


<sup>1</sup>Icahn School of Medicine at Mount Sinai, New York, NY, USA; <sup>2</sup>Emory University School of Medicine, Atlanta, GA, USA; <sup>3</sup>University of Miami Health System, Miami, FL, USA; <sup>4</sup>The University of Texas MD Anderson Cancer Centre, Houston, TX, USA; <sup>5</sup>Indiana University Simon Cancer Center and Roudebush VAMC, Indianapolis, IN, USA; <sup>6</sup>Rutgers Cancer Institute of New Jersey, New Brunswick, NJ, USA; <sup>7</sup>Columbia University Medical Center, New York, NY, USA; <sup>8</sup>Karmanos Cancer Institute, Detroit, MI, USA; <sup>9</sup>Department of Malignant Hematology, H. Lee Moffitt Cancer Center, Tampa, FL, USA; <sup>10</sup>Norton Cancer Institute, Louisville, KY, USA; <sup>11</sup>Swedish Cancer Institute Seattle, WA, USA; <sup>12</sup>Ziekenhuis Netwerk Antwerpen Stuivenberg, Antwerp, Belgium; <sup>13</sup>Rogel Cancer Center, University of Michigan Medical School, Ann Arbor, MI, USA; <sup>14</sup>Knight Cancer Institute, Oregon Health & Science University, Portland, OR, USA; <sup>15</sup>Department of Hematology, College of Medicine, Catholic Hematology Hospital and Leukemia Research Institute, Seoul St. Mary's Hospital, The Catholic University of Korea, Seoul, South Korea; <sup>16</sup>Department of Hematology, Cliniques Universitaires Saint-Luc, Université Catholique de Louvain (UCLouvain), Brussels, Belgium; <sup>17</sup>Department of Internal Medicine III, University Medical Center Mainz, Mainz, Germany; <sup>18</sup>Department of Internal Medicine, Seoul National University College of Medicine, Seoul National University Hospital, Seoul, Korea; <sup>19</sup>Hospital Universitario 12 de octubre, Universidad Complutense, CNIO, Madrid, Spain; <sup>20</sup>Regeneron Pharmaceuticals, Inc., Tarrytown, NY, USA; <sup>21</sup>The Ohio State University Comprehensive Cancer Center, Columbus, OH, USA.

## Background

- Linvoseltamab, REGN5458, is a human BCMA×CD3 bispecific antibody that effectively induces T-cell-mediated cytotoxicity of malignant MM plasma cells
- In the ongoing LINKER-MM1 (NCT03761108) study, linvoseltamab demonstrated encouraging efficacy and a generally manageable safety profile in patients with R/R MM<sup>1</sup>
- Here we report the results at 8.1 months median duration of follow-up from all patients treated with the 200 mg dose that includes 12 Phase 1 patients and 105 Phase 2 patients


Data cut-off date of June 7, 2023.  
Lee HC, et al. American Society of Clinical Oncology (ASCO) Annual Meeting, Chicago, IL, USA, 2-6 June 2023 Oral Presentation.  
BCMA, B-cell maturation antigen; CD, cluster of differentiation; MM, multiple myeloma; R/R MM, relapsed/refractory MM.

Figure 1. Linvoseltamab mechanism of action



## LINKER-MM1 study: Phase 2 design

Figure 2. Linvoseltamab IV dosing schedule for 200 mg phase 2 expansion cohort



- The first full dose was given at week 3 of the study
- For patients who achieved VGPR or better by week 24, dosing frequency decreased from 2 to 4 weeks

### Key eligibility criteria for phase 2:

- Active MM per IMWG criteria
- Prior ≥three lines of therapy and triple-exposed disease (exposed to at least one IMiD + one PI + one anti-CD38 Ab); or triple-refractory disease (refractory to at least one IMiD + one PI + one anti-CD38 Ab)

### Key phase 2 objectives:

- Primary: ORR as determined by a blinded independent review committee (IRC; per IMWG criteria)
- Secondary: ORR by investigator assessment, DOR, PFS, MRD status, and OS

Ab, antibody; DOR, duration of response; IMiD, immunomodulatory drug; IMWG, International Myeloma Working Group; IV, intravenous; MRD, minimal residual disease; ORR, objective response rate; OS, overall survival; PI, proteasome inhibitor; PFS, progression-free survival; VGPR, very good partial response.

## Baseline Characteristics

Table 1. Patient and disease characteristics

n (%), unless otherwise stated	200 mg (n=117)
Age	70 (37–91)
Median (range)	70 (37–91)
≥75	31 (26.5)
Male	64 (54.7)
Race	
White	83 (70.9)
Black or African American	20 (17.1)
Asian	10 (8.5)
Eastern Cooperative Oncology Group (ECOG) performance status	
0	33 (28.2)
1	84 (71.8)
ISS Stage	
I	49 (41.9)
II	41 (35.0)
III	21 (17.9)
Missing	6 (5.1)
Extramedullary plasmacytomas (EMP)	17 (14.5)
High risk cytogenetics*	46 (39.3)
Bone marrow plasma cells (BMPC) percentage	
<50%	64 (54.7)
≥50%	27 (23.1)
Missing	26 (22.2)
Soluble B-cell maturation antigen (BCMA), median (range) ng/mL	365.5 (18.7–4,430.0)

\*Presence of del(17p) and/or translocation t(4;14) and/or translocation t(14;16).

Table 2. Prior treatments


n (%)	200 mg (n=117)
Prior autologous transplant	75 (64.1)
Median prior lines of therapy, n (range)	5 (2–16)
Exposure status	
At least triple-exposed	117 (100)
At least quad-exposed	112 (95.7)
At least penta-exposed	89 (76.1)
Refractory status	
At least triple-refractory	94 (80.3)
At least quad-refractory	74 (63.2)
At least penta-refractory	30 (25.6)
Refractory to last line of therapy <sup>‡</sup>	98 (83.8)

<sup>†</sup>Includes patients with a lack of response or relapse within 60 days of last line of therapy. Triple-exposed/refractory: ≥1 PI, ≥1 IMiD, and ≥1 anti-CD38 Ab. Quad-exposed/refractory: ≥2 PI, ≥1 IMiD, and ≥1 anti-CD38 Ab or ≥1 PI, ≥2 IMiD, and ≥1 anti-CD38 Ab. Penta-exposed/refractory: ≥2 PI, ≥2 IMiD, and 1 anti-CD38 Ab; BMPC, bone marrow plasma cells; IMiD, immunomodulatory drug; PI, proteasome inhibitor.

- Median age was 70 years old, with 37.6% of the population <65, 35.9% 65-74, and 26.5% ≥75 years of age
- EMP were ≥2 cm and excluded patients with only paramedullary plasmacytomas

## Efficacy per IRC assessment

Figure 3. Objective response rate




- ORR was 69.2% with 59% of patients achieving ≥ VGPR and 39% achieving ≥CR
- Median time to ≥PR was 1 month, to ≥VGPR was 2.6 months, and to ≥CR, 7.6 months
- In patients who had prior exposure to the anti-BCMA antibody-drug conjugate belantamab mafodotin (n=10), ORR was 70%
- Among patients with CR or sCR with available MRD data (N=37), 50%+ were MRD-negative at 10<sup>-5</sup>

Data cut-off date of June 7, 2023. Median duration of follow-up: 8.1 months (range 0.2–31.5).

<sup>†</sup>MRD data include clonoSEQ and Euroflow. Eight patients had missing data due to missing specimens or specimen quality. CR, complete response; PR, partial response; sCR, stringent complete response.

Figure 4. Responses per IRC are early, durable, and deepen over time\*



- At 8.1 months median duration of follow-up, 50% of patients are still ongoing core treatment

\*Estimated Kaplan–Meier method. Data cut-off date of June 7, 2023. Median duration of follow-up: 8.1 months (range 0.2–31.5). MR, minimal response; NE, not evaluable; PD, progressive disease; PR, partial response; SD, stable disease.

## Duration of Response Per IRC Assessment

Figure 5. Kaplan–Meier curve of DOR per IRC assessment





Figure 6. Kaplan–Meier curve of PFS<sup>†</sup> per IRC assessment



## Safety

Table 3. Treatment–emergent adverse events

200 mg cohort – (n=117)		
Treatment exposure, median (range), weeks		
TEAEs†, n (%)	Any grade	Grade 3–4
Hematologic TEAEs		
Neutropenia†	38.5	37.6
Anemia†	38.5	30.8
Thrombocytopenia†	17.9	14.5
Lymphopenia†	12.0	11.1
Non-hematologic TEAEs		
CRS	46.2	0.9
Diarrhea	35.0	1.7
Cough	34.2	0
Fatigue	32.5	0
Arthralgia	29.1	0
Hypokalaemia	22.2	2.6
Headache†	21.4	0.9
Nausea	21.4	0
Dyspnea	19.7	0.9
Back pain	17.9	1.7
Vomiting	17.9	0
Constipation	16.2	0
Pyrexia	16.2	0

- The most common Grade 3/4 TEAEs were hematologic events
- Common non-hematologic events were mainly Grade 1/2
- CRS was the most reported TEAE and was mostly mild to moderate (0.9%, Grade 3)
- Adjudicated ICANS\* (any grade) occurred in nine patients (7.7%); mostly after the initial dose and all concurrently with CRS; Grade 3 ICANS occurred in 3 patients, no Grade 4-5 cases
- TEAEs leading to death on treatment or within 30 days following the last dose were reported in 13 patients, with the majority due to infections (10 patients). Two events (one *pneumocystis jirovecii* pneumonia, one pseudomonal sepsis) were considered treatment drug-related, as per the treating physician.

Data cut-off date of June 7, 2023. Median duration of follow-up: 8.1 months (range 0.2–31.5). AEs per NCI-CTCAE v5.0. CRS per ASTCT (Lee DW, et al. Biol Blood Marrow Transplant. 2019;25:625-638). \*Neurologic events consistent with ICANS were reviewed by the sponsor and evaluated according to guidelines in Lee, 2019. †Table includes TEAEs of any grade reported in ≥15% of patients; ‡Composite terms. ASTCT, American Society for Transplantation and Cellular Therapy; CRS, cytokine release syndrome; CTCAE, Common Terminology Criteria for Adverse Events; ICANS, immune effector cell-associated neurotoxicity syndrome; NCI, National Cancer Institute; TEAE, treatment-emergent adverse event.


Table 4. TEAE of interest: Infections

TEAEs, n (%)	Any grade	Grade 3–4
<b>Infections</b>	81 (69.2)	43 (36.8)
Pneumonia	18 (15.4)	16 (13.7)
COVID-19	17 (14.5)	6 (5.1)
Upper respiratory tract	15 (12.8)	2 (1.7)
<b>Opportunistic infections*</b>	10 (8.5)	7 (6.0)
PJP	5 (4.3)	3 (2.6)
CMV infections	2 (1.7)	2 (1.7)
CMV reactivation	3 (2.6)	2 (1.7)

\*One case of progressive multifocal leukoencephalopathy (PML) occurred after the data cutoff date. AEs per NCI-CTCAE v5.0. CMV, cytomegalovirus; COVID-19, coronavirus disease 2019; CTCAE, Common Terminology Criteria for Adverse Events; NCI, National Cancer Institute; PJP, *Pneumocystis jirovecii* pneumonia; TEAE, treatment-emergent adverse event.

- All cases of PJP occurred prior to the institution of mandatory prophylaxis

Figure 7. A temporal analysis of CRS onset, severity and anti IL-6R use



Data cut-off date of June 7, 2023. Median duration of follow-up: 8.1 months (range 0.2–31.5).

The sunburst plot presents the incidence and severity of CRS and use of tocilizumab by patient with each of the first 8 treatment weeks as a separate ring starting with week 1 in the innermost ring. Within each ring the number of patients with each combination of CRS grade and use/no use of tocilizumab is shown by a different color.

- CRS was reported in 46.3% of patients; Grade 1 in 35.9%, Grade 2 in 9.4% of patients, and Grade 3 in 0.9% (one patient)
- Most CRS occurred in the step-up dosing period (most commonly after the first dose) and before the first full dose on week 3
- No Grade 3 or higher CRS occurred after the step-up dosing period
- No Grade 4 or Grade 5 CRS reported
- CRS onset usually occurred on the day of dosing, with resolution within 1 day
- 22 (19%) patients experienced recurrent CRS
- Tocilizumab was administered to 24 (20.5%) patients
- Recurrent CRS after tocilizumab was rare; only one patient experienced CRS after tocilizumab.

## Conclusions

- At a median follow up of 8.1 months, **linvoseltamab induced a high rate of responses (per IRC) that deepened further** since the last data cut (median follow-up of 5.6 months)
  - ❖ At the recommended dose of 200 mg: 69% ORR, 39% ≥CR, and 59% ≥VGPR
  - ❖ Among patients with CR or sCR with available MRD data (N=37), **50% had no measurable residual disease**
- **Responses were early and durable**
  - ❖ Median time to response was 1 month
  - ❖ Median DOR was not reached; **estimated 9-month rate of DOR was 86.8%**
  - ❖ Median PFS was not reached; **estimated 9-month rate of PFS was 72.8%**
- The favorable efficacy profile of linvoseltamab is complemented by a **convenient administration schedule**: Q4W from week 24 onwards for patients who reached VGPR or better
- Linvoseltamab showed a generally manageable safety profile with **no new safety signals with longer follow up**
  - ❖ The most common TEAEs were CRS, neutropenia (37.6% grade 3/4) and anemia (30.8% grade 3/4)
  - ❖ CRS rate was 46.2% (0.9% Grade 3; 0% ≥ Grade 4) and effectively managed with standard of care
- The predictable timing and low rate of CRS allows as per protocol to date an **inpatient monitoring schedule** of 1 day at week 1 and week 2
- **A Phase 3 Trial, LINKER-MM3, is ongoing** in patients with RRMM (NCT05730036)

<sup>1</sup>Lee HC, et al. American Society of Clinical Oncology (ASCO) Annual Meeting, Chicago, IL, USA, 2-6 June 2023 Oral Presentation.

## Acknowledgements

- The authors would like to thank the patients, their families, all other investigators, and all investigational site members involved in the study, especially during the challenging times of the coronavirus pandemic
- This study was funded by Regeneron Pharmaceuticals, Inc.
- Osnat Ben-Shahar, PhD, Regeneron Pharmaceuticals, Inc., assisted with the development and writing of this presentation. Editorial and studio support was provided by OPEN Health Communications and was funded by Regeneron Pharmaceuticals, Inc.

