

- Medical Analysis Report
 -  Diagnosis
 -  Additional Information Needed
 -  Additional Testing Required
 -  Recommended Tests
 -  Prescription & Treatment
 -  Detailed Analysis
 - Medical Report for Mr. Tariq Jamil Khursheed Alam
 - Summary
 - Diagnosis
 - Prescription and Recommendations

Medical Analysis Report

Generated: 2025-09-13 11:15:14

Diagnosis

Likely Dyslipidemia, Non-alcoholic Fatty Liver Disease (NAFLD), and early Chronic Kidney Disease (CKD) with an unexplained acute left-sided pain.

Additional Information Needed

 No

Additional Testing Required

 Yes

Recommended Tests

For the unexplained left-sided pain: Physical examination by a specialist (e.g., orthopedist, neurologist), X-ray/MRI of the left hip and/or lumbar spine. For kidney function: Repeat creatinine, urinalysis, and eGFR calculation.

Prescription & Treatment

Continue current medications (Cmp Nexum 3g (likely Esomeprazole), Tab Spafon, 2EGAP 5mg) as prescribed. Consider initiation of a statin for dyslipidemia, under medical guidance.

Detailed Analysis

 **Disclaimer:** This is not medical advice. Consult your doctor.

Medical Report for Mr. Tariq Jamil Khursheed Alam

Patient Name: Mr. Tariq Jamil Khursheed Alam **Age:** 52 Years **Gender:** Male

Summary

Mr. Tariq Jamil Khursheed Alam, a 52-year-old male, presents with a chief complaint of intense pain in the left side of his body, specifically just above the left leg joint. Review of recent diagnostic reports reveals several key findings:

- **Dyslipidemia:** Your lipid profile indicates elevated total cholesterol, low HDL (good cholesterol), borderline high LDL (bad cholesterol), elevated VLDL, and elevated triglycerides. This combination signifies an increased risk for cardiovascular disease.
- **Non-alcoholic Fatty Liver Disease (NAFLD):** An abdominal ultrasound confirmed the presence of fatty liver (homogeneously raised parenchymal echogenicity). Notably, your liver function tests (LFTs) are currently within normal limits, and the liver is normal in size with no focal masses or biliary duct dilation.

- **Early Kidney Function Impairment:** Your creatinine level is elevated at 1.52 mg/dL, suggesting a reduction in kidney function. However, the ultrasound showed your kidneys to be normal in size, shape, and position with no signs of stones or hydronephrosis.
- **Slightly Elevated White Blood Cell (WBC) Count:** Your WBC count is slightly above the upper normal limit at $10.35 \times 10^9/L$, which could indicate mild inflammation or a reaction to an underlying process.
- **Unremarkable Other Abdominal Organs:** The ultrasound reported all other examined abdominal organs, including the gallbladder, pancreas, spleen, and urinary bladder, as normal and without significant findings.
- **Current Medications:** You are currently prescribed Cmp Nexum 3g (likely Esomeprazole) for acid-related issues, Tab Spafon (likely an antispasmodic/pain reliever), and 2EGAP 5mg (dosage 0-0-1).

Diagnosis

Based on the available information, the likely diagnoses are:

1. **Dyslipidemia:** Characterized by abnormal levels of lipids (fats) in the blood, predisposing to heart disease.
2. **Non-alcoholic Fatty Liver Disease (NAFLD):** Accumulation of fat in the liver, often associated with metabolic risk factors like dyslipidemia.
3. **Early Chronic Kidney Disease (CKD):** Indicated by the elevated creatinine, suggesting a mild but persistent impairment of kidney function.
4. **Unexplained Left-Sided Pain:** The current diagnostic reports do not provide a clear explanation for the intense pain in your left side, just above the left leg joint. This symptom requires further dedicated investigation.

Prescription and Recommendations

1. **Current Medications:** Please continue to take Cmp Nexum 3g (likely Esomeprazole), Tab Spafon, and 2EGAP 5mg exactly as prescribed by your treating physician.
2. **For Dyslipidemia and Cardiovascular Risk:**

- **Lifestyle Changes:** Adopt a heart-healthy diet that is low in saturated and trans fats, cholesterol, and refined sugars. Increase your intake of fruits, vegetables, and whole grains. Engage in regular physical activity, aiming for at least 150 minutes of moderate-intensity exercise per week.
- **Medical Management:** Your doctor may consider initiating a statin medication to help lower your cholesterol and triglyceride levels, given your overall lipid profile and high Cholesterol:HDL-C ratio. Regular monitoring of your lipid profile will be important.

3. For Kidney Function Impairment:

- **Further Evaluation:** It is recommended to have a follow-up blood test to recheck your creatinine level, perform a urinalysis (urine test) to look for protein or blood in the urine, and calculate your estimated Glomerular Filtration Rate (eGFR). These tests will help in better assessing your kidney function and determining the underlying cause of the elevation. Your doctor may also discuss blood pressure control and diabetes screening if not already performed.

4. For Unexplained Left-Sided Pain:

- **Urgent Consultation:** It is crucial to schedule an appointment with your doctor to thoroughly investigate the intense pain in your left side, above the left leg joint. The current reports do not pinpoint its cause, and focused evaluation is needed.
- **Specialist Assessment:** Your doctor may refer you for a physical examination by an orthopedic specialist or neurologist to rule out musculoskeletal, joint, or nerve-related issues in the hip or lower spine.
- **Additional Imaging:** Depending on clinical assessment, imaging such as an X-ray or MRI of the left hip and/or lumbar spine may be necessary to identify the source of the pain.

5. **General Monitoring:** Regular follow-up with your healthcare provider is essential to monitor your chronic conditions, adjust treatments as needed, and address any new symptoms.