

# 3D Line Radiative Transfer & Synthetic Observations with Magritte

Frederik De Ceuster<sup>\*1, 2</sup>, Thomas Ceulemans<sup>2</sup>, Atulit Srivastava<sup>2</sup>,  
Jeremy Yates<sup>2</sup>, Leen Decin<sup>2, 3</sup>, Peter Boyle<sup>4, 5</sup>, and James  
Hetherington<sup>6, 7</sup>

<sup>1</sup> Department of Physics and Astronomy, University College London, London, UK <sup>2</sup> Institute of  
Astronomy, KU Leuven, Leuven, Belgium <sup>3</sup> School of Chemistry, University of Leeds, Leeds, UK <sup>4</sup>  
School of Physics and Astronomy, The University of Edinburgh, Edinburgh, UK <sup>5</sup> Brookhaven  
National Laboratory, NY, USA <sup>6</sup> Department of Computer Science, University College London,  
London, UK <sup>7</sup> The Alan Turing Institute, London, UK

DOI: [10.21105/joss.03905](https://doi.org/10.21105/joss.03905)

## Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Juanjo Bazán](#) ↗

## Reviewers:

- [@gonsie](#)
- [@CFGrote](#)

Submitted: 27 October 2021  
Published: 08 December 2021

## License

Authors of papers retain  
copyright and release the work  
under a Creative Commons  
Attribution 4.0 International  
License ([CC BY 4.0](#)).


## Summary

Magritte is a software library for 3D radiative transfer and synthetic observations, which currently focusses on atomic and molecular line transfer. It is mainly written in C++ and can either be used as a Python package or as a C++ library. To compute the radiation field, a deterministic ray-tracer and a formal solver are employed, i.e., rays are traced through the model and the radiative transfer equation is solved along those rays (De Ceuster et al., 2019). This is in contrast to most radiative transfer solvers which employ (probabilistic) Monte Carlo techniques (Noebauer & Sim, 2019). By virtue of minimal assumptions about the underlying geometric structure of a model, Magritte can handle structured and unstructured input meshes, as well as smoothed-particle hydrodynamics (SPH) data. Furthermore, tools are provided to optimise different input meshes for radiative transfer (De Ceuster et al., 2020).

## Statement of need

Electromagnetic radiation is a key component in many astrophysical simulations. Not only does it dictate what we can or cannot observe, it can provide radiation pressure, efficient heating and cooling mechanisms, and opens up a range of new chemical pathways due to photo-reactions. Magritte can be used as a general-purpose radiative transfer solver, but was particularly designed for line radiative transfer in complex 3D morphologies, such as, for instance, encountered in the stellar winds around evolved stars (see Decin, 2020). Recent high-resolution observations exposed the intricate and intrinsically 3D morphologies of these objects (Decin et al., 2020). The sheer amount of complexity that is observed, makes it difficult to interpret the observations and necessitates the use of 3D hydrodynamics, chemistry and radiative transfer models to study their origin and evolution (El Mellah et al., 2020; Maes et al., 2021; Malfait et al., 2021). Their intricate morpho-kinematics, moreover, makes their appearance in observations far from evident (e.g., Figure 1). Therefore, to study these and other complex morpho-kinematical objects, it is essential to understand how their models would appear in observations. This can be achieved, by creating synthetic observations with Magritte.

\*Corresponding author.



**Figure 1:** Example of a synthetic observation of the  $\text{CO}(v=0, J=1-0)$  transition, created with Magritte for a hydrodynamics model of an asymptotic giant branch (AGB) star, as it is perturbed by a companion (this is model v10E50 in Malfait et al., 2021).

## 38 Future work

39 Currently, Magritte is mainly being used for post-processing hydrodynamics simulations by cre-  
40 ating synthetic observations, such that the models can be compared with real observations.

41 However, Magritte is still under active development. In future work, we aim to improve on the  
 42 computational speed, to clear the path for on-the-fly radiative transfer in those simulations.  
 43 Furthermore, aside from being a practical research tool, we also aim for Magritte to be the  
 44 starting point for further research in computational radiative transfer. Current active research  
 45 topics include: efficient parallelisation and acceleration strategies on modern high-performance  
 46 computing systems, acceleration of convergence in the non-linear coupling between the ra-  
 47 diation field and the medium, and uncertainty quantification in radiative transfer through  
 48 probabilistic numerical methods (e.g., [De Ceuster, 2021](#)).

## 49 Acknowledgements

50 FDC is supported by the EPSRC iCASE studentship programme, Intel Corporation and Cray  
 51 Inc. FDC and LD acknowledge support from the ERC consolidator grant 646758 AEROSOL.

## 52 References

- 53 De Ceuster, F. (2021). *Simulating 3D Radiation Transport* [PhD]. University College London.
- 54 De Ceuster, F., Bolte, J., Homan, W., Maes, S., Malfait, J., Decin, L., Yates, J., Boyle, P.,  
 55 & Hetherington, J. (2020). Magritte, a modern software library for 3D radiative transfer:  
 56 II. Adaptive ray-tracing, mesh construction, and reduction. *Monthly Notices of the Royal*  
 57 *Astronomical Society*, 499(4), 5194–5204. <https://doi.org/10.1093/mnras/staa3199>
- 58 De Ceuster, F., Homan, W., Yates, J., Decin, L., Boyle, P., & Hetherington, J. (2019).  
 59 Magritte, a modern software library for 3D radiative transfer: I. Non-LTE atomic and  
 60 molecular line modelling. *Monthly Notices of the Royal Astronomical Society*, 492(2),  
 61 1812–1826. <https://doi.org/10.1093/mnras/stz3557>
- 62 Decin, L. (2020). Evolution and Mass Loss of Cool Ageing Stars: a Daedalean Story. *arXiv*  
 63 *e-Prints*, arXiv:2011.13472. <http://arxiv.org/abs/2011.13472>
- 64 Decin, L., Montargès, M., Richards, A. M. S., Gottlieb, C. A., Homan, W., McDonald, I., El  
 65 Mellah, I., Danilovich, T., Wallström, S. H. J., Zijlstra, A., Baudry, A., Bolte, J., Cannon,  
 66 E., De Beck, E., De Ceuster, F., de Koter, A., De Ridder, J., Etoka, S., Gobrecht, D.,  
 67 ... Yates, J. (2020). (Sub)stellar companions shape the winds of evolved stars. *Science*,  
 68 369(6510), 1497–1500. <https://doi.org/10.1126/science.abb1229>
- 69 El Mellah, I., Bolte, J., Decin, L., Homan, W., & Keppens, R. (2020). Wind morphology  
 70 around cool evolved stars in binaries: The case of slowly accelerating oxygen-rich outflows.  
 71 *Astronomy and Astrophysics*, 637, A91. <https://doi.org/10.1051/0004-6361/202037492>
- 72 Maes, S., Homan, W., Malfait, J., Siess, L., Bolte, J., De Ceuster, F., & Decin, L. (2021). SPH  
 73 modelling of companion-perturbed AGB outflows including a new morphology classifica-  
 74 tion scheme. *Astronomy & Astrophysics*, 653, A25. <https://doi.org/10.1051/0004-6361/202140823>
- 75
- 76 Malfait, J., Homan, W., Maes, S., Bolte, J., Siess, L., De Ceuster, F., & Decin, L. (2021). SPH  
 77 modelling of wind-companion interactions in eccentric AGB binary systems. *Astronomy &*  
 78 *Astrophysics*, 652, A51. <https://doi.org/10.1051/0004-6361/202141161>
- 79 Noebauer, U. M., & Sim, S. A. (2019). Monte Carlo radiative transfer. *Living Reviews in*  
 80 *Computational Astrophysics*, 5(1), 1. <https://doi.org/10.1007/s41115-019-0004-9>