




AIBECS.jl: A tool for exploring global marine biogeochemical cycles.

Benoît Pasquier^{*1,3}, François W. Primeau², and Seth G. John¹

¹ Department of Earth Sciences, University of Southern California ² Department of Earth System Science, University of California, Irvine ³ (now at) School of Mathematics and Statistics, University of New South Wales, Sydney

DOI: [10.21105/joss.03814](https://doi.org/10.21105/joss.03814)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Kristen Thyng](#) 

Reviewers:

- [@dankelley](#)
- [@zhenwu0728](#)

Submitted: 23 September 2021

Published: 10 December 2021

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

The ocean transports, mixes, and transforms chemical constituents on a multitude of time and length scales. Observations and models are both essential for making sense of this complex system. However, the days of just publishing data without any quantitative modelling are over, increasing pressure on sea-going oceanographers that are expected to be proficient in the use of biogeochemical models. And while the simplest of these models consist of only a few boxes (e.g., the two-box model of [Archer et al., 2000](#)), with solutions that are easily obtained by hand or on a simple desktop computer, the most advanced models have high-resolution three-dimensional meshes that require high-performance computing (HPC) clusters and a considerable amount of computational-science expertise (e.g., the MITgcm, [Campin et al., 2021](#)). A high barrier to entry for modelling in oceanography hinders advances in the field. AIBECS.jl ([Pasquier, 2021](#)) is a [JuliaOcean](#)-affiliated package written in [Julia](#) ([Bezanson et al., 2017](#)), which aims to lower this barrier by providing an easy-to-use, open-source, and modular framework for simulating global marine tracers in steady-state and biogeochemical parameter fitting.

A conceptual model of the cycle of any marine tracer essentially requires two components. (i) A model of how the tracer is transported (be it the ocean currents and eddies, gravitational settling, or a combination of those), and (ii) a model of the local sources and sinks at any location. AIBECS.jl is built on this concept and allows users to build numerical models of marine tracers by selecting a circulation and/or vertical transport of the tracer with particles and local sources and sinks. Tools for generating the steady-state equations, solving them, and then diagnosing and plotting the simulated tracers are also provided, either directly by AIBECS.jl, by its dependencies, or by satellite packages in the AIBECS and Julia ecosystem.

[Box \(1979\)](#) has remarked that, by definition, all models are wrong because they make simplifying assumptions. These assumptions can give rise to model parameters that can capture fundamental characteristics of the system. Once the model of the cycle of a marine tracer is implemented, it is thus natural to try to estimate biogeochemical parameters by minimizing model–observation mismatches, which may improve the skill of the model. Parameter optimization is generally more efficiently performed when first and second derivatives are available, and these derivatives can also be used to estimate parameter sensitivity ([Thacker, 1989](#)). AIBECS.jl was designed with parameter fitting in mind and provides, along with satellite packages, tools for generating first- and second-order derivatives automatically and efficiently.

The AIBECS.jl package builds on the vision of the [AWESOME OCIM](#) (A Working Environment for Simulating Ocean Movement and Elemental cycling within the Ocean Circulation Inverse

^{*}corresponding author

Model, or AO, [John et al., 2020](#)), which was designed to make three-dimensional element-cycling models more broadly accessible. Written in [Julia](#) — chosen for of its combined expressive power and efficiency and offering a truly open-source solution — the AIBECS.jl framework improves on the AO on a number of fronts to target education and research. Simple AIBECS.jl simulations can be produced in minutes in interactive notebooks while resource-hungry projects, e.g., for optimizing models with multiple tracers interacting nonlinearly within a high-resolution mesh, can be easily version-controlled, hosted, and run on HPC clusters.

Through a simple user interface, AIBECS.jl provides access to a variety of steady-state ocean circulation models. These currently include the Ocean Circulation Inverse Model (OCIM) v0.1 ([Primeau et al., 2013](#)), v1.0 ([DeVries, 2014](#); [DeVries & Primeau, 2011](#)), and v2.0 ([DeVries & Holzer, 2019](#)), and the MITgcm-built Ocean Comprehensible Atlas (OCCA) ocean-state estimate model ([Forget, 2010](#)), of which the downloads are handled by the [DataDeps.jl](#) package ([White et al., 2018, 2019](#)). AIBECS.jl also offers classic two-box and three-box models ([Archer et al., 2000](#); [Sarmiento & Gruber, 2006](#)). The [OceanGrids.jl](#) package, on which AIBECS depends, provides the underlying grid configuration types as well as regridding and interpolating routines. Swapping the underlying circulation model and grid requires a single-line-of-code change, facilitating intercomparison projects. As new circulation models that are represented in matrix form are made publicly available, they will be added to the collection. These could include past- and future-ocean circulation models, for paleoceanographic or climate-change studies, for example.

AIBECS.jl also provides extra functionality to facilitate the generation of numerical models. Tooling to simulate gravitational settling of tracers with non-buoyant particles is provided by the `transportoperator` function. In addition, AIBECS provides access to a number of predefined fields that can be used to generate source and sink processes. Fine-resolution (1-arc-minute) topography from the ETOPO1 dataset ([Amante & Eakins, 2009](#)) can be used for a refined interception of particulate fluxes by subgrid topographic features not captured by coarser circulation models. For aeolian deposition, AIBECS.jl includes aerosol-type- and region-of-origin-partitioned dust deposition fields ([Chien et al., 2016](#); [Kok et al., 2021](#)). Datasets for global river discharge ([Dai, 2017](#); [Dai & Trenberth, 2002](#)) and surface groundwater discharge ([Luijendijk et al., 2019, 2020](#)) are included. For hydrothermal-sourced tracers, the helium fluxes from the Earth's mantle computed with the OCIM v1.0 and v2.0 are available when loading the corresponding circulation models ([DeVries, 2014](#); [DeVries & Holzer, 2019](#)). AIBECS.jl also provides access to the data included with the AWESOME OCIM framework ([John et al., 2020](#)), namely data from the Global Ocean Data Analysis Project (GLODAP, [Lauvset et al., 2016](#); [Olsen et al., 2016](#)), P-cycling modelled fields from [Weber et al. \(2018\)](#), nepheloid layers ([Gardner, Richardson, & Mishonov, 2018](#); [Gardner, Richardson, Mishonov, & Biscaye, 2018](#); [Taburet et al., 2019](#)), as well as other data already present within AIBECS or satellite packages. Also useful to global biogeochemistry modelling are data from the World Ocean Atlas ([Garcia et al., 2018](#)) that can be downloaded, assisted by external package [WorldOceanAtlasTools.jl](#) ([Pasquier, 2019](#)). Similarly, GEOTRACES data ([Schlitzer et al., 2018](#)) can be handled by the [GEOTRACES.jl](#) package (although GEOTRACES requires manual download of the data). More advanced usage such as optimization is facilitated by the [F1Method.jl](#) package ([Pasquier, 2020](#)), which provides efficient gradient and Hessian computations of objective functions defined through AIBECS.jl, which can then be directly fed to optimization routines from, e.g., the [Optim.jl](#) package ([Mogensen & Riseth, 2018](#)). Finally, plotting recipes for the [Plots.jl](#) package ([Breloff, 2021](#)), are available.

Internally, AIBECS.jl uses a quasi-Newton solver ([Kelley, 2003](#)) translated from MATLAB to Julia and tailored to the context of marine tracers to solve/simulate tracers. AIBECS uses forward-mode auto-differentiation from the [ForwardDiff.jl](#) package ([Revels et al., 2016](#)) for the nonlinear parts of the system of ordinary differential equations to generate the Jacobian required for the solver. Metadata such as units and prior distributions can be attached to model parameters, which are handled with the help of the [UnPack.jl](#), [FieldMetadata.jl](#), [Flatten.jl](#), [DataFrames.jl](#), [Distributions.jl](#) ([Besançon et al., 2021](#); [Lin et al., 2019](#)), [Unitful.jl](#), and

95 [Bijectors.jl](#) dependencies.

96 The [AIBECS.jl](#) package is registered with Julia's [default package registry \(GENERAL\)](#), such
 97 that installation takes a single line of code from within Julia. The package [documentation](#),
 98 which is built through continuous integration (CI), includes [tutorials](#) and [how-to guides](#) that
 99 are both available online for consultation and as runnable Jupyter notebooks. Continuous
 100 integration through GitHub actions also includes a fairly complete suite of tests.

101 Statement of need

102 Pioneered by [Schlitzer \(1993\)](#) to study the ocean circulation using ventilation tracers such
 103 as radiocarbon (see also [Schlitzer, 2000](#)), the low computational costs of steady-state cir-
 104 culation models allow for efficient optimization and inference/estimation of biogeochemical
 105 parameters. Today, despite reduced resolution and steady-state assumption, data-constrained
 106 matrix-transport models such as the OCIM are at the forefront of oceanographic research.
 107 This is evidenced for example by the growing number of high-profile studies that use such
 108 models for parameter estimation published in recent years (e.g., [DeVries et al., 2017, 2012,](#)
 109 [2013; DeVries, 2014; DeVries & Deutsch, 2014; DeVries & Weber, 2017; Roshan & DeVries,](#)
 110 [2017; Teng et al., 2014; Wang et al., 2019; Weber et al., 2016; Weber & Deutsch, 2010](#)).
 111 However, although the steady-state assumption and matrix representation simplify the simu-
 112 lation of tracers compared to traditional Ocean General Circulation Models (OGCMs), most
 113 studies that employ a steady-state matrix representation of marine cycling remain difficult to
 114 reproduce without significant computer-science and modelling expertise because they are built
 115 on private implementations. Comparisons between different circulation models are moreover
 116 complicated by the lack of standardization across models.

117 Hence, there is a need to facilitate the use of steady-state ocean-circulation models by provid-
 118 ing: (i) an integrated framework for handling a number of different ocean-circulation models
 119 with tools for swapping circulations (including interpolating from one model grid to another),
 120 (ii) a user-friendly interface for translating mathematical models of biogeochemical cycles into
 121 the corresponding code (e.g., for sources, sinks, and vertical transport of tracers), and (iii)
 122 solvers for efficient simulations, optimization, diagnosis, and statistical analysis.

123 AIBECS.jl provides a free, open-source, unified framework for biogeochemical-tracer-modelling
 124 studies that use steady-state circulation models. Among other advantages over existing so-
 125 lutions (i.e., the AO), AIBECS.jl offers better computational efficiency, enhanced versatility,
 126 composability with other Julia packages, and ease of reproducibility (granted by version con-
 127 trol and Julia's package manager) and improved syntax, which are pillars of modern scientific
 128 dissemination. Thus, AIBECS users may include sea-going oceanographers and educators who
 129 will benefit from its simplicity, as well as more experienced modellers who can leverage its
 130 computational advantages. AIBECS.jl has been used for teaching and is currently used for
 131 research focused on marine trace metals.

132 The publication of circulation models as transport matrices from existing general circula-
 133 tion models ([Bardin et al., 2014, 2016; Khatiwala et al., 2005; Khatiwala, 2007; Kvale et al.,](#)
 134 [2017; Zanna et al., 2019](#)), and hopefully, future publication of transport matrix estimates from
 135 standard ocean general circulation models (e.g., [Chamberlain et al., 2019](#)) will increase the
 136 collection of circulations available from AIBECS.jl. Of particular interest to the broader com-
 137 munity to facilitate the simulation of past and future marine biogeochemical states would be
 138 transport matrix models extracted from the Climate Model Intercomparison Project (CMIP),
 139 which includes past and future simulations of the ocean circulation.

140 Further development could include exposing advanced Green-function-based diagnostic tools
 141 (e.g., [Holzer et al., 2021; Pasquier & Holzer, 2018](#)), coupling tracers on different grids,
 142 Newton-Krylov solvers for cyclo-stationary states (e.g., CYCLOCIM: [Huang et al., 2021](#)), or
 143 time-dependent solvers for transient biogeochemical simulations provided, e.g., by the SciML

ecosystem (Rackauckas & Nie, 2017). Bridging packages could be implemented for improved composability with statistical packages (e.g., Turing.jl: Ge et al., 2018), optimization tools, and plotting software (e.g., Makie.jl: Danisch et al., 2021; Danisch & Krumbiegel, 2021).

Acknowledgements

FWP and BP acknowledge funding from the Department of Energy (grant DE-SC0016539) and the National Science Foundation (grant 1658380). BP and SGJ acknowledge funding provided by the Simons Foundation (Award #426570SP to SGJ) and the National Science Foundation (grant 1736896). The authors thank Dr. Zhen Wu and Prof. Dan Kelley for their insightful reviews that have helped improve the software and this manuscript.

References

- Amante, C., & Eakins, B. W. (2009). ETOPO1 1 arc-minute global relief model: Procedures, data sources and analysis. *NOAA Technical Memorandum NESDIS NGDC-24*. <https://doi.org/10.7289/V5C8276M>
- Archer, D. E., Eshel, G., Winguth, A., Broecker, W., Pierrehumbert, R., Tobis, M., & Jacob, R. (2000). Atmospheric pCO₂ sensitivity to the biological pump in the ocean. *Global Biogeochemical Cycles*, 14(4), 1219–1230. <https://doi.org/10.1029/1999GB001216>
- Bardin, A., Primeau, F., & Lindsay, K. (2014). An offline implicit solver for simulating prebomb radiocarbon. *Ocean Modelling*, 73, 45–58. <https://doi.org/10.1016/j.ocemod.2013.09.008>
- Bardin, A., Primeau, F., Lindsay, K., & Bradley, A. (2016). Evaluation of the accuracy of an offline seasonally-varying matrix transport model for simulating ideal age. *Ocean Modelling*, 105, 25–33. <https://doi.org/10.1016/j.ocemod.2016.07.003>
- Besançon, M., Papamarkou, T., Anthoff, D., Arslan, A., Byrne, S., Lin, D., & Pearson, J. (2021). Distributions.jl: Definition and modeling of probability distributions in the JuliaStats ecosystem. *Journal of Statistical Software*, 98(16), 1–30. <https://doi.org/10.18637/jss.v098.i16>
- Bezanson, J., Edelman, A., Karpinski, S., & Shah, V. B. (2017). Julia: A fresh approach to numerical computing. *SIAM Review*, 59(1), 65–98. <https://doi.org/10.1137/141000671>
- Box, G. E. P. (1979). Robustness in the strategy of scientific model building. In R. L. Launer & G. N. Wilkinson (Eds.), *Robustness in statistics* (pp. 201–236). Academic Press. <https://doi.org/10.1016/B978-0-12-438150-6.50018-2>
- Breloff, T. (2021). *Plots.jl* (Version v1.16.3) [Computer software]. Zenodo. <https://doi.org/10.5281/zenodo.4725317>
- Campin, J.-M., Heimbach, P., Losch, M., Forget, G., edhill3, Adcroft, A., amolod, Menemenlis, D., dfer22, Hill, C., Jahn, O., Scott, J., stephdut, Mazloff, M., Fox-Kemper, B., antnguyen13, Doddridge, E., Fenty, I., Bates, M., ... dussin, raphael. (2021). *MIT-gcm/MITgcm: checkpoint67z* (Version checkpoint67z) [Computer software]. Zenodo. <https://doi.org/10.5281/zenodo.1409237>
- Chamberlain, M. A., Matear, R. J., Holzer, M., Bi, D., & Marsland, S. J. (2019). Transport matrices from standard ocean-model output and quantifying circulation response to climate change. *Ocean Modelling*, 135, 1–13. <https://doi.org/10.1016/j.ocemod.2019.01.005>
- Chien, C.-T., Mackey, K. R. M., Dutkiewicz, S., Mahowald, N. M., Prospero, J. M., & Paytan, A. (2016). Effects of African dust deposition on phytoplankton in the western

- 187 tropical Atlantic Ocean off Barbados. *Global Biogeochemical Cycles*, 30(5), 716–734.
188 <https://doi.org/10.1002/2015GB005334>
- 189 Dai, A. (2017). *Dai and trenberth global river flow and continental discharge dataset*. Re-
190 search Data Archive at the National Center for Atmospheric Research, Computational;
191 Information Systems Laboratory. <https://doi.org/10.5065/D6V69H1T>
- 192 Dai, A., & Trenberth, K. E. (2002). Estimates of Freshwater Discharge from Continents:
193 Latitudinal and Seasonal Variations. *Journal of Hydrometeorology*, 3(6), 660–687. [https://doi.org/10.1175/1525-7541\(2002\)003%3C0660:EOFDFC%3E2.0.CO;2](https://doi.org/10.1175/1525-7541(2002)003%3C0660:EOFDFC%3E2.0.CO;2)
- 195 Danisch, S., Krumbiege, J., & others. (2021). *JuliaPlots/makie.jl* (Version v0.15.0) [Com-
196 puter software]. Zenodo. <https://doi.org/10.5281/zenodo.3735092>
- 197 Danisch, S., & Krumbiegel, J. (2021). Makie.jl: Flexible high-performance data visualization
198 for julia. *Journal of Open Source Software*, 6(65), 3349. <https://doi.org/10.21105/joss.03349>
- 200 DeVries, T. (2014). The oceanic anthropogenic CO₂ sink: Storage, air-sea fluxes, and trans-
201 ports over the industrial era. *Global Biogeochemical Cycles*, 28(7), 631–647. <https://doi.org/10.1002/2013GB004739>
- 203 DeVries, T., & Deutsch, C. (2014). Large-scale variations in the stoichiometry of marine
204 organic matter respiration. *Nature Geoscience*, 7(12), 890–894. <https://doi.org/10.1038/ngeo2300>
- 206 DeVries, T., Deutsch, C., Rafter, P. A., & Primeau, F. (2013). Marine denitrification rates
207 determined from a global 3-D inverse model. *Biogeosciences*, 10(4), 2481–2496. <https://doi.org/10.5194/bg-10-2481-2013>
- 209 DeVries, T., & Holzer, M. (2019). Radiocarbon and helium isotope constraints on deep ocean
210 ventilation and mantle-³He sources. *Journal of Geophysical Research: Oceans*, 124(5),
211 3036–3057. <https://doi.org/10.1029/2018JC014716>
- 212 DeVries, T., Holzer, M., & Primeau, F. (2017). Recent increase in oceanic carbon uptake
213 driven by weaker upper-ocean overturning. *Nature*, 542(7640), 215–218. <https://doi.org/10.1038/nature21068>
- 215 DeVries, T., & Primeau, F. (2011). Dynamically and observationally constrained estimates of
216 water-mass distributions and ages in the global ocean. *Journal of Physical Oceanography*,
217 41, 2381–2401. <https://doi.org/10.1175/JPO-D-10-05011.1>
- 218 DeVries, T., Primeau, F., & Deutsch, C. (2012). The sequestration efficiency of the biological
219 pump. *Geophysical Research Letters*, 39(13). <https://doi.org/10.1029/2012GL051963>
- 220 DeVries, T., & Weber, T. (2017). The export and fate of organic matter in the ocean:
221 New constraints from combining satellite and oceanographic tracer observations. *Global*
222 *Biogeochemical Cycles*, 31(3), 535–555. <https://doi.org/10.1002/2016GB005551>
- 223 Forget, G. (2010). Mapping ocean observations in a dynamical framework: A 2004–06 ocean
224 atlas. *Journal of Physical Oceanography*, 40(6), 1201–1221. <https://doi.org/10.1175/2009JPO4043.1>
- 226 Garcia, H. E., Weathers, K., Paver, C. R., Smolyar, I., Boyer, T. P., Locarnini, R. A., Zweng,
227 M. M., Mishonov, A. V., Baranova, O. K., Seidov, D., & Reagan, J. R. (2018). *World*
228 *Ocean Atlas 2018: Vols. Volume 4: Dissolved Inorganic Nutrients (phosphate, nitrate and*
229 *nitrate+nitrite, silicate). Volume 4: Dissolved Inorganic Nutrients (phosphate, nitrate and*
230 *nitrate+nitrite, silicate).*
- 231 Gardner, W. D., Richardson, J., & Mishonov, A. V. (2018). Global assessment of benthic
232 nepheloid layers and linkage with upper ocean dynamics. *Earth and Planetary Science*
233 *Letters*, 482, 126–134. <https://doi.org/10.1016/j.epsl.2017.11.008>

- 234 Gardner, W. D., Richardson, J., Mishonov, A. V., & Biscaye, P. E. (2018). Global compar-
 235 ison of benthic nepheloid layers based on 52 years of nephelometer and transmissome-
 236 ter measurements. *Progress in Oceanography*, 168, 100–111. <https://doi.org/https://doi.org/10.1016/j.pocean.2018.09.008>
 237
- 238 Ge, H., Xu, K., & Ghahramani, Z. (2018). Turing: A language for flexible probabilistic
 239 inference. *International Conference on Artificial Intelligence and Statistics, AISTATS 2018,*
 240 *9-11 April 2018, Playa Blanca, Lanzarote, Canary Islands, Spain*, 1682–1690. <http://proceedings.mlr.press/v84/ge18b.html>
 241
- 242 Holzer, M., Kwon, E. Y., & Pasquier, B. (2021). A new metric of the biological carbon pump:
 243 Number of pump passages and its control on atmospheric pCO₂. *Global Biogeochemical*
 244 *Cycles*, 35(6), e2020GB006863. <https://doi.org/10.1029/2020GB006863>
- 245 Huang, Q., Primeau, F., & DeVries, T. (2021). CYCLOCIM: A 4-d variational assimila-
 246 tion system for the climatological mean seasonal cycle of the ocean circulation. *Ocean*
 247 *Modelling*, 159, 101762. <https://doi.org/10.1016/j.ocemod.2021.101762>
- 248 John, S. G., Liang, H., Weber, T., DeVries, T., Primeau, F., Moore, K., Holzer, M., Ma-
 249 howald, N., Gardner, W., Mishonov, A., Richardson, J., Faugere, Y., & Taburet, G. (2020).
 250 AWESOME OCIM: A simple, flexible, and powerful tool for modeling elemental cycling
 251 in the oceans. *Chemical Geology*, 533, 119403. <https://doi.org/10.1016/j.chemgeo.2019.119403>
 252
- 253 Kelley, C. T. (2003). *Solving nonlinear equations with newton's method* (pp. 1–25). SIAM.
 254 <https://doi.org/10.1137/1.9780898718898.ch1>
- 255 Khatiwala, S. (2007). A computational framework for simulation of biogeochemical tracers in
 256 the ocean. *Global Biogeochemical Cycles*, 21(3). <https://doi.org/10.1029/2007GB002923>
- 257 Khatiwala, S., Visbeck, M., & Cane, M. A. (2005). Accelerated simulation of passive tracers
 258 in ocean circulation models. *Ocean Modelling*, 9(1), 51–69. <https://doi.org/10.1016/j.ocemod.2004.04.002>
 259
- 260 Kok, J. F., Adebisi, A. A., Albani, S., Balkanski, Y., Checa-Garcia, R., Chin, M., Colarco, P.
 261 R., Hamilton, D. S., Huang, Y., Ito, A., Klose, M., Li, L., Mahowald, N. M., Miller, R. L.,
 262 Obiso, V., Pérez García-Pando, C., Rocha-Lima, A., & Wan, J. S. (2021). Contribution
 263 of the world's main dust source regions to the global cycle of desert dust. *Atmospheric*
 264 *Chemistry and Physics*, 21(10), 8169–8193. <https://doi.org/10.5194/acp-21-8169-2021>
- 265 Kvale, K. F., Khatiwala, S., Dietze, H., Kriest, I., & Oschlies, A. (2017). Evaluation of the
 266 transport matrix method for simulation of ocean biogeochemical tracers. *Geoscientific*
 267 *Model Development*, 10(6), 2425–2445. <https://doi.org/10.5194/gmd-10-2425-2017>
- 268 Lauvset, S. K., Key, R. M., Olsen, A., Heuven, S. van, Velo, A., Lin, X., Schirnick, C., Kozyr,
 269 A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M.,
 270 Perez, F. F., Suzuki, T., & Watelet, S. (2016). A new global interior ocean mapped
 271 climatology: The 1°×1° GLODAP version 2. *Earth System Science Data*, 8(2), 325–340.
 272 <https://doi.org/10.5194/essd-8-325-2016>
- 273 Lin, D., White, J. M., Byrne, S., Bates, D., Noack, A., Pearson, J., Arslan, A., Squire, K.,
 274 Anthoff, D., Papamarkou, T., Besançon, M., Drugowitsch, J., Schauer, M., & contributors,
 275 other. (2019). *JuliaStats/Distributions.jl: a Julia package for probability distributions and*
 276 *associated functions*. <https://doi.org/10.5281/zenodo.2647458>
- 277 Luijendijk, E., Gleeson, T., & Moosdorf, N. (2019). *Geospatial data and model results for*
 278 *a global model study of coastal groundwater discharge [Data set]*. PANGAEA. <https://doi.org/10.1594/PANGAEA.907641>
 279
- 280 Luijendijk, E., Gleeson, T., & Moosdorf, N. (2020). Fresh groundwater discharge insignificant
 281 for the world's oceans but important for coastal ecosystems. *Nature Communications*,
 282 11(1). <https://doi.org/10.1038/s41467-020-15064-8>

- 283 Mogensen, P. K., & Riseth, A. N. (2018). Optim: A mathematical optimization package for
284 Julia. *Journal of Open Source Software*, 3(24), 615. <https://doi.org/10.21105/joss.00615>
- 285 Olsen, A., Key, R. M., Heuven, S. van, Lauvset, S. K., Velo, A., Lin, X., Schirnack, C.,
286 Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii,
287 M., Pérez, F. F., & Suzuki, T. (2016). The global ocean data analysis project version 2
288 (GLODAPv2) – an internally consistent data product for the world ocean. *Earth System*
289 *Science Data*, 8(2), 297–323. <https://doi.org/10.5194/essd-8-297-2016>
- 290 Pasquier, B. (2019). *WorldOceanAtlasTools.jl: Download and process world ocean data in*
291 *julia*. <https://doi.org/10.5281/zenodo.2677666>
- 292 Pasquier, B. (2021). *AIBECS.jl: The ideal tool for exploring global marine biogeochemical*
293 *cycles* (Version v0.10.10) [Computer software]. Zenodo. [https://doi.org/10.5281/zenodo.](https://doi.org/10.5281/zenodo.2864051)
294 [2864051](https://doi.org/10.5281/zenodo.2864051)
- 295 Pasquier, B. (2020). *F1Method.jl* (Version v0.4.0) [Computer software]. Zenodo. <https://doi.org/10.5281/zenodo.4284133>
- 296
297 Pasquier, B., & Holzer, M. (2018). The number of past and future regenerations of iron
298 in the ocean and its intrinsic fertilization efficiency. *Biogeosciences*, 15(23), 7177–7203.
299 <https://doi.org/10.5194/bg-15-7177-2018>
- 300 Primeau, F., Holzer, M., & DeVries, T. (2013). Southern Ocean nutrient trapping and the
301 efficiency of the biological pump. *Journal of Geophysical Research: Oceans*, 118, 2547–
302 2564. <https://doi.org/10.1002/jgrc.20181>
- 303 Rackauckas, C., & Nie, Q. (2017). Differentialequations.jl—a performant and feature-rich
304 ecosystem for solving differential equations in julia. *Journal of Open Research Software*,
305 5(1).
- 306 Revels, J., Lubin, M., & Papamarkou, T. (2016). Forward-mode automatic differentiation in
307 Julia. *arXiv:1607.07892 [cs.MS]*. <https://arxiv.org/abs/1607.07892>
- 308 Roshan, S., & DeVries, T. (2017). Efficient dissolved organic carbon production and ex-
309 port in the oligotrophic ocean. *Nature Communications*, 8(1). [https://doi.org/10.1038/](https://doi.org/10.1038/s41467-017-02227-3)
310 [s41467-017-02227-3](https://doi.org/10.1038/s41467-017-02227-3)
- 311 Sarmiento, J. L., & Gruber, N. (2006). *Ocean biogeochemical dynamics* (p. 503). Princeton
312 Univ. Press, Princeton; Oxford.
- 313 Schlitzer, R. (1993). Determining the mean, large-scale circulation of the atlantic with the
314 adjoint method. *Journal of Physical Oceanography*, 23(9), 1935–1952. [https://doi.org/](https://doi.org/10.1175/1520-0485(1993)023%3C1935:DTMLSC%3E2.0.CO;2)
315 [10.1175/1520-0485\(1993\)023%3C1935:DTMLSC%3E2.0.CO;2](https://doi.org/10.1175/1520-0485(1993)023%3C1935:DTMLSC%3E2.0.CO;2)
- 316 Schlitzer, R. (2000). Applying the adjoint method for biogeochemical modeling: Export of par-
317 ticulate organic matter in the world ocean. *GEOPHYSICAL MONOGRAPH-AMERICAN*
318 *GEOPHYSICAL UNION*, 114, 107–124.
- 319 Schlitzer, R., Anderson, R. F., Dodas, E. M., Lohan, M., Geibert, W., Tagliabue, A., Bowie, A.,
320 Jeandel, C., Maldonado, M. T., Landing, W. M., Cockwell, D., Abadie, C., Abouchami,
321 W., Achterberg, E. P., Agather, A., Aguiar-Islas, A., Aken, H. M. van, Andersen, M.,
322 Archer, C., ... Shelley, Rachel. (2018). The GEOTRACES Intermediate Data Product
323 2017. *Chemical Geology*, 493, 210–223.
- 324 Taburet, G., Sanchez-Roman, A., Ballarotta, M., Pujol, M.-I., Legeais, J.-F., Fournier, F.,
325 Faugere, Y., & Dibarboure, G. (2019). DUACS DT2018: 25 years of reprocessed sea
326 level altimetry products. *Ocean Science*, 15(5), 1207–1224. [https://doi.org/10.5194/](https://doi.org/10.5194/os-15-1207-2019)
327 [os-15-1207-2019](https://doi.org/10.5194/os-15-1207-2019)
- 328 Teng, Y.-C., Primeau, F., Moore, J. K., Lomas, M. W., & Martiny, A. C. (2014). Global-scale
329 variations of the ratios of carbon to phosphorus in exported marine organic matter. *Nature*
330 *Geoscience*, 7(12), 895–898. <https://doi.org/10.1038/ngeo2303>

- 331 Thacker, W. C. (1989). The role of the hessian matrix in fitting models to measurements.
332 *Journal of Geophysical Research: Oceans*, 94(C5), 6177–6196. <https://doi.org/10.1029/JC094iC05p06177>
333
- 334 Wang, W.-L., Moore, J. K., Martiny, A. C., & Primeau, F. (2019). Convergent estimates
335 of marine nitrogen fixation. *Nature*, 566(7743), 205–211. <https://doi.org/10.1038/s41586-019-0911-2>
336
- 337 Weber, T., Cram, J. A., Leung, S. W., DeVries, T., & Deutsch, C. (2016). Deep ocean
338 nutrients imply large latitudinal variation in particle transfer efficiency. *Proceedings of*
339 *the National Academy of Sciences*, 113(31), 8606–8611. <https://doi.org/10.1073/pnas.1604414113>
340
- 341 Weber, T., & Deutsch, C. (2010). Ocean nutrient ratios governed by plankton biogeography.
342 *Nature*, 467(7315), 550–554. <https://doi.org/10.1038/nature09403>
- 343 Weber, T., John, S., Tagliabue, A., & DeVries, T. (2018). Biological uptake and reversible
344 scavenging of zinc in the global ocean. *Science*, 361(6397), 72–76. <https://doi.org/10.1126/science.aap8532>
345
- 346 White, L., Togneri, R., Liu, W., & Bennamoun, M. (2018). DataDeps.jl: Repeatable data
347 setup for replicable data science. *CoRR*, abs/1808.01091. <http://arxiv.org/abs/1808.01091>
348
- 349 White, L., Togneri, R., Liu, W., & Bennamoun, M. (2019). DataDeps.jl: Repeatable data
350 setup for reproducible data science. *Journal of Open Research Software*, 7. <https://doi.org/10.5334/jors.244>
351
- 352 Zanna, L., Khatiwala, S., Gregory, J. M., Ison, J., & Heimbach, P. (2019). Global reconstruction
353 of historical ocean heat storage and transport. *Proceedings of the National Academy of*
354 *Sciences*, 116(4), 1126–1131. <https://doi.org/10.1073/pnas.1808838115>