

INVERSE TRIGONOMETRIC FUNCTIONS – CLASS 12 SUPER STUDY GUIDE

1. THEORY IN SIMPLE WORDS (WITH VISUALS & ANALOGIES)

★ 1.1 What Are Inverse Trigonometric Functions?

Think of trigonometric functions (sin, cos, tan...) as **machines**:

Angle \rightarrow sin machine \rightarrow **Ratio**
(θ) ($\sin \theta$)

Inverse trigonometric functions **reverse the machine**:

Ratio \rightarrow \sin^{-1} machine \rightarrow **Angle**
($\frac{1}{2}$) ($\sin^{-1} \frac{1}{2} = \pi/6$)

💡 Super Simple Analogy

- Trigonometric function = “What is the ratio for this angle?”
 - Inverse trigonometric function = “What is the angle for this ratio?”
-

★ 1.2 Why Do We Need Principal Values?

A single ratio (like $\sqrt{2}/2$) comes from **many angles**:
 $\pi/4, 7\pi/4, 9\pi/4 \dots$

👉 Too many answers = confusion = not suitable for a function.
So mathematicians **restrict the domain** of trig functions to make them **one-one**, and hence invertible.

Visual Memory Trick

Imagine cutting a **slice** from a big unit-circle “pizza” so that every ratio gives **only one unique angle**.

★ 1.3 Principal Value Branches (IMPORTANT!)

Function	Principal Value Range	Easy Visual
$\sin^{-1} x$	$[-\pi/2, \pi/2]$	Right/Left tilt, no full rotation

Function	Principal Value Range	Easy Visual
$\cos^{-1} x$	$[0, \pi]$	Top semicircle only
$\tan^{-1} x$	$(-\pi/2, \pi/2)$	Up-Down strip
$\cot^{-1} x$	$((0, \pi)$	Rightwards exclusive
$\sec^{-1} x$	$([0, \pi] - \{\pi/2\})$	Top semicircle skipping vertical
$\operatorname{cosec}^{-1} x$	$[-\pi/2, \pi/2] - \{0\}$	Tilt up/down, no horizontal

★ 1.4 How to Think of Inverse Trig Graphs

All inverse trig graphs:

- Are **smooth curves**
- Are **monotonic**
- Show **angles on y-axis** and **ratios on x-axis**

Example for $\sin^{-1}x$:

Starts at $-\pi/2$ and ends at $\pi/2$.

★ 1.5 Common Identity Patterns

Think of them as “angle-simplifiers”.

- $\sin(\sin^{-1}x) = x$
- $\sin^{-1}(\sin x) = \text{principal value of } x$
(Same rule for all others.)

2. KEY CONCEPTS & FORMULAS (QUICK REVISION TABLES)

★ Big Formula Table

Basic Identities

Expression	Simplified Value
$\sin(\sin^{-1} x)$	x
$\cos(\cos^{-1} x)$	x
$\tan(\tan^{-1} x)$	x

Expression	Simplified Value
$\sin^{-1}(\sin x)$	PV of x
$\cos^{-1}(\cos x)$	PV of x
$\tan^{-1}(\tan x)$	PV of x

★ Angle Transformation Formulas (VERY IMPORTANT FOR BOARDS)

Formula	Value
$\sin^{-1}x + \cos^{-1}x$	$\pi/2$
$\tan^{-1}x + \cot^{-1}x$	$\pi/2$
$\tan^{-1}x + \tan^{-1}y$	$\tan^{-1}((x+y)/(1-xy))$
$\tan^{-1}x - \tan^{-1}y$	$\tan^{-1}((x-y)/(1+xy))$

★ Sign & Quadrant Memory Trick

- $\sin^{-1} x \rightarrow$ angle between -90° to 90°
- $\cos^{-1} x \rightarrow$ angle between 0° to 180°
- $\tan^{-1} x \rightarrow$ angle between -90° to 90°

🎯 Mnemonic:

"S & T stay small, C goes to the ceiling."

3. SOLVED NUMERICAL PROBLEMS (STEP-BY-STEP)

★ TYPE 1: Evaluate Inverse Trig Values

Q1: Find $\sin^{-1}(1/2)$

Step 1: Think: "sin what = $1/2$?"

= $\pi/6$

Step 2: Check if $\pi/6$ in principal range: YES

Final Answer: $\pi/6$

Q2: Calculate $\cos^{-1}(-\sqrt{3}/2)$

$$\cos \theta = -\sqrt{3}/2 \rightarrow \text{angle} = 5\pi/6$$

(\because cos is negative in II quadrant, allowed in $[0, \pi]$)

★ TYPE 2: Simplify Expressions

Q3: Find $\sin(\sin^{-1} x + \cos^{-1} x)$

$$\sin(A + B) = \sin(\pi/2) = 1$$

✓ Final Ans: 1

Q4: Show $\sin^{-1}x + \cos^{-1}x = \pi/2$

$$\text{Let } \sin^{-1}x = \theta \rightarrow \sin \theta = x$$

$$\cos^{-1}x = \pi/2 - \theta$$

$$\text{Add: } \theta + (\pi/2 - \theta) = \pi/2 \quad \checkmark$$

★ TYPE 3: \tan^{-1} Identities

Q5: $\tan^{-1}(1) + \tan^{-1}(2)$

Use formula:

$$\tan^{-1}x + \tan^{-1}y = \tan^{-1}((x+y)/(1-xy))$$

$$= \tan^{-1}((1+2)/(1-2))$$

$$= \tan^{-1}(3/-1) = \tan^{-1}(-3)$$

Angle lies in $(-\pi/2, \pi/2) \Rightarrow$ valid.



4. PREVIOUS YEARS' BOARD QUESTIONS (SOLVED)

(Based on typical CBSE patterns)

★ PYQ 1: Evaluate $\sin(\sin^{-1}(3/5) + \cos^{-1}(4/5))$

$$\cos^{-1}(4/5) = \theta \Rightarrow \cos \theta = 4/5 \Rightarrow \sin \theta = 3/5$$

$$\text{So } \sin^{-1}(3/5) + \cos^{-1}(4/5) = \pi/2$$

$$\text{Therefore } \sin(\pi/2) = 1$$

★ PYQ 2: Find the value of $\tan^{-1}(1/2) + \tan^{-1}(1/3)$

Use identity:

$$\tan^{-1}x + \tan^{-1}y = \tan^{-1}((x+y)/(1-xy))$$

$$= \tan^{-1}((1/2 + 1/3)/(1 - 1/6))$$

$$= \tan^{-1}((5/6)/(5/6))$$

$$= \tan^{-1}(1)$$

$$= \pi/4$$

★ PYQ 3: Prove $\sin^{-1}x + \sin^{-1}y = \sin^{-1}(x\sqrt{1-y^2} + y\sqrt{1-x^2})$

Standard identity. Proof involves $\sin(A + B)$ formula.

★ Frequently Asked Concepts (Based on trends)

- ✓ Principal values (VERY frequent)
 - ✓ \tan^{-1} identities
 - ✓ Simplification of composite expressions
 - ✓ Graph-based questions (rare but possible)
-

⚡ 5. QUICK REVISION NOTES (1–2 PAGES)

★ DEFINITIONS

- Inverse trig functions give **angle** for a **ratio**
 - Need principal values → make function one-one
 - Use identity $\sin^{-1}x + \cos^{-1}x = \pi/2$ everywhere
-

★ IMPORTANT VALUES

x	$\sin^{-1}x$	$\cos^{-1}x$	$\tan^{-1}x$
0	0	$\pi/2$	0
1	$\pi/2$	0	$\pi/4$
-1	$-\pi/2$	π	$-\pi/4$

★ GRAPH SHAPES

- $\sin^{-1}x \rightarrow$ S-curve
- $\cos^{-1}x \rightarrow$ decreasing curve

- $\tan^{-1} x \rightarrow$ sigmoid curve
-

★ DO THESE ALWAYS

- ✓ Check principal value
 - ✓ Use identity charts
 - ✓ Convert to standard angles
 - ✓ For negative inputs \rightarrow use symmetry:
 - $\sin^{-1}(-x) = -\sin^{-1}x$
 - $\tan^{-1}(-x) = -\tan^{-1}x$
 - $\cos^{-1}(-x) = \pi - \cos^{-1}x$
-



6. PREDICTED / LIKELY QUESTIONS FOR BOARD 2025

Short Answer

1. State principal value of $\tan^{-1}(-\sqrt{3})$.
2. Simplify $\sin^{-1}x + \cos^{-1}x$.
3. Plot rough graph of $\sin^{-1}x$.

Long/Numerical

1. Prove the identity involving $\sin^{-1}x + \sin^{-1}y$.
 2. Evaluate $\tan^{-1}a + \tan^{-1}b$.
 3. Solve composite expression:
 $\tan^{-1}(2) - \tan^{-1}(3)$
-



7. EXAM TIPS & TRICKS

★ Golden Rule

- 👉 Always check the principal value range first.
This removes 90% errors.

★ Common Mistakes

- ✗ Writing multiple answers (only one PV allowed)
 - ✗ Using wrong quadrant
 - ✗ Mixing up \tan^{-1} identities
 - ✗ Forgetting denominators in formulas
-

★ Time-Saving Flowchart

Start with given expression



Convert values to standard angles



Apply inverse trig identities



Check principal value range



Final simplified answer






8. SUPER KID-FRIENDLY VISUAL RECAP

- Inverse trig = Angle Finder Machine
- Principal value = Special slice of circle
- $\sin^{-1}x \rightarrow$ small tilt angles
- $\cos^{-1}x \rightarrow$ top semi circle
- $\tan^{-1}x \rightarrow$ slanted S-curve



Memory Colors

-  $\sin^{-1} \rightarrow$ green (gentle slope)
-  $\cos^{-1} \rightarrow$ blue (cool upper region)
-  $\tan^{-1} \rightarrow$ yellow (smooth curve)