

Precipitation Change in Beaufort, NC

[https://github.com/tarokatayama/KatayamaNgenziThornton_ENV872_
EDA_FinalProject.git](https://github.com/tarokatayama/KatayamaNgenziThornton_ENV872_EDA_FinalProject.git)

Karen Thornton, Lambert Ngenzi, Taro Katayama

Contents

1	Rationale and Research Questions	5
2	Dataset Information	6
3	Exploratory Analysis	7
4	Wrangle Data for Analysis	11
5	Analysis	22
6	Summary and Conclusions	25
7	Resources	26

List of Tables

1	1-year/24hr Events Over Year	13
2	Significant Events Over Year	19

List of Figures

1	Total Monthly Precipitation	11
2	Sig events (early decade)	14
3	Overall Precipitation (early decade)	15
4	Sig events (late decade)	17
5	Overall Precipitation (late decade)	18
6	Overall Precipitation by decade	20
7	1 year 24 hr storm event decade comparison	21
8	Decomposed time series	23

1 Rationale and Research Questions

Question 1: Has there been a significant increase in precipitation in Beaufort, NC from 1980 to 2016?

Question 2: Has there been a significant increase in 1-year/24h precipitation event in Beaufort, NC from decade to decade (1997 to 2006 and 2007 to 2016)?

Null Hypothesis 1: There is no significant change in precipitation from 1980 to 2016.

Null Hypothesis 2: There is no significant change in 1-year/24h precipitation events from decade to decade (1997 to 2006 and 2007 to 2016).

2 Dataset Information

Data was originally attained online from DAYMET for a Hydrology assignment during Fall Semester. Since then, DAYMET has been taken down, so the excel sheet from class was used. The data shows historical precipitation data from Beaufort, NC for the years 1980 to 2016. This dataset was chosen because it is the most complete and consistent precipitation dataset that we could find.

“Significant 1-year/24hr Precipitation Events” are considered 1 year/24h events using the NOAA threshold of 3.66 inches for Beaufort, NC. This means that in 24h the probability of it raining 3.66 inches or more should only happen once a year. These events are considered large storm events.

Get the working directory

```
# Get your working directory  
getwd()
```

```
# Load your datasets  
Beaufort_RAW<-read.csv("./Data/Raw/Beaufort_precip_1980-present_HUC_030203010503_dayMet_
```

Load all packages

```
# Load your packages  
#install.packages(webshot)  
library(webshot)  
library(tidyverse)  
library(dplyr)  
library(lubridate)  
#library(mapview)  
#mapviewOptions(fgb = FALSE)  
#library(sf)  
library(knitr)  
library(Kendall)  
library(zoo)  
library(gridExtra)  
library(formatR)  
options(scipen = 4) # limit number of digits
```

3 Exploratory Analysis

Initial Wrangling

```
knitr::opts_chunk$set(tidy.opts=list(width=50), tidy=TRUE)
```

```
#Structure of the dataset
```

```
str(Beaufort_RAW)
```

```
## 'data.frame': 13514 obs. of 5 variables:
```

```
## $ Date : Factor w/ 13514 levels "1980-01-01"
```

```
## $ Area.Weighted.Mean.Precipitation..mm.per.day.: num 0 0 0 12 9 0 0 7 9 0 ...
```

```
## $ year : int 1980 1980 1980 1980 1980 1980 1980
```

```
## $ month : int 1 1 1 1 1 1 1 1 1 1 ...
```

```
## $ day_of_month : int 1 2 3 4 5 6 7 8 9 10 ...
```

```
#Head of the dataset
```

```
head(Beaufort_RAW)
```

```
## Date Area.Weighted.Mean.Precipitation..mm.per.day. year month
```

```
## 1 1980-01-01 0 1980 1
```

```
## 2 1980-01-02 0 1980 1
```

```
## 3 1980-01-03 0 1980 1
```

```
## 4 1980-01-04 12 1980 1
```

```
## 5 1980-01-05 9 1980 1
```

```
## 6 1980-01-06 0 1980 1
```

```
## day_of_month
```

```
## 1 1
```

```
## 2 2
```

```
## 3 3
```

```
## 4 4
```

```
## 5 5
```

```
## 6 6
```

```
#Dimension of dataset
```

```
dim(Beaufort_RAW)
```

```
## [1] 13514 5
```

```
#Class of the dataset
```

```
class(Beaufort_RAW)
```

```
## [1] "data.frame"
```

```
# Column names of the dataset
```

```
colnames(Beaufort_RAW)
```

```
## [1] "Date"
```

```
## [2] "Area.Weighted.Mean.Precipitation..mm.per.day."
```

```
## [3] "year"
```

```
## [4] "month"
```

```
## [5] "day_of_month"
```

```
#summarize the data
```

```
summary(Beaufort_RAW)
```

```
##          Date          Area.Weighted.Mean.Precipitation..mm.per.day.
## 1980-01-01:      1      Min.      : 0.00
## 1980-01-02:      1      1st Qu.: 0.00
## 1980-01-03:      1      Median : 0.00
## 1980-01-04:      1      Mean    : 4.14
## 1980-01-05:      1      3rd Qu.: 2.00
## 1980-01-06:      1      Max.    :195.00
## (Other)      :13508      NA's    :9
##          year          month          day_of_month
## Min.      :1980      Min.      : 1.000      Min.      : 1.00
## 1st Qu.:1989      1st Qu.: 4.000      1st Qu.: 8.00
## Median :1998      Median : 7.000      Median :16.00
## Mean      :1998      Mean      : 6.522      Mean      :15.73
## 3rd Qu.:2007      3rd Qu.:10.000      3rd Qu.:23.00
## Max.      :2016      Max.      :12.000      Max.      :31.00
##
```

```
# Rename columns to logical names
```

```
names(Beaufort_RAW)[1] <- "Date"
```

```
names(Beaufort_RAW)[2] <- "Mean_Precip_mm"
```

```
#Set Date as a Date
```

```
Beaufort_RAW$Date<-as.Date(Beaufort_RAW$Date, "%Y-%m-%d")
```

```
# Set your ggplot theme
```

```
Theme <- theme_classic(base_size = 9) +  
  theme(axis.text = element_text(color = "black"),  
        legend.position = "right")  
theme_set(Theme)
```

```
#Save processed Raw to processed folder
```



```
write.csv(Beaufort_RAW, row.names = FALSE,
         file = "./Data/Processed/Beaufort_Processed.csv")

#Read in processed data
Beaufort_Processed<-read.csv("./Data/Processed/Beaufort_Processed.csv")
```

Create a Map of Beaufort for Visualization Aid

```
# Create a Beaufort point using long and lat
# mapviewOptions(fgb=FALSE)

# Beaufort_pt <- st_point(c(-76.6515, 34.7192))

# Convert the the two points into spatial data
# Beaufort_Pol <- st_sfc(Beaufort_pt, crs = 4326)

# view the map mapview(Beaufort_Pol)

# repeat the steps for the next layer
# Beaufort_Town <- st_as_sf(Beaufort_Pol)

# mapview(Beaufort_Town, col.regions = 'red')

# Load NC counties info NC_data <-
# st_read('./Data/Spatial/cb_2018_us_county_20m.shp')
# %>% filter(STATEFP == 37)

# Visualize MC data mapview(NC_data)

# Filter Beaufort county and map it
# Beaufort_County <- NC_data %>% filter(NAME %in%
# 'Carteret')

# mapview(Beaufort_County, col.regions =
# 'orange')

# Combine all map mapview(NC_data, alpha.regions
# = 0.2) + mapview(Beaufort_County, col.regions =
# 'orange') + mapview(Beaufort_Town, col.regions
# = 'red')

# plot in ggplot to put in presentation ggplot()+
# geom_sf(data = NC_data, aes(), alpha=0.2)+
# geom_sf(data = Beaufort_County, fill= 'orange',
```

```
# alpha = 0.7)+ geom_sf(data= Beaufort_Town, col  
# = 'red')+ ggtitle('Beaufort, NC')
```

4 Wrangle Data for Analysis

```
# Create a monthly mean precipitation and total  
# monthly precipitation dataset for Beaufort.  
# This dataset is only for visualization  
# purposes.  
Beaufort_Clean <- Beaufort_Processed %>%  
  group_by(year, month) %>%  
  summarise(meanmonthlyprecip = mean(Mean_Precip_mm),  
            sumMonthlyPrecip = sum(Mean_Precip_mm)) %>%  
  mutate(Date = my(paste0(month, "-", year)))  
  
# Plot total monthly precipitation data to see  
# rough trend (Clean dataset)  
ggplot(Beaufort_Clean, aes(x = Date, y = sumMonthlyPrecip)) +  
  geom_line() + labs(y = "Sum Monthly Precip. (mm)",  
                    caption = "Total Monthly Precipitation") + geom_smooth(method = lm)
```




Figure 1: Total Monthly Precipitation

Create a dataset for the early decade (1997-2006)

```
# This dataset has a 10 year time frame with
# precipitation in inches (1997-01-01 to
# 2006-12-31) and significant 1-year/24hr
# precipitation events in magnitude and number of
# significant precipitation events

Beaufort_early <- Beaufort_Processed %>%
  mutate(PrecipInches = Mean_Precip_mm * 0.0394) %>%
  filter(Date > ("1996-12-31"), Date < ("2007-01-01")) %>%
  mutate(sigPrecip = ifelse(PrecipInches > 3.66,
    PrecipInches, 0), NumSigPrecip = ifelse(PrecipInches >
    3.66, 1, 0)) %>%
  select(Date, year, month, day_of_month, PrecipInches,
    sigPrecip, NumSigPrecip) %>%
  drop_na()

# Create a Beaufort early dataset, where
# non-significant 1-year/24h precipitation events
# were classified as NA.
Beaufort_earlyNo0Precip <- Beaufort_Processed %>%
  mutate(PrecipInches = Mean_Precip_mm * 0.0394) %>%
  filter(Date > ("1996-12-31"), Date < ("2007-01-01")) %>%
  mutate(sigPrecip = ifelse(PrecipInches > 3.66,
    PrecipInches, NA), NumSigPrecip = ifelse(PrecipInches >
    3.66, 1, 0)) %>%
  select(Date, year, month, day_of_month, PrecipInches,
    sigPrecip, NumSigPrecip) %>%
  drop_na()

# Summary of number of 1-year/24hr events per
# year
Beaufort_early_summary <- Beaufort_early %>%
  group_by(year) %>%
  summarise(SigPrecipEvents = sum(NumSigPrecip))
```

```

# Create a table of number of 1-year/24hr events
# per year.
EarlyTable <- kable(Beaufort_early_summary, caption = "1-year/24hr Events Over Year",
  col.names = c("Year", "1-year/24hr event"))
EarlyTable

```

Table 1: 1-year/24hr Events Over Year

Year	1-year/24hr event
1997	0
1998	1
1999	1
2000	0
2001	0
2002	0
2003	0
2004	0
2005	1
2006	1

```

# Create a figure with number and magnitude of
# significant events per year.
Plot_early_sig <- ggplot(Beaufort_earlyNo0Precip, aes(x = Date,
  y = sigPrecip)) + geom_point() + ylim(c(0, 8)) +
  theme(axis.text.x = element_text(angle = 60, hjust = 1)) +
  labs(y = "1-year/24hr storm events (in)", x = "Date")
Plot_early_sig

```




Figure 2: Sig events (early decade)

```

# Create a figure showing the overall
# precipitation for the early decade
Beaufort_early$Date <- as.Date(Beaufort_early$Date,
    "%Y-%m-%d")

Plot_early_overall <- ggplot(Beaufort_early, aes(x = Date,
    y = PrecipInches)) + geom_point() + scale_x_date(date_labels = ("%m-%Y")) +
    labs(y = "Precipitation (in)", x = "Date")
Plot_early_overall

```




Figure 3: Overall Precipitation (early decade)

Create a dataset for the late decade (2007-2016)

```

# This dataset has a 10 year time frame,
# precipitation in inches (2007-01-01 to
# 2016-12-30) and significant 1-year/24hr
# precipitation events in magnitude and number of
# significant precipitation events

```

```

Beaufort_Late <- Beaufort_Processed %>%
  mutate(PrecipInches = Mean_Precip_mm * 0.0394) %>%
  filter(Date > "2006-12-31") %>%
  mutate(sigPrecip = ifelse(PrecipInches > 3.66,
    PrecipInches, 0), NumSigPrecip = ifelse(PrecipInches >
    3.66, 1, 0)) %>%
  select(Date, year, month, day_of_month, PrecipInches,
    sigPrecip, NumSigPrecip) %>%
  drop_na()

# Create a Beaufort late dataset, where non
# significant 1-year/24hr precipitation events
# were classified as NA.
Beaufort_LateNo0Precip <- Beaufort_Processed %>%
  mutate(PrecipInches = Mean_Precip_mm * 0.0394) %>%
  filter(Date > "2006-12-31") %>%
  mutate(sigPrecip = ifelse(PrecipInches > 3.66,
    PrecipInches, NA), NumSigPrecip = ifelse(PrecipInches >
    3.66, 1, 0)) %>%
  select(Date, year, month, day_of_month, PrecipInches,
    sigPrecip, NumSigPrecip) %>%
  drop_na()

# Summary of number of significant 1-year.24hr
# events per year (Late)
Beaufort_late_summary <- Beaufort_Late %>%
  group_by(year) %>%
  summarise(SigPrecipEvents = sum(NumSigPrecip))

```



```
# Create a figure with number and magnitude of
# significant 1-year/24hr events per year.
Plot_late_sig <- ggplot(Beaufort_LateNo0Precip, aes(x = Date,
  y = sigPrecip)) + geom_point() + ylim(c(0, 8)) +
  theme(axis.text.x = element_text(angle = 60, hjust = 1)) +
  labs(y = "1-year/24h storm events (in) ", x = "Date")
Plot_late_sig
```




Figure 4: Sig events (late decade)

```

# Create a figure showing the overall
# precipitation for the late decade
Beaufort_Late$Date <- as.Date(Beaufort_Late$Date, "%Y-%m-%d")

Plot_late_overall <- ggplot(Beaufort_Late, aes(x = Date,
  y = PrecipInches)) + geom_point() + scale_x_date(date_labels = ("%m-%Y")) +
  labs(y = "Precipitation (in)", x = "Date")
Plot_late_overall

```




Figure 5: Overall Precipitation (late decade)

```

# Create a table of number of significant
# 1-year/24hr events per year.
LateTable <- kable(Beaufort_late_summary, caption = "Significant Events Over Year",
  col.names = c("Year", "1-year/24hr event"))
LateTable

```

Table 2: Significant Events Over Year

Year	1-year/24hr event
2007	2
2008	1
2009	2
2010	2
2011	2
2012	0
2013	1
2014	2
2015	3
2016	2

Compare overall precipitation for each decade

```
grid.arrange(Plot_early_overall, Plot_late_overall,  
             ncol = 2)
```




Figure 6: Overall Precipitation by decade

Compare the 1-year/24 hour events for each decade

```
grid.arrange(Plot_early_sig, Plot_late_sig, ncol = 2)
```




Figure 7: 1 year 24 hr storm event decade comparison

5 Analysis

Perform t-test and seasonal Mann-Kendall for overall dataset

```
t.test(Beaufort_Processed$Mean_Precip_mm)

##
## One Sample t-test
##
## data: Beaufort_Processed$Mean_Precip_mm
## t = 43.492, df = 13504, p-value < 2.2e-16
## alternative hypothesis: true mean is not equal to 0
## 95 percent confidence interval:
## 3.953508 4.326684
## sample estimates:
## mean of x
## 4.140096

# significant (p-value=2.2e-16) <- This means
# that there is a significant change in
# precipitation over the years 1980 to 2016

# Using Seasonal Mann-Kendall to look at trend
# excluding seasonality. This will be a better
# indicator of significant change than the
# t-test. That being said, both tests were run.
Beaufort_RAW_2 <- Beaufort_Processed

# Used linear interpolation to fill missing data
# for precipitation data.
Beaufort_RAW_2$Mean_Precip_mm <- na.approx(Beaufort_RAW_2$Mean_Precip_mm)

# Created a time series analysis of the
# precipitation data at Beaufort.
firstday <- day(first(Beaufort_RAW_2$Date))
firstmonth <- month(first(Beaufort_RAW_2$Date))
firstyear <- year(first(Beaufort_RAW_2$Date))

Beaufort_TS <- ts(Beaufort_RAW_2$Mean_Precip_mm, start = c(firstyear,
  firstmonth, firstday), frequency = 365)

# Decomposed the time series to see components.
```

```
Beaufort_decompose <- stl(Beaufort_TS, s.window = "periodic")
plot(Beaufort_decompose)
```




Figure 8: Decomposed time series

```
# Ran a seasonal Mann Kendall to see if there is
# a change in precipitation over the time of the
# data frame. Used seasonal Mann-Kendall to
# exclude seasonality of precipitation.
Beaufort.trend <- Kendall::SeasonalMannKendall(Beaufort_TS)
Beaufort.trend
```

```
## tau = 0.0189, 2-sided pvalue =0.0056612
```

```
# Significant! (p-value = 0.0056612) <- This
# means that when you exclude seasonality there
# is still a significant change in precipitation
# from 1980 to 2016
```

T-test was run to compare the two decades

```
# Here we are looking to see if there is a change  
# in precipitation amount comparing two decades  
# (1999-2006) & (2007-2016)  
t.test(Beaufort_early$PrecipInches, Beaufort_Late$PrecipInches)  
  
##  
## Welch Two Sample t-test  
##  
## data: Beaufort_early$PrecipInches and Beaufort_Late$PrecipInches  
## t = -0.64906, df = 7133.8, p-value = 0.5163  
## alternative hypothesis: true difference in means is not equal to 0  
## 95 percent confidence interval:  
## -0.02812072 0.01413102  
## sample estimates:  
## mean of x mean of y  
## 0.1627598 0.1697546
```

```
# not significant (p-value=0.5163)<- meaning  
# precipitation amount not significantly  
# different over the two decades
```

```
# Here we compared the significant 1-year/24hr  
# precipitation events for the two decades  
# (1999-2006) & (2007-2016).  
t.test(Beaufort_early$sigPrecip, Beaufort_Late$sigPrecip)
```

```
##  
## Welch Two Sample t-test  
##  
## data: Beaufort_early$sigPrecip and Beaufort_Late$sigPrecip  
## t = -2.7068, df = 5451.7, p-value = 0.006815  
## alternative hypothesis: true difference in means is not equal to 0  
## 95 percent confidence interval:  
## -0.028681849 -0.004586864  
## sample estimates:  
## mean of x mean of y  
## 0.005537589 0.022171945
```

```
# significant! (p-value=0.006815) <- There are  
# more statistically significant 1 year  
# precipitation events in later decade.
```


6 Summary and Conclusions

After careful analysis, it has been concluded that there is a significant increase in precipitation from 1980 to 2016 in Beaufort, NC. When comparing decade 1 (1997 to 2006) to decade 2 (2007 to 2016) there was no significant increase in precipitation. That being said, when looking at significant 1-year/24hr storm events with a threshold of 3.66 inches there was a significant increase from decade 1 to decade 2.

While there isn't enough information to draw conclusions based on precipitation alone, our prediction is climate change as the driver behind the increase in 1-year/24hr storm events between decades. This project shows the importance of tracking large storm events in future climate change studies.

It would also be interesting to use a longer period of data (more than 30 years) to see if there is any new or different trends that could not be captured with our data.

7 Resources

NOAA Website https://hdsc.nws.noaa.gov/hdsc/pfds/pfds_map_cont.html?bkmrk=nc