

Drug Search and Physician Hazard:
An Investigation into Addict Behavior and Policy Remedies

Tate Mason*
University of Georgia
Athens, Georgia

October 19, 2025

Abstract

Keywords:

*Tate.Mason@uga.edu

1 Introduction

1.1 Background and Motivation

Over the last few decades, prescription drug abuse has become a significant and growing public health concern across the globe. The misuse of prescription drugs, specifically opioids, benzodiazepines, and stimulants, has led to the question of how to mitigate the practice of physician search. Physician search refers to the practice of patients seeking multiple doctors in the hope of "scoring" a prescription to continue their addiction. While there are preventative measures in place, like registries of offenders, the practice persists. Coinciding with this question, it would be of great use to ascertain the incentive for physicians to enable the misuse of drugs, gaining a repeat source of revenue.

This paper seeks to understand the interplay of addicts and physician search, in hope of policy remedies which are more effective than a list. Further, the analysis of prescriber responsibility is also of great importance. Being able to understand how morals and fiscal incentives contrast in this case could help to implement policy which negates the opportunity for prescribers to mis-prescribe a sensitive drug. Finally, this paper will look to gain an insight into potential rehabilitation remedies which may help take the onus off both addicts and physicians.

1.2 Problem Statement

To ascertain the interplay between physician and addict behavior, it would be of interest to model physician WTP, in a sense, with their reputation. That is, how much risk are physicians willing to onboard for the profits associated with increased and repeat prescriptions? This would be interesting for many reasons, since physicians have many reasons to uphold their standing from liability, licensure, and even word of mouth advertisement. The last one may be increased as addicts see a pathway to drug access, but that would also increase liability and risk of loss of their licensure.

For addicts, it is interesting to inquire about their search cost allocation, as well as habit based expenditure on health. Tolerance necessarily grows with prolonged use of drugs, thus modeling this sort of depletion of both search and health budgets is of interest to understand the impact on patients of finding a willing prescriber.

From these two avenues, I think there is a worthwhile question to answer. That is, what is the risk threshold physicians will take on in exchange for profit as well as understanding time and health costs applied to addicts once they find a willing prescriber.

1.3 Contributions

1.4 Paper Organization

2 Related Work

2.1 Previous Approaches

2.2 Limitations of Existing Work

3 Model

3.1 Addicts

There is a continuum of addicts in this model such that $\mathcal{A} = \{a_1, a_2, \dots, a_N\}$, each with an individual type $\gamma_i \in \{\gamma_l, \gamma_h\}$, corresponding to high or low predisposition to addiction. Each addict maximizes their utility over an infinite horizon, with discount factor β . Each addict also has a tolerance for risk of being cut off from search, r_i , which evolves according to their type and consumption behavior. Each period, they choose to search for a physician, and will successfully receive a prescription with probability $m(\theta)$, where θ is the measure of physicians willing to prescribe. Searching comes at a cost s_i , which depletes their health budget, and increases their risk of being cut off. Each period, their risk evolves according to $r'_i = r_i + \epsilon_a$, where $\epsilon_a \sim N(\mu_a, \sigma_a^2)$. Addicts optimize over the following: In the first period:

$$V_a^0(\gamma_i, r_i) = \{u(c_i) + \beta \mathbb{E}_t(1 - a(\theta))V_a(\gamma_i, r_i) + \beta \mathbb{E}_t a(\theta)V_a(\gamma_i, r'_i)\} \quad (1)$$

s.t.

$$u(c) = \frac{c_i^{1-\sigma}}{1-\sigma}$$

$$r'_i = r_i + \epsilon_a$$

In proceeding periods:

$$V_a^m(\gamma_i, r_i) = \max\{V_a^s(\gamma_i, r_i), V_a^f(\gamma_i, r'_i)\} \quad (2)$$

s.t.

$$V_a^s(\gamma_i, r_i) = \{u(c_i) + \beta V_a^m(\gamma_i, r'_i)\}$$

$$V_a^f(\gamma_i, r'_i) = \{u(c_i) + \mathbb{E}_t[\beta(1 - a(\theta))V_a(\gamma_i, r'_i) + \beta \mathbb{E}_t[a(\theta)V_a(\gamma_i, r'_i)]]\}$$

$$u(c) = \frac{c_i^{1-\sigma}}{1-\sigma}$$

$$r'_i = r_i + \epsilon_a$$

$$c_i = \gamma_i p_{it} + r_i^{1/\gamma_i} \times \sum s_{it}$$

Where V_a^0 is the value function for an addict in their first period of search, and V_a^m is the

value function for an addict in proceeding periods, if they choose to find a new doctor, V_a^f , or to stay with their current match, V_a^s . $u(c_i)$ is the utility from consumption, with CRRA parameter σ . Risk evolves as before, and consumption is a function of risk tolerance and type, with higher type addicts consuming more for a given risk level due to their predisposition. Addicts will continue to search until they are cut off, which happens when their risk exceeds a threshold set by physicians. Finally, the constraint on consumption is made up of prescription amount, p_{it} , risk tolerance, and search cost.

3.2 Doctors

There is a continuum of doctors such that $\mathcal{M} = \{m_1, m_2, \dots, m_K\}$ who value profit. They will exist in the model until they have overprescribed, taking on too much risk, $p_{jt} > r_{jt}$. Doctors can be matched with many addicts, with $p_{jt} = \sum_{i=1}^I c_{it}$, for all patients $I \subset \mathcal{A}$ matched to them. They will drop patients when their consumption needs threaten their risk constraint, thus allowing them to bring in a new, lower risk, patient. Doctors will prescribe to addicts with probability $m(\theta)$, where θ is the measure of physicians willing to prescribe. Each period, their risk evolves according to $r'_{jt} = r_{jt} + \epsilon_d$, where $\epsilon_d \sim N(\mu_d, \sigma_d^2)$.

The doctor's optimization is as follows

$$\begin{aligned} \max_{v_j} & v_j - i_j r_{jt} \\ \text{s.t.} & \\ v_j = & r_{jt}^2 \sum_{i=1}^I c_{it} + \xi \end{aligned} \tag{3}$$

Where v_j is a visit cost, which is initially a flat ξ and will evolve according to how many patients they prescribe to. i_j is a flat physician's insurance cost which is raised by amount of risk. This can be thought of as malpractice insurance, or something to that effect. Basically, they want to maximize visit revenue, net of their risk adjusted fixed cost.

Doctors who do not prescribe will have a different optimization, which is simply maximizing their revenue from visits, net of risk cost, which is 0 since they do not prescribe.

3.3 Estimation

To estimate, using method of simulated moments, I would need to get moments of the variance of risk for each type of γ_i , which I think I will classify as either low, average, or high. I am still working on this part, and will hope to have more in the next couple of weeks. I have been looking over literature using the Roy model since I think it makes the most sense to use for a simpler case like this. Hopefully, I will have more for this soon, but I think the bones are there. I am still just looking for more understanding of the estimation process.

3.4 Problem Formulation

3.5 Proposed Approach

4 Experimental Evaluation

4.1 Experimental Setup

4.2 Datasets

4.3 Results

4.4 Analysis

5 Discussion

5.1 Interpretation of Results

5.2 Limitations

5.3 Future Work