

Complete Record Edges, Path Holonomy, and Recovery-Correctable Readouts

Tau Core Technical Paper II (Series Paper III)

Jozsef Olcsak

Independent researcher

Draft v0.1 — Last revised: 11 August 2026

Abstract

We isolate a finite technical question arising in morphological readout frameworks: under what conditions can a source-side action separation be identified exactly with an observer-carrier Uhlmann chord? A reversible *complete record edge* is defined as a source-owned, trace- and dagger-preserving relation between occupied finite record algebras, equipped with one common action unit and no terminal bypass. Primitive-defect extensivity then forces an exact Trace–Gram law on the supported endpoint pair. Composable complete edges form a groupoid, but endpoint-only descent is valid only when relative holonomy is null under a source-frozen operational descriptor or belongs to a declared Uhlmann gauge.

We then separate this reversible theorem from irreversible record formation. A three-qubit bit-flip construction gives an explicit non-isomorphic CPTP map with an exact recovery on its occupied logical operator system. Monotonicity in both directions proves preservation of fidelity and hence of the Uhlmann chord on that system. By contrast, commuting records alone cannot distinguish reversible transport from dephasing. This yields a seven-part occupation certificate that adds either a coherent off-diagonal witness or a certified recovery test to independent action, tomography, and timing measurements. For nonexact recovery we derive an explicit Uhlmann-chord loss bound from a uniform trace-norm error, and hence from a full-code diamond-norm certificate. The results are representation and identifiability theorems. They do not prove that a physical parent source creates or occupies complete edges, that a laboratory carrier realizes the proposed common-action law, or that Tau Core is empirically validated. An upstream cyclic-support theorem can select the complete source packet inside the declared source-irredundant MVP class, but it does not by itself prove invertible edge transport, exact recovery, or finite action–chord identity. This is the third stage of a seven-stage foundation programme: Paper I fixes the architecture, Paper II isolates finite source sufficiency, the present paper asks what survives record transport, Paper IV isolates parent-law realization, Paper V develops temporal descent and Paper VI the quantum terminal. Paper VII is issued as the focused VII-A–VII-C triptych. The regular local clock factorization and GR proper-time limit are conditionally closed downstream of the record map; they neither strengthen exact recovery here nor prove physical edge occupation. In the common morphology-conditioned architecture, the record map is the finite resolution/acceptance stage after observer access and typed terminal precursors; recovery tests whether distinctions lost there remain correctable on the declared occupied support.

Dependency and ownership box.

Inputs from earlier papers: Paper I supplies the common morphology-conditioned readout architecture and finite action variable; Technical Paper I (series Paper II) supplies a frozen typed source signature and occupied support. Their unrestricted physical realization and occupation remain open.

Results owned here: sufficient complete-edge conditions for finite Trace–Gram transport; descriptor-frozen path holonomy; exact and approximate recovery statements for irreversible CP maps; the recovery-correctable toy model; and the FOC-7 operational certificate. This paper does not infer its source support from transport success. The downstream scalar-15/type-II ownership and functional-RG closure are imported only as branch status; an RG regulator is an analysis scheme and is not a complete record edge or an observer quantizer in the sense defined here.

Atemporal parent-realization terminology. Because the Tau parent is atemporal, phrases such as *Nature selects* do not denote an external agent or a later choice. Their precise meaning is the internal physical implication

$$(B_{\tau}^{\text{phys}}, s_U, \mathcal{L}_{\text{parent}}) \implies \mathfrak{P},$$

together with occupation of the support of the resulting packet $\mathfrak{P}$. A theorem inside a declared completion proves only class-relative realization. It does not prove this unrestricted base–seed arrow, and agreement of finitely many terminal readouts cannot by itself establish ambient-source exhaustivity. Current-facing statements therefore use *parent-law realization* and *physical occupation*; any retained legacy selection language is shorthand for this same distinction.

Contents

1	Question and claim boundary	4
2	Finite record packet	5
3	Record descent in the primitive-readout architecture	6
4	Complete source-owned record edges	7
5	Operational descriptor completeness	7
6	Paths and holonomy	8
7	Irreversible CP relations	8
8	A recovery-correctable CP toy model	10
9	Why commuting records are insufficient	11
10	Why recovery is the relevant physical distinction	11
11	A quantitative nonrecoverable dephasing control	12
12	Carrier classes and admissible claims	12
13	Proof and falsification dependency	12
13.1	Recovery-error budgets before timing	13
13.2	Frozen operational decision protocol	14

14 FOC-7 occupation certificate	14
15 What remains open	15
15.1 Later enriched-source status	15
16 Conclusion	15
A Fidelity convention	16




Figure 1: Dependency direction of the foundation sequence. Paper III starts from a frozen occupied source support; recovery or chord preservation cannot be used to infer that the physical parent selected it.

1 Question and claim boundary

Let a stabilized source/body construction assign a dimensionless finite action separation

$$x_A = \frac{\mathcal{X}_s}{\mathcal{A}_*}, \quad (1)$$

while an observer carrier assigns density operators ρ_u, ρ_v and the minimum-lift Uhlmann chord

$$x_Q = d_{U, \text{chord}}^2(\rho_u, \rho_v) = \min_U \|W_v - W_u U\|_{\text{HS}}^2 = 2(1 - \sqrt{F(\rho_u, \rho_v)}). \quad (2)$$

The target identity is $x_A = x_Q$. Equality of infinitesimal Hessians is not enough: a finite response can have the same tangent metric but differ by nonlinear terms. This paper therefore asks which *finite source relations* are sufficient for the identity, and how this statement changes for irreversible completely positive (CP) maps.

The status is deliberately narrow:

- complete-edge existence is an assumption about an occupied source relation, not a theorem that the unrestricted physical parent law realizes it;
- the Trace–Gram result is exact only on the declared occupied support;
- descriptor-null holonomy is operational and claim-relative, not an ontological erasure rule;
- the CP example is a recovery-correctable toy model, not an identified physical Tau carrier;
- no numerical constant, Standard-Model spectrum, or empirical anomaly is derived here.

Series notation ledger.




Figure 2: The finite physical fork. Exact Trace–Gram transport follows on a reversible complete edge. An irreversible CP relation requires a recovery theorem; otherwise contraction or an additional finite defect remains.

Symbol	Series-wide meaning	Primary treatment
s	source/seed packet prior to observer descent	Paper I
M_τ	stabilized morphological body	Paper I
Desc_O	observer-indexed physical descent/readout	Paper I
Π_{phys}	selected coherent source representation	Paper II
x_A	normalized finite source/body action separation	Papers I, III
x_Q	observer-carrier Uhlmann chord separation	Papers I, III
$\mathcal{C}, \mathcal{R}$	record map and recovery map	Paper III

2 Finite record packet

Let the occupied record support decompose into finitely many central blocks,

$$\mathcal{A}_Q = \bigoplus_{\alpha=1}^m M_{n_\alpha}(\mathbb{C}), \quad \tau(a) = \sum_{\alpha} p_{\alpha} \frac{\text{Tr}(a_{\alpha})}{n_{\alpha}}, \quad p_{\alpha} > 0, \quad \sum_{\alpha} p_{\alpha} = 1. \quad (3)$$

We assume a finite special dagger-Frobenius packet $(\mathcal{A}_Q, \mu, \eta, \Delta = \mu^\dagger, \epsilon = \eta^\dagger)$ with

$$\mu\Delta = I_{\mathcal{A}_Q}. \quad (4)$$

Specialness makes the primitive copy map isometric in the inherited trace metric. It is a strong normalization condition; a merely normalizable Frobenius algebra has $\mu\Delta = L$ for a positive central loop operator L , leaving a possible block-dependent scale.

Definition 1 (Finite recovery completeness). *A source law is finite recovery-complete on an occupied family when its finite action separation depends only on the source-owned minimum-lift chord,*

$$x_A = F(x_Q), \quad F(0) = 0, \quad (5)$$

and no additional positive stiffness remains on the declared source support while being invisible to that lift. This is a sufficient-statistic assumption for finite action separation, not a consequence of a positive local Hessian alone.

The admissible domain for exact extensivity is

$$\mathcal{D}_2 = \{(Y_1, Y_2) : Y_i \geq 0, Y_1 + Y_2 \leq 2, Y_1, Y_2 \text{ arise from admitted orthogonal source slots}\}. \quad (6)$$

The upper bound follows from the squared chord convention. Extensivity means

$$F(Y_1 + Y_2) = F(Y_1) + F(Y_2), \quad (Y_1, Y_2) \in \mathcal{D}_2, \quad (7)$$

with one universal continuous F on the connected occupied orbit. Merely assigning separate functions F_α to blocks would not imply a common finite law.

Lemma 1 (Trace–Gram response on the admissible domain). *If Eq. (5) is continuous, has local unit $F'(0) = 1$, and satisfies Eq. (7) on a connected admissible interval, then $F(Y) = Y$ throughout that interval.*

Proof. The restricted Cauchy equation gives rational homogeneity on every admissible subinterval. Continuity extends it to real arguments. Hence $F(Y) = cY$, and $F'(0) = 1$ fixes $c = 1$. $\square$

The assumptions are material. The family

$$F_\lambda(Y) = Y + \lambda Y^2 \quad (8)$$

has the same local unit but composition defect

$$F_\lambda(Y_1 + Y_2) - F_\lambda(Y_1) - F_\lambda(Y_2) = 2\lambda Y_1 Y_2. \quad (9)$$

Thus local metric agreement does not establish finite A8b.

3 Record descent in the primitive-readout architecture

The common architecture of Paper I is

$$Y_O^{(a)} = Q_O^{(a)} \circ T^{(a)} \circ A_O[M_\tau]. \quad (10)$$

The present paper studies the finite record realization of the final stage, not an independent physical layer. Observer position, extent and relational access are already contained in A_O ; $T^{(a)}$ supplies a typed continuous precursor; and $Q_O^{(a)}$ supplies stable operational classes. Consequently a recovery map is meaningful only on a declared state family, operator system, code subspace or correctable algebra inherited from the same body/access packet.

This distinction separates the primitive readouts. For time, ordered changes between distinct stable $Q_O^{(T)}$ -classes are candidate ticks, while the positive clock form determines duration. For a quantum-capable readout, several configurations may occupy one ordinary resolved class while a source-owned phase terminal retains their relative holonomy and coherent composition. Noninjectivity or finite capacity alone yields only classical coarse graining. Gravity, matter, radiation and gauge records likewise require their own typed precursors before the record edge acts.

The stacked precursor rank remains bounded by the common observer access:

$$\text{rank} \begin{bmatrix} D(T^{(1)} \circ A_O) \\ \vdots \\ D(T^{(n)} \circ A_O) \end{bmatrix} \leq \text{rank } A_O. \quad (11)$$

Thus preserving several named records does not by itself prove preservation of independent source information. Complete-edge and recovery theorems below remain conditional on physical occupation of this common typed support.

4 Complete source-owned record edges

Definition 2 (Complete record edge). *Let u, v be occupied fibers with finite special dagger-Frobenius record algebras $(\mathcal{A}_{Q,u}, \tau_u)$ and $(\mathcal{A}_{Q,v}, \tau_v)$. A source relation $e : u \rightarrow v$ is a complete record edge on a declared support when its realization:*

- (E1) *is an invertible multiplication-, unit- and dagger-intertwiner;*
- (E2) *preserves the inherited metric, faithful trace and common action unit;*
- (E3) *transports the full declared endpoint record pair;*
- (E4) *is source-owned and contains no terminal-only bypass copy; and*
- (E5) *belongs to the same primitive-extensive occupied edge orbit.*

This definition characterizes sufficient structure. It does not derive edge existence or invertibility.

Proposition 1 (Exact finite identity on a complete edge). *On a finite recovery-complete, primitive-extensive packet, every occupied complete edge restricts to a faithful trace-preserving $*$ -isomorphism*

$$\Pi_e : \mathcal{A}_{Q,u} \rightarrow \mathcal{A}_{Q,v}, \quad \tau_v(\Pi_e(a)) = \tau_u(a), \quad (12)$$

whose standard-form lift is isometric. For its supported endpoint pair,

$$\boxed{\frac{\mathcal{X}_s(e)}{\mathcal{A}_*} = \min_U \|W_v - W_u U\|_{\text{HS}}^2 = d_{U, \text{chord}}^2(\rho_u, \rho_v).} \quad (13)$$

Proof. The algebraic intertwiners and invertibility give a faithful $*$ -isomorphism. Metric and common-unit transport exclude a rescaled trace. Therefore

$$\langle L^2(\Pi_e)a, L^2(\Pi_e)b \rangle_v = \tau_v(\Pi_e(a)^\dagger \Pi_e(b)) \quad (14)$$

$$= \tau_v(\Pi_e(a^\dagger b)) = \tau_u(a^\dagger b), \quad (15)$$

so the standard-form lift is isometric. Source ownership places the amplitude residual in the once-counted primitive defect ledger. Finite recovery completeness and the previous lemma then force $x_A = x_Q$. $\square$

A rescaled trace, a terminal-only amplitude copy, or the nonlinear response (8) violates a distinct hypothesis. The theorem therefore packages a strong but auditable sufficient realization rather than showing that every stable record process has this form.

5 Operational descriptor completeness

Holonomy cannot be declared null by inspecting density matrices alone. The descriptor must be fixed before paths are compared.

Definition 3 (Claim-complete operational descriptor). *For an occupied fiber u , a descriptor protocol $\mathfrak{D}_u^{(C)}$ is complete relative to claim C when it freezes:*

- (D1) *the occupied operator system $\mathcal{S}_{\text{occ}}$;*
- (D2) *an informationally complete effect frame on that system, or an explicit restricted-observable label;*
- (D3) *estimators for the retained action and metric data;*

(D4) every typed terminal and calibration used by C ;

(D5) the admitted gauge subgroup and path family;

(D6) equivalence margins δ_j , confidence level and multiple-test correction.

For two path outputs x_p, x_q , descriptor-nullity is accepted only by a predeclared equivalence test,

$$\text{CI}_{1-\alpha}[D_j(x_p) - D_j(x_q)] \subseteq [-\delta_j, \delta_j] \quad \text{for every frozen component } j. \quad (16)$$

Failure to reject equality is not evidence of nullity. Completeness here is claim-relative and never means access to all parent information.

If the effect frame is informationally complete on $\mathcal{S}_{\text{occ}}$, equality of all ideal probabilities identifies the state on that system. Finite data replace equality by Eq. (16). A central automorphism is not automatically gauge: block permutations, weights and typed roles remain physical whenever the frozen descriptor detects them.

6 Paths and holonomy

Composable complete edges form a groupoid $\mathcal{G}_{\text{rec}}$. A path $p = e_n \circ \dots \circ e_1 : u \rightarrow v$ induces $\Pi_p = \Pi_{e_n} \circ \dots \circ \Pi_{e_1}$. For two paths $p, q : u \rightarrow v$, define relative holonomy

$$H_{p,q} = \Pi_q^{-1} \Pi_p. \quad (17)$$

Corollary 1 (Pathwise identity and endpoint descent). *Every complete path obeys Eq. (13) pathwise. An endpoint-only law descends on a declared occupied family exactly when $H_{p,q}$ belongs to the frozen Uhlmann gauge or is null under the claim-complete descriptor. Otherwise the path class is retained physical data.*

Proof. Faithful trace-preserving $*$ -isomorphisms are closed under composition and inversion. Right-unitary Uhlmann gauge leaves $WW^\dagger$ and the minimized chord invariant. Any descriptor-visible relative automorphism distinguishes the two transported records and cannot consistently be quotiented. $\square$

For a concrete control, take three fibers carrying $M_2(\mathbb{C})$, with normalized trace, and edges

$$\Pi_e = \text{Ad}_X : u \rightarrow v, \quad \Pi_f = \text{Ad}_X : v \rightarrow w, \quad \Pi_g = \text{Ad}_Z : u \rightarrow w. \quad (18)$$

For $p = f \circ e$ and $q = g$, $\Pi_p = \text{id}$ while $H_{p,q} = \text{Ad}_Z$. On diagonal densities and effects this holonomy is null. If $|+\rangle\langle+|$ and an X -basis effect are frozen into the descriptor, Ad_Z maps $|+\rangle$ to $|-\rangle$ and becomes visible. The null decision therefore belongs to the predeclared operational claim, not to an analyst's post-hoc choice.

7 Irreversible CP relations

A generic detector, decohering evolution or rank-reducing instrument is not a complete edge. If $\mathcal{C}$ is CPTP, fidelity monotonicity gives

$$F(\mathcal{C}(\rho), \mathcal{C}(\sigma)) \geq F(\rho, \sigma), \quad d_U^2(\mathcal{C}(\rho), \mathcal{C}(\sigma)) \leq d_U^2(\rho, \sigma). \quad (19)$$

Exact finite A8b can therefore be lost. It is restored on an occupied domain if the map is exactly recoverable there. The domain must be typed:

Definition 4 (Recovery domains). *Let $P\mathcal{H}$ be a code subspace. We distinguish:*

Three-vertex descriptor-frozen holonomy control




Figure 3: A three-vertex control. The same relative automorphism is null on a diagonal record family but visible once a coherent X -basis witness is frozen into the endpoint descriptor.

(R1) a state family $\mathfrak{S} \subseteq \mathcal{D}(P\mathcal{H})$;

(R2) a unital self-adjoint linear operator system $\mathcal{S}_{\text{op}} \subseteq \mathcal{B}(P\mathcal{H})$;

(R3) a correctable unital $*$ -subalgebra $\mathcal{A}_{\text{corr}} \subseteq \mathcal{B}(P\mathcal{H})$; and

(R4) the full code algebra $\mathcal{B}(P\mathcal{H})$.

State-family recovery means equality only on $\mathfrak{S}$. Operator-system recovery means equality of the linear maps on $\mathcal{S}_{\text{op}}$. Algebra recovery preserves all states and observables encoded by $\mathcal{A}_{\text{corr}}$. Full-code recovery is the special case $\mathcal{A}_{\text{corr}} = \mathcal{B}(P\mathcal{H})$. None of these scopes may be silently enlarged after the test.

Proposition 2 (Recovery sandwich). *Let $\mathcal{C}$ and $\mathcal{R}$ be CPTP maps satisfying*

$$(\mathcal{R} \circ \mathcal{C})(\rho) = \rho \quad (20)$$

for every state in a frozen family $\mathfrak{S}$. Then for every pair $\rho, \sigma \in \mathfrak{S}$,

$$F(\mathcal{C}(\rho), \mathcal{C}(\sigma)) = F(\rho, \sigma), \quad d_U^2(\mathcal{C}(\rho), \mathcal{C}(\sigma)) = d_U^2(\rho, \sigma). \quad (21)$$

Proof. Monotonicity under $\mathcal{C}$ gives one inequality. Applying monotonicity under $\mathcal{R}$ and using Eq. (20) gives the reverse inequality. Equality of fidelity implies equality of the chord convention in Eq. (2). $\square$

This statement is closely related to quantum error-correction conditions [1] and to the broader sufficiency/recovery viewpoint [2], but only the explicit recovery identity is needed below.

Proposition 3 (Explicit approximate-recovery chord bound). *Let $\mathfrak{S}$ be a frozen state family and suppose*

$$\sup_{\rho \in \mathfrak{S}} \|(\mathcal{R}\mathcal{C})(\rho) - \rho\|_1 \leq \eta. \quad (22)$$

This follows, in particular, from $\|\mathcal{R}\mathcal{C} - \text{id}\|_\diamond \leq \eta$ on the full code algebra. For $\rho, \sigma \in \mathfrak{S}$, define

$$x_{\text{in}} = d_U^2(\rho, \sigma), \quad x_{\text{out}} = d_U^2(\mathcal{C}\rho, \mathcal{C}\sigma). \quad (23)$$

Then

$$0 \leq x_{\text{in}} - x_{\text{out}} \leq x_{\text{in}} - [\max\{0, \sqrt{x_{\text{in}}} - 2\sqrt{\eta}\}]^2. \quad (24)$$

Equivalently,

$$x_{\text{in}} - x_{\text{out}} \leq \begin{cases} x_{\text{in}}, & x_{\text{in}} \leq 4\eta, \\ 4\sqrt{\eta x_{\text{in}}} - 4\eta, & x_{\text{in}} > 4\eta. \end{cases} \quad (25)$$

Proof. Write $d(\rho, \sigma) = \sqrt{d_U^2(\rho, \sigma)}$, the Bures chord metric. The Fuchs–van de Graaf inequality [3] gives

$$d(\rho, \mathcal{R}\mathcal{C}\rho)^2 = 2(1 - \sqrt{F(\rho, \mathcal{R}\mathcal{C}\rho)}) \leq \|\rho - \mathcal{R}\mathcal{C}\rho\|_1 \leq \eta, \quad (26)$$

and likewise for σ . The triangle inequality and contractivity under $\mathcal{R}$ imply

$$\sqrt{x_{\text{in}}} \leq \sqrt{x_{\text{out}}} + 2\sqrt{\eta}. \quad (27)$$

Contractivity under $\mathcal{C}$ also gives $x_{\text{out}} \leq x_{\text{in}}$. Solving these two inequalities for x_{out} yields Eqs. (24) and (25). $\square$

The bound is useful only relative to the frozen domain. It becomes weak when $x_{\text{in}} \lesssim 4\eta$, precisely where recovery error can hide the full input separation. It still concerns only the Uhlmann side; it does not make the once-counted source action equal to either chord.

The estimate is exact in the limit $\eta = 0$ and becomes vacuous in the first branch when $x_{\text{in}} \leq 4\eta$. It is a uniform worst-case consequence of trace-norm recovery plus metric contractivity; no claim of universal optimality is made for a specified channel, code, or state pair. A smaller state-dependent error may give a sharper bound, but it must be certified on the same predeclared state family. A diamond-norm certificate applies to the full code including reference systems, whereas the state-family hypothesis in Eq. (22) must not be silently promoted to that stronger scope.

8 A recovery-correctable CP toy model

Let $\mathcal{H}_L = \mathbb{C}^2$, $\mathcal{H}_P = (\mathbb{C}^2)^{\otimes 3}$, and define the encoding isometry

$$V|0\rangle = |000\rangle, \quad V|1\rangle = |111\rangle. \quad (28)$$

Let X_i flip physical qubit i , $P_0 = VV^\dagger$, and $P_i = X_i P_0 X_i$. These four syndrome subspaces are mutually orthogonal. For $0 \leq p \leq 1$, define

$$\mathcal{C}_p(\rho) = (1-p)V\rho V^\dagger + \frac{p}{3} \sum_{i=1}^3 X_i V \rho V^\dagger X_i. \quad (29)$$

This is CPTP and, for $0 < p < 1$, is not a $*$ -isomorphism from the logical matrix algebra onto its noisy image.

Define a decoder by syndrome measurement and correction,

$$\mathcal{R}_{\text{dec}}(\sigma) = V^\dagger \left[P_0 \sigma P_0 + \sum_{i=1}^3 X_i P_i \sigma P_i X_i \right] V + \text{Tr}(P_{\text{fail}} \sigma) |0\rangle\langle 0|, \quad (30)$$

where $P_{\text{fail}} = I - P_0 - \sum_i P_i$ makes the map trace preserving outside the occupied range.

Proposition 4 (Exact recovery-correctable non-isomorphic edge). *For every logical state ρ ,*

$$\mathcal{R}_{\text{dec}} \circ \mathcal{C}_p(\rho) = \rho. \quad (31)$$

Consequently Eq. (21) holds for all logical state pairs and all $p \in [0, 1]$.

Proof. Each summand of Eq. (29) lies in one orthogonal syndrome subspace. The corresponding term in Eq. (30) projects onto that subspace, applies the matching correction, and decodes with $V^\dagger$. Every branch therefore returns its probability weight times ρ ; the weights sum to one. The recovery-sandwich proposition gives fidelity and chord equality. $\square$

The toy model recovers the full logical algebra $\mathcal{B}(\mathcal{H}_L) = M_2(\mathbb{C})$, not merely a commuting state family. It proves that invertibility of the physical CP map is unnecessary for exact finite metric transport on a certified code. It does not show that the body action assigns the same x_A , nor that this code is a physical Tau carrier.

9 Why commuting records are insufficient

Let $\mathcal{D} \subset M_2(\mathbb{C})$ be the diagonal algebra and compare

$$\Pi_{\text{rev}} = \text{id}, \quad \Pi_{\text{irr}}(a) = P_0 a P_0 + P_1 a P_1. \quad (32)$$

Both maps agree on every diagonal density and diagonal effect. Yet Π_{irr} destroys off-diagonal coherence and has no CPTP inverse on the full matrix algebra.

Proposition 5 (Commuting-record reversibility no-go). *No protocol restricted to the commuting record algebra $\mathcal{D}$ can identify whether the occupied process is Π_{rev} or Π_{irr} . Reversibility requires either a coherent witness outside $\mathcal{D}$ or a certified recovery test on a larger predeclared operator system.*

Proof. The two channels induce identical probabilities for every state and effect in $\mathcal{D}$. Therefore every experiment confined to that algebra has the same likelihood under both hypotheses. An off-diagonal state/effect or a recovery test separates them. $\square$

10 Why recovery is the relevant physical distinction

The preceding results separate three statements that are often conflated. First, an output may retain enough classical statistics to reproduce a chosen endpoint table. Second, a channel may preserve all pairwise state distances on a declared family. Third, a physical operation may admit a recovery on the full operator system carrying that family. Only the third statement certifies that the apparently retained information was not silently converted into a smaller commuting record.

This distinction matters for a morphological readout proposal because the target identity $x_A = x_Q$ concerns a finite state geometry, not merely one probability distribution. If the source-side action distinguishes coherent directions that the observer process dephases, then a diagonal endpoint test can report agreement while the claimed common geometry has already been lost. The recovery domain therefore belongs to the physical claim:

$$\text{action support} \supseteq \mathcal{S}_{\text{occ}} \xrightarrow{\mathcal{C}} \mathcal{C}(\mathcal{S}_{\text{occ}}) \xrightarrow{\mathcal{R}} \mathcal{S}_{\text{occ}}. \quad (33)$$

If the last arrow is certified only on a state family, no conclusion about an unmeasured coherent observable follows. If it holds on a correctable algebra, all encoded states and effects of that algebra are retained. This is why the paper keeps state-family, operator-system, algebra and full-code claims separate rather than using “information preserving” as an untyped phrase.

11 A quantitative nonrecoverable dephasing control

Consider the qubit dephasing family

$$\mathcal{D}_\lambda(\rho) = \frac{1+\lambda}{2}\rho + \frac{1-\lambda}{2}Z\rho Z, \quad 0 \leq \lambda \leq 1. \quad (34)$$

It fixes every diagonal state. For the coherent pair

$$\rho_+ = |+\rangle\langle+|, \quad \rho_- = |-\rangle\langle-|, \quad (35)$$

the input states are orthogonal and hence

$$x_{\text{in}} = 2. \quad (36)$$

Their outputs have Bloch vectors $\pm\lambda\hat{x}$. The squared fidelity is

$$F(\mathcal{D}_\lambda\rho_+, \mathcal{D}_\lambda\rho_-) = 1 - \lambda^2, \quad (37)$$

so the output chord is

$$x_{\text{out}}(\lambda) = 2 \left(1 - \sqrt{1 - \lambda^2}\right). \quad (38)$$

Thus $x_{\text{out}}(0) = 0$, $x_{\text{out}}(1) = 2$, and every $0 \leq \lambda < 1$ strictly contracts the coherent separation. Meanwhile all diagonal preparations and effects are exactly independent of λ .

Proposition 6 (Continuous hidden-contraction family). *No descriptor restricted to the diagonal algebra can estimate the coherent chord loss in Eq. (38). A single predeclared X -coherence witness identifies λ and therefore the lost finite separation.*

Proof. The restriction of $\mathcal{D}_\lambda$ to the diagonal algebra is the identity for every λ , proving non-identifiability there. On the input ρ_+ , the expectation of X is $\text{Tr}[X\mathcal{D}_\lambda(\rho_+)] = \lambda$. Substitution into Eq. (38) reconstructs the chord. $\square$

This control is deliberately simpler than the three-qubit recovery example. Together they bracket the decision: non-isomorphic and noisy need not mean information-losing on a code, while perfect agreement on a classical record need not mean preservation of the occupied quantum geometry.

12 Carrier classes and admissible claims

The theorem does not require one preferred hardware platform. It requires a carrier whose operational scope matches the claimed recovery scope. Table 1 lists the minimum evidence for broad candidate classes; none is asserted to be a Tau carrier.

The table also clarifies effect size. A QOR timing test is meaningful only after the finite geometry has been certified with uncertainty smaller than the separation under test. If η_{rec} makes the upper bound in Eq. (25) comparable to x_{in} , then the carrier cannot distinguish a Tau-specific finite identity from ordinary channel contraction, regardless of timing precision.

13 Proof and falsification dependency

The logical order of the paper is finite:

typed occupied support $\Downarrow$ complete edge or certified recovery $\Downarrow$ Uhlmann-chord preservation $\Downarrow$ plus independent source action $x_A = x_Q$ $\Downarrow$ plus held record and ΔH QOR timing test.	(39)
--	------

Candidate class		Natural record support	Missing decisive certificate
Superconducting spin qubit	or	Full two-level operator system with controllable coherent bases	Independent source-action reconstruction and certified recovery/decoherence budget
Trapped ion or metastable atom	or	Long-lived internal states with single-shot discrimination	Same-carrier action curvature and held-record timing on a frozen protocol
Photon polarization/path		High-quality tomography and explicit path holonomy	Metastable one-record hold and separation of propagation loss from source geometry
Macroscopic record	pointer	Stable, redundant commuting outcomes	Coherent witness or recovery on a larger operator system; diagonal stability alone is insufficient

Table 1: Candidate carrier classes and the evidence still needed before a finite record packet can be promoted. Familiar hardware supplies controls, not the Tau-specific action–Uhlmann identity.

Target chord x_*	Recovery error η	Maximum chord loss	Fractional loss
1.00	10^{-2}	0.36	36.0%
1.00	10^{-3}	0.1225	12.25%
0.25	10^{-3}	0.0592	23.68%
0.25	10^{-4}	0.0196	7.84%

Table 2: Worst-case losses from Eq. (25). Smaller target separations demand substantially stronger recovery certificates.

Failure at an earlier row cannot be repaired by a successful later timing fit. Conversely, failure of exact recovery need not reject every finite readout model: it selects the approximate branch and requires the explicit loss ledger of Eq. (24). The decisive falsification targets are therefore typed: failure of coherent/recovery evidence rejects complete transport on the tested support; failure of $x_A = x_Q$ rejects the common-action identification; and a timing violation rejects the occupied QOR completion only after both upstream certificates have passed.

13.1 Recovery-error budgets before timing

For a target input chord x_* , define the worst-case unresolved fractional loss

$$r_{\text{loss}}(x_*, \eta) = \frac{x_* - [\max\{0, \sqrt{x_*} - 2\sqrt{\eta}\}]^2}{x_*}. \quad (40)$$

The recovery certificate is informative at fractional tolerance r_0 only if $r_{\text{loss}}(x_*, \eta) \leq r_0$. In the nonvacuous branch this is equivalent to

$$\frac{4\sqrt{\eta x_*} - 4\eta}{x_*} \leq r_0. \quad (41)$$

This inequality must be checked before a timing run. It prevents a small observed body–state mismatch from being interpreted as new temporal physics when ordinary channel uncertainty already permits the full discrepancy.

For illustration, Table 2 evaluates the certified worst-case loss. These are dimensionless mathematical controls, not predicted experimental errors.

The scaling is important: at fixed η , the relative uncertainty grows as the target chord decreases. A laboratory proposal should therefore not begin with the smallest technically resolvable separation. It should first demonstrate the common-action identity at two or more

separations for which the recovery uncertainty is decisively subdominant, and only then test the downstream timing relation.

13.2 Frozen operational decision protocol

A reproducible decision can be organized without assuming the Tau conclusion:

- (P1) Declare the state family or operator system, the source-action estimator and the allowed gauge before collecting endpoint data.
- (P2) Characterize $\mathcal{C}$ and either certify a recovery map or freeze a coherent witness that detects departure from the commuting restriction.
- (P3) Choose at least two nonzero target separations satisfying Eq. (41) at the declared tolerance.
- (P4) Reconstruct x_A and x_Q independently and perform an equivalence test, not a failure-to-reject test, for the same-action identity.
- (P5) Only after P1–P4 pass, measure ΔH and the held-record timing endpoint used in the QOR inequality.
- (P6) Run dephasing, drift, detector and nonlinear-response controls under the same analysis freeze.

The corresponding verdicts are asymmetric. Failure of P2 demotes the carrier scope; failure of P4 rejects finite A8b on that scope; failure of P5 rejects the occupied timing completion if the upstream certificates remain valid. Passing all steps identifies one finite carrier packet within the tested model class, not the complete Tau parent ontology.

14 FOC-7 occupation certificate

The preceding no-go upgrades the finite occupation certificate. A candidate carrier passes *FOC-7* only if one predeclared protocol establishes:

- (F1) a metastable, held, single-shot record with a frozen discrimination error;
- (F2) independent source-side reconstruction of x_A , without tomography or timing data;
- (F3) independent tomographic reconstruction of x_Q ;
- (F4) $x_A = x_Q$ within predeclared equivalence margins at at least two nonzero separations;
- (F5) either a coherent off-diagonal witness distinguishing the complete operator system from its commuting restriction, or a certified recovery map satisfying $\|\mathcal{RC} - \text{id}\|_\diamond \leq \eta_{\text{rec}}$ on that system;
- (F6) downstream timing with independently measured energy spread; and
- (F7) controls for decoherence, drift, detector response and nonlinear finite corrections.

Passing FOC-7 would identify the tested finite packet within its frozen model class. It would not by itself prove the entire parent ontology. Failure at F5 specifically prevents commuting data from being promoted into evidence for a reversible complete edge.

15 What remains open

15.1 Later enriched-source status

Inside the primary source-irredundant enriched completion, the traced dagger-Frobenius two-leg record source, canonical represented trace and primitive-extensive finite Trace–Gram law are now constructed conditionally on the occupied complete ROOT path packet. Thus their internal construction is no longer the principal gap. The cyclic generated-support criterion, together with nonzero record unfolding and actual stable-record existence, now selects complete source-packet occupation inside the declared source-irredundant MVP carrier class. This does not yet supply the stronger transport hypotheses of this paper: complete-edge invertibility, descriptor-complete holonomy descent, or exact recovery on a stated operator system must still be certified. Moreover, an inequivalent energy-sharing ownership completion has the same current narrow source reduct, so transport or recovery success cannot identify the unrestricted physical parent retroactively.

The later scalar-15/type-II branch does not change this transport boundary. Its class-relative base–seed reconstruction closes which source packet owns the selected branch, and its exact functional RG equation closes how that packet is evaluated across scales. Neither statement proves reversible record transport, descriptor-complete holonomy descent or exact recovery. In particular, the smooth RG cutoff must not be identified with the observer-source finite-resolution map: the former is a removable calculation scheme for an exact flow, whereas the latter defines operational readout classes and must be physically sourced.

The internal mathematics now distinguishes three cases:

$$\begin{aligned} \text{complete reversible edge} &\Rightarrow \text{exact finite Trace–Gram,} \\ \text{recovery-correctable CP edge} &\Rightarrow \text{exact chord on the code,} \\ \text{generic irreversible CP edge} &\Rightarrow \text{contraction or extra defect.} \end{aligned} \tag{42}$$

The decisive physical question is not settled by these implications:

Does the physical base–seed source create and occupy either the complete dagger-Frobenius edge packet or a recovery-correctable non-isomorphic packet, and does its once-counted action assign the same finite separation as the observer amplitude geometry?

The represented normalized trace and primitive extensivity are supplied conditionally by the enriched traced source packet, rather than by local Hessian positivity alone. Their unrestricted physical occupation remains an input. Finite recovery completeness still says that the chord is a sufficient statistic for action separation and remains separately falsifiable outside the complete occupied path class.

Equation (24) controls approximate recovery on the frozen state family. It is not an exact A8b theorem: the finite source action remains independent, and experiment-specific promotion still requires a declared code, error model and reconstruction uncertainty.

16 Conclusion

The subsequent quantum-terminal closure also fixes the scope of the transport results. For a logical source component separated from its inaccessible complement by the occupied source hypergraph, a disjoint-source valuation forces a block-diagonal parent Hessian and the complement carries no logical input information; exact recovery then follows. A crossing occupied hyperedge supplies the permitted interaction boundary, where entanglement, leakage and local decoherence may occur while the joint state remains represented. This does not assert unrestricted parent-law realization and physical occupation of the valuation: nonfaithful terminal

descent can hide a parent cross term, so complete-edge or recovery success on a restricted code cannot establish global parent additivity.

Complete source-owned record edges provide a concise sufficient theorem for finite action–Uhlmann identity, while path holonomy states exactly when that identity can descend to endpoint-only data. The operational descriptor makes the null decision testable and prevents post-hoc removal of visible path information.

Irreversibility does not end the route: the three-qubit construction proves that a non-isomorphic CP map can preserve the Uhlmann chord on a certified recoverable code. But commuting record agreement is insufficient to decide between this structure, a reversible edge, and dephasing. FOC-7 therefore requires coherent or recovery evidence in addition to action, tomography and timing measurements.

The paper closes the internal finite alternatives, not their unrestricted parent-law realization. Physical source occupation, common-action ownership and a laboratory carrier remain open empirical and source-theoretic problems.

This closes the series-level handoff without reversing it: architecture precedes source identification, and source identification precedes transport. An exact recovery theorem can preserve an occupied packet, but it cannot prove that the physical base–seed parent law generates, realizes and occupies that packet in the first place.

A Fidelity convention

We use squared Uhlmann fidelity

$$F(\rho, \sigma) = \left[\text{Tr} \sqrt{\sqrt{\rho} \sigma \sqrt{\rho}} \right]^2. \quad (43)$$

The minimum Hilbert–Schmidt purification chord is

$$\min_U \|W_\rho - W_\sigma U\|_{\text{HS}}^2 = 2(1 - \sqrt{F}), \quad (44)$$

following Uhlmann’s purification geometry [4]. This is not the squared Bures distance convention used in every part of the literature; all equalities in this paper use the displayed definition.

References

- [1] Emanuel Knill and Raymond Laflamme. Theory of quantum error-correcting codes. *Physical Review A*, 55:900–911, 1997.
- [2] Dénes Petz. Sufficient subalgebras and the relative entropy of states of a von neumann algebra. *Communications in Mathematical Physics*, 105:123–131, 1986.
- [3] Christopher A. Fuchs and Jeroen van de Graaf. Cryptographic distinguishability measures for quantum-mechanical states. *IEEE Transactions on Information Theory*, 45(4):1216–1227, 1999.
- [4] Armin Uhlmann. The transition probability in the state space of a *-algebra. *Reports on Mathematical Physics*, 9(2):273–279, 1976.