

5. Linear Programming

Roberto Amadini

Department of Computer Science and Engineering, University of Bologna, Italy

Combinatorial Decision Making and Optimization

2nd cycle degree programme in Artificial Intelligence

University of Bologna, Academic Year 2023/24



- Let's face (combinatorial) **decision making** and **optimization** from a different perspective
 - different paradigm, same **goal**: modeling and solving hard real-world optimization problems subject to different constraints
- Less “AI-oriented” and “logic-oriented”, more “**math-oriented**”
- Less “constraints-centered”, more “**inequalities**-centered”
- **Relaxations** and **cutting-planes** rather than propagation and search

Operations research

- **Operation Research (OR)** is a well-established field based on mathematical techniques for enhancing complex **decision-making**
- Originated in first half of 20th century for military purposes, nowadays OR finds **application** in several fields, e.g.:
 - Finance
 - Manufacturing and Logistics
 - Simulations and stochastic models
 - Transportation
 - ...
- OR strongly influenced by **linear programming** techniques and its variants (ILP, MIP, NLP...)
 - As for CP, “programming” does not mean “coding” in this context...

Linear programming

- Linear programming (LP) is based on systems of linear (in-)equalities
- We typically resort to LP when we need an optimal allocation for a limited number of resources
- LP is among the most relevant scientific advances of last century: several applications in disparate fields — not only scientific fields
 - Agriculture, sports, marketing, environment etc.
 - Delta claimed 100.000.000\$ saving per year using LP
 - H. Markowitz won Nobel prize for using LP to optimize portfolio profit
- Let's start with a toy example from MiniZinc tutorial
 - <https://www.minizinc.org/doc-2.5.5/en/modelling.html#an-arithmetic-optimisation-example>


Baking cakes

```
1 % We know how to make two sorts of cakes. A banana cake which takes 250g of
2 % self-raising flour, 2 mashed bananas, 75g sugar and 100g of butter, and a
3 % chocolate cake which takes 200g of self-raising flour, 75g of cocoa, 150g
4 % sugar and 150g of butter. We can sell a chocolate cake for $4.50 and a
5 % banana cake for $4.00. And we have 4kg self-raising flour, 6 bananas,
6 % 2kg of sugar, 500g of butter and 500g of cocoa. How many of each sort of
7 % cake should we bake for the fete to maximise the profit
8
9 var 0..100: b; % no. of banana cakes
10 var 0..100: c; % no. of chocolate cakes
11
12 % flour
13 constraint 250*b + 200*c <= 4000;
14 % bananas
15 constraint 2*b <= 6;
16 % sugar
17 constraint 75*b + 150*c <= 2000;
18 % butter
19 constraint 100*b + 150*c <= 500;
20 % cocoa
21 constraint 75*c <= 500;
22
23 % maximize our profit
24 solve maximize 400*b + 450*c;
25
26 output ["no. of banana cakes = \"(b)\n",
27         "no. of chocolate cakes = \"(c)\n"];
```

Geometric interpretation

- This is a linear problem: all the constraints are **linear inequalities** and we optimize a **linear function**
- Only 2 variables involved: we can **geometrically represent** the problem
 - In 2 dim: equalities \equiv **straight-lines**, inequalities \equiv **half-plane**
 - In n dim: equalities \equiv **hyperplanes**, inequalities \equiv **half-spaces**
- **Feasible solution** \equiv assignment satisfying all the constraints \equiv points within the **intersection** of all half-spaces defined by inequalities
- Set of all solutions \equiv set of all feasible points \equiv **feasible region**
 - It is a **convex polyhedron**: it may be empty, bounded or unbounded
- **GOAL**: find a point within the feasible region where the objective function has **maximal** value

Geometric interpretation



Where is the feasible region? What point is optimal?

Drawn with <https://www.desmos.com/calculator>

Geometric interpretation



- By “*tuning the isolines*” of the objective function we find **optimal solution** $(3, 4/3)$, having **optimal value** $z = 400 \cdot 3 + 450 \cdot 4/3 = 1800\$$
- This is **inconsistent** with the model specification, where $b, c \in \mathbb{Z}$
- No worries, for now let’s assume that we can sell slices of cake
 - If not, what would be the optimal solution?

Brewery problem

- Let's see another toy example: the **brewery problem**
 - From <https://www.cs.princeton.edu/courses/archive/spr03/cs226/lectures/lp-4up.pdf>
- A small brewery needs to produce ale and (lager) beer with **limited resources**:

Beverage	Corn	Hops	Malt	Profit
Ale	5	4	35	13
Beer	15	4	20	23
Q.ty available	480	160	1190	

- How can they **maximize profits**?
 - Devote all resources to ale: 34 barrels of ale \rightarrow 442\$
 - Devote all resources to beer: 32 barrels of beer \rightarrow 736\$
 - 7.5 barrels of ale, 29.5 barrels of beer \rightarrow 776\$
 - 12 barrels of ale, 28 barrels of beer \rightarrow 800\$
 - ...

Brewery problem

- A small brewery needs to produce ale and (lager) beer with **limited resources**:


Beverage	Corn	Hops	Malt	Profit
Ale	5	4	35	13
Beer	15	4	20	23
Q.ty available	480	160	1190	

- Let's formulate this as a **LP problem**:

$$\begin{array}{llll} \text{maximize} & 13A & + & 23B \\ \text{subject to} & 5A & + & 15B \leq 480 \\ & 4A & + & 4B \leq 160 \\ & 35A & + & 20B \leq 1190 \\ & A & , & B \geq 0 \end{array}$$


Geometry

The **feasible region** is:




Geometry

We need to maximize $13A + 23B$:




Geometry

Observation: the optimal solution necessarily occurs at **extreme point**



Geometry

- **Extreme point property**: if an optimal solution for a LP problem exists, then there is one at the **extreme point** of its feasible region
- **Good news**: the number of extreme points is **finite**
- **Bad news**: the number of extreme points can be **exponential**
 - E.g., the **n -dimensional hypercube** has exactly 2^n vertices



Canonical form

- A LP problem in **canonical form** has the form:

$$\begin{array}{ll}\max & \sum_{j=1}^n c_j x_j \\ \text{s.t.} & \sum_{j=1}^n a_{i,j} x_j \leq b_i \quad 1 \leq i \leq m \\ & x_j \geq 0 \quad 1 \leq j \leq n\end{array}$$

- m = no. of linear **constraints**, n = no. of non-negative **variables**
- $a_{i,j}, b_i, c_j \in \mathbb{R}$ are known **parameters**
- $\sum_{j=1}^n c_j x_j$ is the **objective function** to maximize
 - subject to m linear inequalities $\sum_{j=1}^n a_{i,j} x_j \leq b_i$
- **Matrix form**: $\max c \cdot x$ s.t. $Ax \leq b$ and $x \geq 0$
 - $c = \langle c_1, \dots, c_n \rangle$, $x = \langle x_1, \dots, x_n \rangle^t$, $b = \langle b_1, \dots, b_m \rangle^t$,

$$A = \begin{pmatrix} a_{1,1} & \dots & a_{1,n} \\ \dots & \dots & \dots \\ a_{m,1} & \dots & a_{m,n} \end{pmatrix}$$

Standard form

- A LP problem in **standard form** has the form:

$$\begin{array}{ll}\max & \sum_{j=1}^n c_j x_j \\ \text{s.t.} & \sum_{j=1}^n a_{i,j} x_j = b_i \quad 1 \leq i \leq m \\ & x_j \geq 0 \quad 1 \leq j \leq n\end{array}$$

- **Matrix form**: $\max c \cdot x$ s.t. $Ax = b$ and $x \geq 0$
- We can easily convert from canonical to **equivalent** standard form with **m slack variables** $y_1, \dots, y_m \geq 0$
 - From n -dimensional to **$(n + m)$ -dimensional** problem
- $\sum_{j=1}^n a_{i,j} x_j \leq b_i \implies \sum_{j=1}^n a_{i,j} x_j + y_i = b_i, y_i \geq 0$ for $i = 1, \dots, m$
 - Objective function does not change

Example

- The above brewery problem is already in **canonical form**:

$$\begin{array}{llll} \max & 13A & + & 23B \\ \text{s.t.} & 5A & + & 15B \leq 480 \\ & 4A & + & 4B \leq 160 \\ & 35A & + & 20B \leq 1190 \\ & A & , & B \geq 0 \end{array}$$


- The corresponding **standard form** is:

$$\begin{array}{llllllllll} \max & 13A & + & 23B & & & & & & \\ \text{s.t.} & 5A & + & 15B & + & S_C & & & & = 480 \\ & 4A & + & 4B & & & + & S_H & & = 160 \\ & 35A & + & 20B & & & & & + & S_M = 1190 \\ & A & , & B & , & S_C & , & S_H & , & S_M \geq 0 \end{array}$$

- How to find an optimal solution with $n > 2$ variables?

Symplex algorithm

- **Symplex algorithm** developed by G. Dantzig in 1947
- General idea: start at some **extreme point** and iteratively move to a **neighboring** one that **doesn't decrease** the objective value
 - If no such extreme point exists, we found an **optimal** solution




- It exploits **linear algebra** properties

Basis

- Given a LP problem P in standard form, a **basis** of P is a subset $\mathcal{B} = \{x_{i_1}, \dots, x_{i_m}\}$ of $m \leq n$ variables s.t. **columns** A^{i_1}, \dots, A^{i_m} form a $m \times m$ **invertible matrix** $A_{\mathcal{B}}$
 - E.g., if $A = \begin{pmatrix} 1 & 2 & 2 \\ 1 & 2 & 1 \end{pmatrix}$ then $\mathcal{B} = \{x_1, x_2\}$ is not a basis because $A_{\mathcal{B}} = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix}$ is not invertible, while $\{x_1, x_3\}$ and $\{x_2, x_3\}$ are basis
- We can rewrite P by separating **basic** from **non-basic** variables:
 $\max(c_{\mathcal{B}}x_{\mathcal{B}} + c_{\mathcal{N}}x_{\mathcal{N}})$ s.t. $A_{\mathcal{B}}x_{\mathcal{B}} + A_{\mathcal{N}}x_{\mathcal{N}} = b$ and $x_{\mathcal{B}}, x_{\mathcal{N}} \geq 0$
 - $\mathcal{N} = \{x_1, \dots, x_n\} - \mathcal{B}$ are the non-basic variables of P
- By setting $x_{\mathcal{N}} = 0$, P becomes $\max(c_{\mathcal{B}}x_{\mathcal{B}})$ s.t. $A_{\mathcal{B}}x_{\mathcal{B}} = b$ hence $x_{\mathcal{B}} = A_{\mathcal{B}}^{-1}b \in \mathbb{R}^m$ with objective value $c_{\mathcal{B}}A_{\mathcal{B}}^{-1}b$. This solution is called a **basic solution** for \mathcal{B}

Basis

- A basic solution for \mathcal{B} is **feasible** iff $(\forall_{i=1}^m) (x_{\mathcal{B}})_i \geq 0$
 - **Basic feasible solution (BFS)** \equiv extreme point of feasible region
- A BFS for \mathcal{B} is **non-degenerate** iff $(\forall_{i=1}^m) (x_{\mathcal{B}})_i > 0$
 - If BFS non-degenerate, then it's represented by a **unique** basis
- Simplex method **iteratively** considers BFS $\tilde{x}_1, \tilde{x}_2, \dots$ s.t. $c\tilde{x}_k \geq c\tilde{x}_{k-1}$



Brewery example

- The brewery example in **standard form** is:

$$\begin{array}{llllllllll} \max & 13A & + & 23B & & & & & & \\ \text{s.t.} & 5A & + & 15B & + & S_C & & & & = 480 \\ & 4A & + & 4B & & & + & S_H & & = 160 \\ & 35A & + & 20B & & & & & + & S_M = 1190 \\ & A & , & B & , & S_C & , & S_H & , & S_M \geq 0 \end{array}$$

- Let's first pick an arbitrary feasible basis, e.g., $\mathcal{B} = \{S_C, S_H, S_M\}$
 - $A_{\mathcal{B}}$ is the 3×3 identity matrix, $\mathcal{N} = \{A, B\}$ so $A = B = 0$
- The BFS for \mathcal{B} is $S_C = 480, S_H = 160, S_M = 1190$ with obj. value 0
 - Feasible, but we may improve it with a new basis \mathcal{B}' adjacent to \mathcal{B}
 - $\mathcal{B}' = \mathcal{B} \cup \{x^{in}\} - \{x^{out}\}$ with $x^{in} \notin \mathcal{B}$ and $x^{out} \in \mathcal{B}$

Pivoting

- How to select entering variable x^{in} and leaving variable x^{out} ?
- We can choose the x^{in} which “increases more” the objective value
 - x^{in} can increase from 0 ($x^{in} \in \mathcal{N}$) to a value ≥ 0 ($x^{in} \in \mathcal{B}'$)
 - In the brewery example, a unit increase in A increases the obj. value of 13; a unit increase in B increases the obj. value of 23: $x^{in} = B$
- Then, choose x^{out} by ensuring that $\mathcal{B}' = \mathcal{B} \cup \{x^{in}\} - \{x^{out}\}$ is a feasible basis
 - Minimum ratio rule: x^{out} is in the row i minimizing ratio $\beta_i^{in} / \alpha_i^{in}$ where $\alpha_i^{in}, \beta_i^{in}$ are the x^{in} coefficient and known term in the i -th row
 - Otherwise, \mathcal{B}' not feasible
 - In the brewery ex. $\min\{480/15, 160/4, 1190/20\} = \min\{32, 40, 59.5\} = 32$ so $x^{out} = S_C$: the new basis will be $\mathcal{B}' = \{B, S_H, S_M\}$
 - Then, x^{in} is derived and its value replaced in all other equations

Pivoting

- Selecting x^{in} , x^{out} resp. means choosing a **pivot column** and a **pivot row** from the **tableau** representation of $\max(Z)$ s.t.:

$$\begin{array}{rclclclclcl} 13A & + & 23B & & & - & Z & = & 0 \\ \hline 5A & + & 15B & + & S_C & & & = & 480 \\ 4A & + & 4B & & & + & S_H & = & 160 \\ 35A & + & 20B & & & & + & S_M & = & 1190 \\ \hline A & , & B & , & S_C & , & S_H & , & S_M & \geq 0 \end{array}$$

- The row of x^{out} (in this case, S_C) is the “**most restrictive**”, i.e., the first one to be violated if B increases too much:
 - In row 1, B can be increased up to $480/15 = 32$
 - In row 2, B can be increased up to $160/4 = 40$
 - In row 3, B can be increased up to $1190/20 = 59.5$

Pivoting

- Selecting x^{in} , x^{out} resp. means choosing a **pivot column** and a **pivot row** from the **tableau** representation of $\max(Z)$ s.t.:

$$\begin{array}{rclclclcl}
 13A & + & 23B & & & - & Z & = & 0 \\
 \hline
 5A & + & 15B & + & S_C & & & = & 480 \\
 4A & + & 4B & & & + & S_H & = & 160 \\
 35A & + & 20B & & & & + & S_M & = & 1190 \\
 \hline
 A & , & B & , & S_C & , & S_H & , & S_M & \geq 0
 \end{array}$$

- From x^{out} row we get $B = \frac{480-5A-S_C}{15} = 32 - \frac{1}{3}A - \frac{1}{15}S_C$ and we **substitute** it (remember Gauss-Jordan?) in all other equations:

$$\begin{array}{rclclclcl}
 \frac{16}{3}A & - & \frac{23}{15}S_C & & & - & Z & = & -736 \\
 \hline
 \frac{1}{3}A & + & B & + & \frac{1}{15}S_C & & & = & 32 \\
 \frac{8}{3}A & & & + & \frac{4}{15}S_C & + & S_H & = & 32 \\
 \frac{85}{3}A & & & - & \frac{4}{3}S_C & & + & S_M & = & 550 \\
 \hline
 A & , & B & , & S_C & , & S_H & , & S_M & \geq 0
 \end{array}$$

Pivoting

- Current basis $\{B, S_H, S_M\}$. What row/column should we select now?

$$\begin{array}{rclclclclcl}
 \frac{16}{3}A & - & \frac{23}{15}S_C & & & - & Z & = & -736 \\
 \hline
 \frac{1}{3}A & + & B & + & \frac{1}{15}S_C & & & = & 32 \\
 \frac{8}{3}A & & & - & \frac{4}{15}S_C & + & S_H & = & 32 \\
 \frac{85}{3}A & & & - & \frac{4}{3}S_C & & + & S_M & = & 550 \\
 \hline
 A & , & B & , & S_C & , & S_H & , & S_M & \geq 0
 \end{array}$$

Pivoting

- Current basis $\{B, S_H, S_M\}$. What row/column should we select now?

$$\begin{array}{rclclclclcl}
 \frac{16}{3}A & - & \frac{23}{15}S_C & & & - & Z & = & -736 \\
 \hline
 \frac{1}{3}A & + & B & + & \frac{1}{15}S_C & & & = & 32 \\
 \frac{8}{3}A & & & - & \frac{4}{15}S_C & + & S_H & = & 32 \\
 \frac{85}{3}A & & & - & \frac{4}{3}S_C & & + & S_M & = & 550 \\
 \hline
 A & , & B & , & S_C & , & S_H & , & S_M & \geq 0
 \end{array}$$

- A contributes more than S_C in increasing Z : 1st column chosen ($x^{in} = A$)
 - The coefficients in the obj. function row are called **reduced costs** or sometimes relative profits

Pivoting

- Current basis $\{B, S_H, S_M\}$. What row/column should we select now?

$$\begin{array}{rcccccccl}
 \frac{16}{3}A & - & \frac{23}{15}S_C & & & - & Z & = & -736 \\
 \hline
 \frac{1}{3}A & + & B & + & \frac{1}{15}S_C & & & = & 32 \\
 \frac{8}{3}A & & & - & \frac{4}{15}S_C & + & S_H & = & 32 \\
 \frac{85}{3}A & & & - & \frac{4}{3}S_C & & + & S_M & = & 550 \\
 \hline
 A & , & B & , & S_C & , & S_H & , & S_M & \geq 0
 \end{array}$$

- A contributes more than S_C in increasing Z : $x^{in} = A$
- The ratios for A are $\{32 \cdot 3, 32 \cdot \frac{3}{8}, 550 \cdot \frac{3}{85}\} = \{96, 12, 19.41 \dots\}$: 2nd row chosen ($x^{out} = S_H$)

Pivoting

- Current basis $\{B, S_H, S_M\}$. What row/column should we select now?

$$\begin{array}{rcccccccl}
 \frac{16}{3}A & - & \frac{23}{15}S_C & & & - & Z & = & -736 \\
 \hline
 \frac{1}{3}A & + & B & + & \frac{1}{15}S_C & & & = & 32 \\
 \frac{8}{3}A & & & - & \frac{4}{15}S_C & + & S_H & = & 32 \\
 \frac{85}{3}A & & & - & \frac{4}{3}S_C & & + & S_M & = & 550 \\
 \hline
 A & , & B & , & S_C & , & S_H & , & S_M & \geq 0
 \end{array}$$

- A contributes more than S_C in increasing Z : $x^{in} = A$
- The ratios for A are $\{32 \cdot 3, 32 \cdot \frac{3}{8}, 550 \cdot \frac{3}{85}\}$: $x^{out} = S_H$
- New basis: $\{A, B, S_M\}$. We derive $A = \frac{3}{8} \cdot (32 + \frac{4}{15}S_C - S_H) = 12 + \frac{1}{10}S_C - \frac{3}{8}S_H$ and substitute it in the other equations

Pivoting

- Current basis $\{A, B, S_M\}$. What row/column should we select now?

$$\begin{array}{rcccccccl}
 & - & S_C & - & 2S_H & & - & Z & = -800 \\
 \hline
 & & B & + & \frac{1}{10}S_C & + & \frac{1}{8}S_H & & = 28 \\
 A & & & - & \frac{1}{10}S_C & + & \frac{3}{8}S_H & & = 12 \\
 & & & - & \frac{25}{6}S_C & - & \frac{85}{8}S_H & + & S_M = 210 \\
 \hline
 A, & B, & S_C, & S_H, & S_M & \geq 0
 \end{array}$$

Optimality

- Current basis $\{A, B, S_M\}$. What row/column should we select now?

$$\begin{array}{rclclclcl}
 & - & S_C & - & 2S_H & & - & Z & = & -800 \\
 \hline
 & & B & + & \frac{1}{10}S_C & + & \frac{1}{800}S_H & & = & 28 \\
 A & & - & \frac{1}{10}S_C & + & \frac{1}{800}S_H & & & = & 12 \\
 & & - & \frac{25}{6}S_C & - & \frac{85}{8}S_H & + & S_M & = & 210 \\
 \hline
 A & , & B & , & S_C & , & S_H & , & S_M & \geq 0
 \end{array}$$

- All the **reduced costs** are ≤ 0 : increasing the value of corresponding variables won't increase the obj. value
- We cannot improve the current feasible solution \rightarrow we reached an **optimal** solution:
 - $S_C = S_H = 0$
 - $A = 12, B = 28, S_M = 210$
 - $-S_C - 2S_H - Z = -800 \implies Z = 800 - S_C - 2S_H = 800$

Optimality

- The simplex method performs an “**optimality check**”: if **all** the reduced costs are ≤ 0 , we reached an optimal solution
- This condition is **sufficient**: for any optimal solution there is at least a basis s.t. all the reduced costs are ≤ 0
- ...But it's **not necessary**: we may reach an optimal solution even if some reduced cost is > 0
- E.g., $\max(x_1)$ s.t. $x_3 = 1 - x_2, x_4 = -x_1, x_i \geq 0$ with basis $\mathcal{B} = \{x_3, x_4\}$ corresponds to solution $x_3 = 1, x_1 = x_2 = x_4 = 0$.
- If we switch to $\mathcal{B} = \{x_3, x_1\}$ we get $\max(-x_4)$ s.t. $x_3 = 1 - x_2, x_1 = -x_4$: optimality condition is OK but last solution not improved

Optimal region

- The **feasible region** for a LP problem P in **canonical** form is a set $\mathcal{F}_P = \{x \in \mathbb{R}^n \mid Ax \leq b, x \geq 0\}$ denoting a **convex polyhedron**
- The **optimal region** for a LP problem P in standard form is a set of solutions $\mathcal{O}_P = \{x^* \in \mathcal{F}_P \mid cx^* \geq cx, \forall x \in \mathcal{F}_P\}$
- Clearly $\mathcal{O}_P \subseteq \mathcal{F}_P$ and $\mathcal{F}_P = \emptyset \Rightarrow \mathcal{O}_P = \emptyset$
- If \mathcal{O}_P is **finite**, then $|\mathcal{O}_P| = 1$ (hence $|\mathcal{O}_P| > 1 \Rightarrow \mathcal{O}_P$ **infinite**)
 - if $x_1, x_2 \in \mathcal{O}_P$ and $x_1 \neq x_2$ then all points in segment $\overline{x_1 x_2}$ are in \mathcal{O}_P because $x_1, x_2 \in \mathcal{F}_P$ which is convex
- Is there any problem P such that $\mathcal{F}_P \neq \emptyset \wedge \mathcal{O}_P = \emptyset$?

Unboundedness

- Simply consider $\max(x)$ s.t. $x \geq 0$: we have $\mathcal{F}_P = [0, +\infty)$ but there is no $x^* \in \mathcal{R}$ s.t. $x^* \geq x$ for all $x \in \mathcal{F}_P$: $\mathcal{O}_P = \emptyset$
- In these cases P is said **unbounded**: no optimal solution exists
 - \mathcal{F}_P is an **unbounded polyhedron**
- Simplex method also performs an “**unboundedness check**”
- With the tableau method seen above, we must ensure that **no column** j has reduced cost $\gamma_j > 0$ and coefficients $\alpha_{i,j} \leq 0$ for $i = 1, \dots, m$
 - In the literature you can find different but equivalent formulations


Unboundedness

- For example, consider $\max(x - y + 2)$ s.t. $x + y \geq 0$, $2x + y \geq -1$, $x, y \geq 0$. The tableau of the problem can be written as:

$$\begin{array}{rclclcl}
 & x & - & y & & - & Z & = & -2 \\
 \hline
 - & x & - & y & + & S_1 & & = & 0 \\
 - & 2x & - & y & & & + & S_2 & = & 1 \\
 \hline
 & x & , & y & , & S_1 & , & S_2 & \geq & 0
 \end{array}$$

- With $\mathcal{B} = \{S_1, S_2\}$ we have a BFS with $x = y = 0$ and obj. value 2
 - Degenerate solution ($S_1 = 0$)
- Reduced cost of x is $1 > 0$ and its row coefficients are $-1, -2 \leq 0$: the problem is **unbounded**: x can be arbitrarily increased **within** the feasible region

Unboundedness



- E.g., from each feasible solution (\tilde{x}, \tilde{y}) we can always derive a **better** solution $(\tilde{x} + k, \tilde{y})$ for each $k > 0$

Simplex steps

Given LP P in **standard form**, we can (roughly) summarize the simplex method as:

0. Let $k \leftarrow 0$, let \mathcal{B}_0 a **feasible base** for P and go to 1.
1. If BFS of \mathcal{B}_k is **optimal** then **STOP**, else go to 2.
2. If P **unbounded** then **STOP**, else go to 3.
3. Select an **entering** variable $x^{in} \notin \mathcal{B}_k$ and go to 4.
4. Select a **leaving** variable $x^{out} \in \mathcal{B}_k$ and go to 5.
5. Let $\mathcal{B}_{k+1} = \mathcal{B}_k \cup \{x^{in}\} - \{x^{out}\}$ and **reformulate** P accordingly.
Let $k \leftarrow k + 1$ and go back to 1.

- The former baking example in standard form is:

$$\begin{array}{llllll} \max & 400B & + & 450C & & \\ \text{s.t.} & 250B & + & 200C & + & S_1 = 4000 \\ & 2B & & & + & S_2 = 6 \\ & 75B & + & 150C & + & S_3 = 2000 \\ & 100B & + & 150C & + & S_4 = 500 \\ & & & 75C & + & S_5 = 500 \\ & B & , & C & , & S_i \geq 0 \quad i = 1, \dots, 5 \end{array}$$

- Exercise:** Find the optimal solution through the simplex method.

Simplex properties

- If **all** the possible BFS are **non-degenerate**, the simplex method always **terminates** in a finite number of steps
 - The no. of vertices is **finite** and at each step we move from one vertex to another always **strictly improving** the obj. value
- Otherwise, possible **stalling**: we repeatedly change base without improving the obj. value (common in large scale applications)
 - **Termination** can be guaranteed with **anti-cycling rules** preventing possible loops $\mathcal{B}_k \rightarrow \mathcal{B}_{k+1} \rightarrow \dots \rightarrow \mathcal{B}_k$
- **Worst-case** time-complexity of simplex method is $O(2^n)$ but in practice is typically **polynomial**

Other LP algorithms

- Khachiyan's Ellipsoid Algorithm (1979)
 - Proves that LP problem belongs to \mathcal{P} , but not practical
- Karmakar's algorithm (1984): 1st practical polynomial-time algorithm employing a interior point method
 - traverses feasible region “internally” instead of exploring extreme points
- In general interior point methods scale better than simplex for (very) large problems
 - Not so suitable for MIP problems including discrete variables

Two-phase method

- So far we assumed that we can always find a **feasible base**
 - How to choose an initial **feasible base** B_0 ?
 - What if the problem is **unsatisfiable**?
- The **two-phase method** finds (if any) an initial base for a standard problem P by first solving an “**artificial**” **problem** P' derived from P by adding **fresh variables** s_1, \dots, s_m
- **2nd phase** problem P : $\max(cx)$ s.t. $Ax = b, x \geq 0$
- **1st phase** problem P' : $\max(-\sum_{i=1}^m s_i)$ s.t.
 - $\sum_{j=1}^n a_{i,j}x_j + s_i = b_i$ for $i \in \{k \in \{1, \dots, m\} \mid b_k \geq 0\}$
 - $\sum_{j=1}^n a_{i,j}x_j - s_i = b_i$ for $i \in \{k \in \{1, \dots, m\} \mid b_k < 0\}$
 - $s_i, x_j \geq 0$

Two-phase method

- 1st phase problem P' : $\max(-\sum_{i=1}^m s_i)$ s.t.
 - $\sum_{j=1}^n a_{i,j}x_j + s_i = b_i$ for $i \in \{k \in \{1, \dots, m\} \mid b_k \geq 0\}$
 - $\sum_{j=1}^n a_{i,j}x_j - s_i = b_i$ for $i \in \{k \in \{1, \dots, m\} \mid b_k < 0\}$
 - $s_i, x_j \geq 0$

- E.g., if P is:

$$\max(x_1 + 2x_2) \quad \text{s.t.}$$

$$-x_1 - x_2 + x_3 = -1$$

$$x_1 + x_2 + x_4 = 2$$

$$x_1, \dots, x_4 \geq 0$$

- Then P' is:

$$\max(-s_1 - s_2) \quad \text{s.t.}$$

$$-x_1 - x_2 + x_3 - s_1 = -1$$

$$x_1 + x_2 + x_4 + s_2 = 2$$

$$x_1, \dots, x_4, s_1, s_2 \geq 0$$

Two-phase method

- Note that objective $-\sum_{i=1}^m s_i$ is always ≤ 0 and $\mathcal{B}' = \{s_1, \dots, s_m\}$ is always a **feasible basis** corresponding to BFS $x_j = 0, s_i = |b_i|$
 - Hence, $\mathcal{F}_{P'} \neq \emptyset$ and $\mathcal{O}_{P'} \neq \emptyset$ (P' upper-bounded by 0)

- So we **reformulate** P' w.r.t. \mathcal{B}' . In the example above we get:

$$\max(-3 - 2x_1 + 2x_2 - x_3 + x_4) \text{ s.t.}$$


$$s_1 = 1 - x_1 - x_2 + x_3$$

$$s_2 = 2 - x_1 - x_2 - x_4$$

$$x_1, \dots, x_4, s_1, s_2 \geq 0$$

- Then we solve P' with the simplex method. A nice property is that $\mathcal{F}_P \neq \emptyset \iff \sum_{i=1}^m s_i = 0$:
 - If the optimal value of P' is < 0 , then P is **unsatisfiable**,
 - Otherwise, from the basis corresponding to the optimal solution of P' we get an **initial basis** for P by **removing** s_i variables

Two-phase method



Entrepreneur problem

- Let's now tackle the brewery problem from a different angle

Beverage	Corn	Hops	Malt	Profit
Ale	5	4	35	13
Beer	15	4	20	23
Q.ty available	480	160	1190	

- An **entrepreneur** wants to buy individual resources (corn, hops, malt) **from** brewer at **minimum cost**
- The brewer won't sell resources if $5C + 4H + 35M < 13$ (Ale profit) and $15C + 4H + 20M < 23$ (Beer profit)
- What would be the **minimum unit cost** for corn (C), hops (H), and malt (M) given the resource availability and brewer's constraints?

Entrepreneur problem

- Let's now tackle the brewery problem from a different angle


Beverage	Corn	Hops	Malt	Profit
Ale	5	4	35	13
Beer	15	4	20	23
Q.ty available	480	160	1190	

- The entrepreneur LP problem can be formulate as:

$$\begin{array}{llllllll} \text{minimize} & 480C & + & 160H & + & 1190M & & \\ \text{subject to} & 5C & + & 4H & + & 35M & \geq & 13 \\ & 15C & + & 4H & + & 20M & \geq & 23 \\ & C & , & H & , & M & \geq & 0 \end{array}$$

- Optimal solution: $C = 1, H = 2, M = 0$ with total cost 800\$
 - Exercise: transform in canonical and standard form

Entrepreneur problem



Duality

- The entrepreneur problem is the **dual** of the brewery problem
- Price **evaluation** rather than resource allocation
- Two different perspectives, but **same optimal value**
 - Brewer knows that can **earn at most 800\$**
 - Entrepreneur knows that has to **spend at least 800\$**
- The **duality** concept is important (not only) in LP and can be extended to general (N)LP problems

Duality

- Let $P : \max(cx)$ s.t. $Ax = b, x \geq 0$ with $b \in \mathbb{R}^m, x \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n}$ the **primal** problem
- Its **dual** problem $\mathcal{D}(P) : \min(by)$ s.t. $A^t y \geq c$ with $y \in \mathbb{R}^m$ has:
 - a variable y_i for each constraint $\sum_{j=1}^n a_{i,j}x_j = b_i$ of P , $i = 1, \dots, m$
 - a constraint $\sum_{i=1}^m a_{j,i}y_i \leq c_j$ for each variable x_j of P , $j = 1, \dots, n$
- Exercise:** find the dual of following primal problem:

$$\begin{array}{rcllclclcl} \max & x_1 & + & x_2 & & & & & \\ \text{s.t.} & 3x_1 & + & 2x_2 & + & x_3 & & & = 5 \\ & 4x_1 & + & 5x_2 & & & + & x_4 & = 4 \\ & & & + & x_2 & & & + & x_5 & = 2 \\ & x_1 & , & x_2 & , & x_3 & , & x_4 & , & x_5 & \geq 0 \end{array}$$

Duality

- Let $P : \max(cx)$ s.t. $Ax = b, x \geq 0$ with $b \in \mathbb{R}^m, x \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n}$ the **primal** problem
- Its **dual** problem $\mathcal{D}(P) : \min(by)$ s.t. $A^t y \geq c$ with $y \in \mathbb{R}^m$ has:
 - a variable y_i for each constraint $\sum_{j=1}^n a_{i,j}x_j = b_i$ of P , $i = 1, \dots, m$
 - a constraint $\sum_{i=1}^m a_{j,i}y_i \leq c_j$ for each variable x_j of P , $j = 1, \dots, n$
- Exercise:** find the dual of following primal problem:

$$\begin{array}{llllllll} \max & x_1 & + & x_2 & & & & \\ \text{s.t.} & 3x_1 & + & 2x_2 & + & x_3 & & = 5 & (y_1) \\ & 4x_1 & + & 5x_2 & & & + & x_4 & = 4 & (y_2) \\ & & + & x_2 & & & & + & x_5 & = 2 & (y_3) \\ & x_1 & , & x_2 & , & x_3 & , & x_4 & , & x_5 & \geq 0 \\ & (con_1) & , & (con_2) & , & (con_3) & , & (con_4) & , & (con_5) \end{array}$$

Duality

- Let $P : \max(cx)$ s.t. $Ax = b, x \geq 0$ with $b \in \mathbb{R}^m, x \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n}$ the **primal** problem
- Its **dual** problem $\mathcal{D}(P) : \min(by)$ s.t. $A^t y \geq c$ with $y \in \mathbb{R}^m$ has:
 - a variable y_i for each constraint $\sum_{j=1}^n a_{i,j}x_j = b_i$ of P , $i = 1, \dots, m$
 - a constraint $\sum_{i=1}^m a_{j,i}y_i \leq c_j$ for each variable x_j of P , $j = 1, \dots, n$
- Exercise:** find the dual of following primal problem:

$$\begin{array}{llll} \max & x_1 + x_2 & \text{s.t.} & \implies \min & 5y_1 + 4y_2 + 2y_3 & \text{s.t.} \\ 3x_1 + 2x_2 + x_3 & = 5 & (y_1) & 3y_1 + 4y_2 & \geq 1 \\ 4x_1 + 5x_2 + x_4 & = 4 & (y_2) & 2y_1 + 5y_2 + y_3 & \geq 1 \\ x_2 + x_5 & = 2 & (y_3) & y_1 & \geq 0 \\ x_1, x_2, x_3, x_4, x_5 & \geq 0 & & y_2 & \geq 0 \\ & & & y_3 & \geq 0 \end{array}$$

Duality

- Let $P : \max(cx)$ s.t. $Ax = b, x \geq 0$ with $b \in \mathbb{R}^m, x \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n}$ the **primal** problem
- Its **dual** problem $\mathcal{D}(P) : \min(by)$ s.t. $A^t y \geq c$ with $y \in \mathbb{R}^m$ has:
 - a variable y_i for each constraint $\sum_{j=1}^n a_{i,j}x_j = b_i$ of P , $i = 1, \dots, m$
 - a constraint $\sum_{i=1}^m a_{j,i}y_i \leq c_j$ for each variable x_j of P , $j = 1, \dots, n$
- What about the **dual** of the **dual**?

$$\begin{array}{llll} \max & x_1 + x_2 & \text{s.t.} & \implies \\ & 3x_1 + 2x_2 + x_3 = 5 & (y_1) & \\ & 4x_1 + 5x_2 + x_4 = 4 & (y_2) & \\ & x_2 + x_5 = 2 & (y_3) & \\ & x_1, x_2, x_3 \geq 0 & & \end{array} \quad \begin{array}{llll} \min & 5y_1 + 4y_2 + 2y_3 & \text{s.t.} & \implies \\ & 3y_1 + 4y_2 \geq 1 & (x_1) & \\ & 2y_1 + 5y_2 + y_3 \geq 1 & (x_2) & \\ & y_1 \geq 0 & (x_3) & \\ & y_2 \geq 0 & (x_4) & \\ & y_3 \geq 0 & (x_5) & \end{array}$$

Duality properties

- $\mathcal{D}(\mathcal{D}(P)) = P$: the dual of the dual is the **primal**
- **Weak duality**: the cost of **any** feasible primal solution is **at most** the cost of **any** feasible dual solution: $(\forall x \in \mathcal{F}_P, \forall y \in \mathcal{F}_{\mathcal{D}(P)}) \quad cx \leq by$
 - by is an **upper bound** for obj. value of P
 - cx is a **lower bound** for obj. value of $\mathcal{D}(P)$
 - by “decreases” until a **minimum** value eventually reached ↘
 - cx “increases” until a **maximum** value eventually reached ↗
 - If P unbounded, $\mathcal{D}(P)$ unfeasible: $(\mathcal{F}_P \neq \emptyset \wedge \mathcal{O}_P = \emptyset) \Rightarrow \mathcal{F}_{\mathcal{D}(P)} = \emptyset$
 - if $\mathcal{D}(P)$ unbounded, P unfeasible: $(\mathcal{F}_{\mathcal{D}(P)} \neq \emptyset \wedge \mathcal{O}_{\mathcal{D}(P)} = \emptyset) \Rightarrow \mathcal{F}_P = \emptyset$
- **Strong duality**: if primal and dual are feasible they have **same optimal** cost: $\mathcal{F}_P, \mathcal{F}_{\mathcal{D}(P)} \neq \emptyset \Rightarrow (\forall x^* \in \mathcal{O}_P, \forall y^* \in \mathcal{O}_{\mathcal{D}(P)}) \quad cx^* = by^*$
 - Remember brewery example?

Possible cases

- Note that primal unbounded \implies dual unfeasible but in general
primal unfeasible $\not\Rightarrow$ dual unbounded
 - E.g., $P : \max(2x_1 - x_2)$ s.t. $x_1 - x_2 \leq 1, -x_1 + x_2 \leq -2, x_1, x_2 \geq 0$ is unfeasible, and so is $\mathcal{D}(P)$
 - Exercise: build the dual, prove unsatisfiability with 2-phase method
- In summary (✓ = possible, ✗ = impossible):

	$\mathcal{D}(P)$ bounded	$\mathcal{D}(P)$ unbounded	$\mathcal{D}(P)$ unfeasible
P bounded	✓	✗	✗
P unbounded	✗	✗	✓
P unfeasible	✗	✓	✓

Dual simplex

- We can avoid to compute $\mathcal{D}(P)$ to apply the (primal) simplex on $\mathcal{D}(P)$ by running on P the dual simplex
 - C.E. Lemke, 1954
- Primal simplex: from feasible to optimal basis, preserving feasibility
- Dual simplex: from “optimal basis” (reduced costs ≤ 0 , not necessarily feasible) to feasible basis, while preserving optimality
 - x^{out} = variable with minimum value
 - x^{in} = variable with maximum ratio
- Primal-dual: hybrid approach

Why duality?

- **Theoretical** purposes
 - E.g., finding a **feasible** solution is **as hard as** finding the **optimal** one: if we can find a solution for $\max(cx)$ s.t. $Ax = b, x \geq 0$ in $T(n, m)$ time, we can find one for $Ax = b, x \geq 0, A^t y \geq c, cx = by$ in $O(T(n, m))$
- **Prove infeasibility** of the primal problem (via dual simplex)
- **Bounding** the objective function (maybe with parallel solving)
 - Primal gives **lower** bound, dual gives **upper** bound
- **Exploiting** alternative/hybrid approaches
 - (primal-)dual simplex, (logic-based) Benders' decomposition
- **Sensitivity analysis**

Sensitivity analysis

- **Sensitivity analysis** refers to how the optimal solution of a problem is affected by **changes** in the input **parameters**
 - **Post-optimality** analysis
- If x^* is the optimal solution for standard LP problem P , will x^* be still **feasible** and/or **optimal** for a **perturbed** problem $\bar{P} \neq P$?
- \bar{P} can be obtained from P by **altering**:
 - The **known term**: $b \rightsquigarrow \bar{b} = b + \Delta b$
 - The **objective function** coefficients: $c \rightsquigarrow \bar{c} = c + \Delta c$
 - The **variables** coefficients: $A \rightsquigarrow \bar{A} + \Delta A$

Changing known term

- **Known term:** Changing $b \rightsquigarrow \bar{b} = b + \Delta b$ can affect both the **feasibility** and **optimality** of current solution


Beverage	Corn	Hops	Malt	Profit
Ale	5	4	35	13
Beer	15	4	20	23
Q.ty available	480	160	1190	

- E.g., for brewery problem if corn availability **decreases** from 480 to 479 the optimal solution $A = 12, B = 28$ is **not feasible** anymore and the objective value drops from 800 to 790 ($A = 13, B = 27$)
 - If hops decreases 1 unit, objective value drops to 787 ($A = 11, B = 28$)
 - If malt decreases up to 210 units, $A = 12, B = 28$ **still optimal**
 - In fact, $S_C = S_H = 0$ and $S_M = 210$ in the original optimal solution

Changing known term

- In general, **increasing b** doesn't affect feasibility, but can **improve** the obj. value
 - $\Delta b < 0 \rightarrow$ narrowing \mathcal{F}_P , $\Delta b > 0 \rightarrow$ extending \mathcal{F}_P
- For the brewery problem:
 - Increasing corn of ≥ 10 units improves the profit
 - Increasing hops of ≥ 4 units improves the profit
 - Increasing malt only **never improves** the profit
- Changing known term of $P \equiv$ changing objective function of $\mathcal{D}(P)$
- We can compute the impact of Δb on P **without re-solving** P : if \mathcal{B} is an **optimal basis** for P , $\bar{x}_{\mathcal{B}} = A_{\mathcal{B}}^{-1}\bar{b} = A_{\mathcal{B}}^{-1}b + A_{\mathcal{B}}^{-1}\Delta b$
 - If feasible ($\bar{x}_{\mathcal{B}} \geq 0$), the objective value $c_{\mathcal{B}}A_{\mathcal{B}}^{-1}\bar{b}$ can change as well

Geometric interpretation



$$0 \leq x \leq 100$$

$$0 \leq y \leq 100$$

$$250x + 200y \leq 4000$$


$$2x \leq 6$$

$$75x + 150y \leq 2000$$

$$100x + 150y \leq 500$$

$$75y \leq 500$$

$$400x + 450y = 0$$



Baking example: what ingredients should we (*not*) **buy** first to increase profit? If we can't buy, which ones should we **sell** without altering profit?

Changing costs

- **Cost coefficients:** Changing $c \rightsquigarrow \bar{c} = c + \Delta c$ can't affect **feasibility**, but may involve:
 - Loss of **optimality** of current solution
 - Different **objective value** for current solution (still optimal)
- E.g., in the brewery example if the profit of beer is **39** instead of 23 the optimal solution will be still **$A = 12, B = 28$**
 - But total profit would be **1248** instead of 800!
 - The same applies if beer profit drops to **14** (total profit 548)
- If beer profit is outside **$[14, 39]$** the solution changes as well
 - E.g., if beer profit = 13 then **$A = 26, B = 14$** is more convenient
- Again, we don't need to re-solve P to assess the impact of Δc

Changing constraints

- The impact of changing a coefficient $A \rightsquigarrow \bar{A} = A + \Delta A$ depends on whether $\bar{a}_{i,j}$ refers to a variable x_j in the **optimal basis**:
- If not, $x_j = 0$: current solution still feasible, its value won't change.
 - Objective function can change: we **may lose optimality**
- If yes, we can't say much: we need to **re-solve** with \bar{A}
- E.g., if the ale production would now require 4 units of corn instead of 5, then $A = 12, B = 28$ still feasible but no more optimal
 - $A = 11, B = 29$ would be better in this case (total profit 810)

Take-home messages

- Linear programming (LP) is one of the main areas of Operations Research (OR) field
- LP is about solving problems with linear constraints/objective function in a canonical or standard form
- The feasible region for a LP is a convex polyhedron
 - it can be empty, bounded, unbounded
- The simplex algorithm is a well-known method to tackle LP problems
 - Worst-case exponential, typically polynomial
 - Worst-case polynomial algorithms exist for LP problems

Take-home messages

- Simplex works by moving from one **extreme point** of the feasible region to a “not-worse one” up to an optimal point
 - **Interior point** methods traversing feasible region may scale better
- **Two-phase method** to check feasibility and get initial basis
- We can solve the **dual** of a LP problem: different perspective, **same optimal value**
- **Sensitivity analysis** to assess the effect of **perturbations** of the original problem
 - **Post-optimality** analysis, often doable without re-solving the problem

Resources

- <https://www.cs.princeton.edu/courses/archive/spr03/cs226/lectures/lp-4up.pdf>
- <https://www.coursera.org/lecture/solving-algorithms-discrete-optimization>
- ...