

Emergent Topologies and Quantum Field Mechanics: A Framework for Spacetime Engineering, Zero-Point Energy, and Traversable Wormholes

The Ontological Shift: From Smooth Manifolds to Quantum Field Topologies

For over a century, the general theory of relativity has treated spacetime as a continuous, pseudo-Riemannian manifold, a passive geometric stage warped by the presence of classical mass and energy. However, the synthesis of quantum information science, holography, and background-independent quantum gravity necessitates a radical ontological reframing. Spacetime is no longer viewed as a fundamental geometric background; rather, it is an emergent macroscopic property arising from the complex entanglement structure of microscopic quantum degrees of freedom. This paradigm shift requires transitioning from continuous mathematical topology to discrete quantum field mechanics, where locality, causality, and dimensionality are merely phase transitions of a fundamental quantum network. In this reframed ontology, the manipulation of spacetime—such as the theoretical folding of space to create traversable wormholes—is inherently a problem of quantum field mechanics. By modifying the entanglement structure of the vacuum and exploiting localized negative energy densities, the macroscopic geometry of the universe can theoretically be re-engineered. This report exhaustively details the theoretical mechanisms for emergent spacetime topologies, the holographic protocols for spacetime folding (specifically the Gao-Jafferis-Wall protocol), the stabilization of zero-point energy via squeezed vacuum states, and the rigorous constraints imposed by Quantum Energy Inequalities (QEIs). Furthermore, it incorporates relevant experimental and theoretical research originating from the University of Oregon in Eugene, culminating in a structured research protocol designed to guide future investigations into spacetime manipulation.

Topological Emergence of Spacetime

Quantum Graphity and the Condensation of Locality

To reframe mathematical topology into quantum field mechanics, one must address how a local, low-dimensional spacetime emerges from a pre-geometric, seemingly structureless state. "Quantum Graphity," a background-independent model of emergent locality, models spacetime as a dynamical graph of N vertices, where the states of the system correspond to the connectivity of these vertices rather than coordinates in a pre-existing space. In this framework, the physical degrees of freedom live on the edges of a complete graph, and the dynamics are governed by a Hamiltonian that is initially invariant under the full symmetric group acting on the vertices.

The Hamiltonian governing this discrete topology typically takes the form:

$$H = H_{\{\text{links}\}} + H_{\{\text{vertices}\}} + H_{\{\text{loops}\}} + H_{\{\text{hop}\}}$$

At ultra-high energies, such as those hypothesized at the birth of the universe or deep within a gravitational singularity, the graph remains maximally connected. Every node is adjacent to every other node, meaning there is no macroscopic concept of distance, locality, or dimensionality, and the physics is entirely invariant under permutation symmetry. However, as the system cools into low-energy states, it undergoes a profound thermodynamic phase transition. The connectivity-sensitive Hamiltonian favors configurations that break permutation symmetry, causing the highly connected graph to settle into a local, ordered, low-dimensional phase.

In this stable, low-energy sector, the suppression of causal loops induces an effective Lorentzian-type geometry with a defined causal structure and a macroscopic signature of the form $(-, +, +, +)$. This topological phase transition maps directly onto condensed matter systems. The discrete spacetime often crystallizes into a honeycomb lattice structure, analogous to graphene, where the Planck-scale discreteness can be approximated at large distances by continuum field theories, such as the Dirac equation. The emergence of space, therefore, is a geometric response to the correlation networks of the underlying quantum tension fields, proving that spatial dimensions are a consequence of field mechanics rather than a prerequisite for them.

Tensor Networks and Holographic Kinematic Space

The bridge between discrete topologies and emergent quantum gravity is rigorously defined by the Anti-de Sitter/Conformal Field Theory (AdS/CFT) correspondence and its tensor network realizations. The Multi-Scale Entanglement Renormalization Ansatz (MERA) serves as a direct quantum mechanical translation of topological geometry.

A MERA network implements real-space renormalization on the wavefunction of a quantum many-body system, utilizing a hierarchical structure of isometric tensors and disentanglers to systematically remove short-range entanglement at each increasing length scale. Structurally, MERA operates as a discretization of kinematic space—the geometry of CFT intervals—which, in holographic theories, is directly identified with the space of bulk geodesics in the dual AdS spacetime. The extra dimension of the AdS bulk corresponds precisely to the renormalization scale within the MERA network.

Through the Ryu-Takayanagi formula, the entanglement entropy $S(A)$ of a boundary region A is proportional to the area of the minimal bulk surface γ_A homologous to A :

$$S(A) = \frac{\text{Area}(\gamma_A)}{4G_N}$$

In the discrete topology of the tensor network, this entropy is bounded by the minimal number of bonds cut to separate region A from the rest of the network, explicitly linking quantum entanglement to geometric connectivity. Thus, reframing the topology into quantum field mechanics reveals that the metric of spacetime is defined by the density of unitary quantum gates and the entanglement entropy produced per unit scale along the continuous MERA (cMERA) flow. Spacetime is, fundamentally, the geometric skeleton of quantum entanglement.

Quantum Field Mechanics of Spacetime Folding

The ER=EPR Conjecture and Thermofield Double States

To manipulate or "fold" spacetime, one must manipulate the underlying entanglement structure. The ER=EPR conjecture posits an exact equivalence between Einstein-Rosen (ER) bridges (spatial wormholes) and Einstein-Podolsky-Rosen (EPR) quantum entanglement. In this framework, the quantum state of a double-sided AdS black hole corresponds precisely to a Thermofield Double (TFD) state in the dual quantum mechanical boundary system:

$$|TFD\rangle = \frac{1}{\sqrt{Z}} \sum_n e^{-\beta E_n / 2} |E_n\rangle_L \otimes |E_n\rangle_R$$

The two boundary CFTs (Left and Right) are maximally entangled but non-interacting. Geometrically, their dual description is an eternal black hole with a spatial wormhole connecting the two interiors. However, classically, this Einstein-Rosen bridge is non-traversable; the wormhole throat stretches faster than light can cross it, maintaining absolute causal separation between the two boundaries.

The Gao-Jafferis-Wall (GJW) Protocol for Traversability

To render the wormhole traversable—effectively folding spacetime to allow information and matter to cross between originally disconnected regions—one must violate the Averaged Null Energy Condition (ANEC). The ANEC dictates that the integral of the stress-energy tensor along any complete null geodesic must be non-negative: $\int_{\gamma} T_{\mu\nu} k^\mu k^\nu d\lambda \geq 0$.

Gao, Jafferis, and Wall (GJW) demonstrated that this topological barrier can be breached using quantum field mechanics. By introducing a non-local double-trace deformation that couples the two asymptotic boundary CFTs, one can generate negative energy shockwaves in the bulk spacetime. The boundary interaction takes the form:

$$\delta H(t) = \int d^{d-1}x \, h(t, \vec{x}) \, O_L(-t, \vec{x}) \, O_R(t, \vec{x})$$

where $O_{L,R}$ are scalar primary operators acting on the left and right boundaries, and $h(t, \vec{x})$ is the coupling function.

When this coupling is activated, the quantum interactions between the boundaries induce a negative averaged null energy in the bulk ($\int_{\gamma} T_{VV} dV < 0$). This negative energy backreacts on the background geometry, causing the event horizons of the black hole to shift inward. In terms of Kruskal coordinates, the shift $\Delta U < 0$ allows a signal originating from the left boundary to fall into the wormhole, avoid the singularity, and emerge in the right boundary's exterior.

This holographic teleportation protocol is heavily governed by information scrambling and out-of-time-ordered correlators (OTOCs). The quantum channel capacity of the GJW traversable wormhole is quantitatively bounded by the time derivative of the OTOC, reflecting the operator size growth in the holographic dual. If the operator growth proceeds at the maximum rate bounded by Einstein gravity, the wormhole achieves maximum channel capacity for teleportation.

Traversable Wormhole Protocol	Channel Properties	Operational Characteristics
Scalar Double-Trace (Original GJW)	Constant coupling h , single primary operator.	Induces negative energy shockwaves; bounded by Planck time windows.
Conserved Current U(1)	Diffusive pole behavior, power-law tail scrambling.	Extended traversal windows dependent on diffusion constants.

Traversable Wormhole Protocol	Channel Properties	Operational Characteristics
Gravitational Shear & Sound Channels	Metric perturbations $h_{\{xt\}}$, $h_{\{xz\}} \neq 0$.	Sound channels show late-time power-law remnants and exponential decay based on sound speed.
Defect-Induced Traversability	Non-invertible symmetry defects in large- c CFTs.	Localized boundary modifications enforcing selection rules beyond standard group symmetries.
SYK Model Implementation	Majorana fermion coupling in $0+1$ dimensions.	Teleportation fidelity peaks post-scrambling; verifiable on quantum simulators.

Recent advancements have expanded the GJW protocol to gravitational shear and sound channels within the hydrodynamic approximation. First-order gravitational perturbations backreact on the metric at second-order, meaning that dynamical metric perturbations alone can facilitate information transfer in a purely gravitational setting. Due to its propagating nature, the sound channel exhibits unique traversability profiles; at low sound speeds, it shows late-time power-law remnants akin to vector diffusive probes. At higher sound speeds, it prefers exponential decay, while superluminal sound channels render the wormhole entirely non-traversable due to the fleeting, unstable duration of the opening. Further theoretical extensions involve multi-boundary black holes, where traversability between any two asymptotic regions can be triggered using double-trace deformations in a "hot limit" (where the temperatures related to all horizons are large), providing a pathway for multi-node topological networks.

Stabilizing Energy Input and Output: Approaching Zero-Point Energy

To actualize the mathematical folding of spacetime described by the GJW protocol in non-holographic, physical laboratory conditions, one must engineer and stabilize physical sources of negative energy density. Classical matter adheres strictly to established energy conditions (Weak, Null, Dominant, Strong), meaning macroscopic negative energy is strictly forbidden by classical general relativity. However, quantum field theory inherently allows for localized suppression of vacuum fluctuations, plunging the local energy density below the zero-point energy of the vacuum state through quantum coherence effects.

Squeezed Vacuum States and Quantum Optics

The most viable experimental pathway to generating the negative energy density required for ANEC violation and spacetime manipulation involves squeezed states of light. The vacuum state of the electromagnetic field is not empty; it consists of zero-point fluctuations. While a standard coherent state (like a laser) possesses quantum noise equal in all quadratures, a squeezed vacuum state reduces the quantum uncertainty in one field quadrature below the standard quantum limit, at the necessary expense of increasing it in the conjugate quadrature to satisfy the Heisenberg uncertainty principle. When the electromagnetic field is manipulated via highly non-linear optical media, it undergoes

parametric down-conversion, altering the fundamental structure of the vacuum such that the energy density is no longer spatially homogeneous. For specific phases, the variance of the electric field drops below that of the bare vacuum. During these phases, the local energy density (ρ_E) becomes strictly negative relative to the zero-point background.

The equation governing the energy density of a squeezed vacuum state for a specific mode exhibits interference terms that drop below zero. By continuously injecting squeezed vacuum into a target spatial region—such as a Casimir cavity or a specially configured Mach-Zehnder interferometer—one generates an effective negative energy gradient. If properly stabilized, this localized negative energy density fulfills the physical requirement to shift the horizons of microscopic topological fluctuations, functioning as a real-world analog to the double-trace deformation in the GJW protocol. The technological challenge lies in sustaining this state without the surrounding positive energy fluctuations overwhelming the localized negative energy well.

Absolute Constraints: Quantum Energy Inequalities (QEIs)

While quantum field mechanics permits the existence of negative energy densities via squeezed states or Casimir effects, nature imposes draconian penalties on its magnitude and duration. If unbounded negative energy were allowed, it would trigger catastrophic macroscopic instabilities, violating the second law of thermodynamics and allowing the unchecked formation of closed timelike curves (time machines) and macroscopic warp drives.

To prevent this architectural collapse of causality, quantum field theory strictly enforces Quantum Energy Inequalities (QEIs). First introduced by Ford and Roman, and subsequently generalized through rigorous algebraic quantum field theory by Fewster, Smith, and Verch, QEIs are state-independent lower bounds on the weighted average of the stress-energy tensor.

The Ford-Roman Bounds and the Microlocal Spectrum Condition

A generic worldline QEI for a massless scalar field limits the locally averaged negative energy density ρ sampled over a characteristic timescale t_0 by a smooth weighting function:

$$\rho \geq -\frac{3}{32} \frac{\hbar c^3}{\pi^2 t_0^4}$$

This inverse fourth-power scaling implies an extreme physical tradeoff: one may create a highly intense pulse of negative energy, but it must be exceedingly brief. Conversely, long-lasting negative energy fields must be vanishingly weak. This bounds the extent to which energy conditions can be violated, proving that quantum negative energy acts more like a high-interest temporal loan than a permanent energy source.

Fewster and Verch proved that QEIs are inexorably linked to the Microlocal Spectrum Condition, which encodes the Hadamard condition for physically reasonable quantum states. A Hadamard state ensures that the short-distance singularity structure of the two-point function mirrors the Minkowski vacuum, which is required to yield finite fluctuations in the Wick-ordered stress-energy tensor. If a quantum state attempts to harbor negative energy beyond the QEI limits, the thermodynamic stability (passivity) of the system is destroyed. The state ceases to be Hadamard, leading to infinite, uncontrollable fluctuations in the stress-energy tensor, rendering the spacetime fundamentally unstable.

To extend these concepts to null vectors (crucial for analyzing light-like traversal of wormholes), physicists developed the Smeared Null Energy Condition (SNEC) and the Double Smeared Null

Energy Condition (DSNEC). Since the un-smearred null energy is unbounded from below, integrating the null energy over two smeared null directions provides a finite lower bound, ensuring that causality-violating geometries are heavily suppressed even in advanced field theories. Furthermore, recent advancements have even extended QEIs into non-commutative spacetimes, ensuring that these energy constraints remain robust even at fundamental scales where standard manifold geometry breaks down.

Implications for Macroscopic Spacetime Engineering

The application of QEIs to the geometry of traversable wormholes and warp drives severely restricts their physical realizability. To maintain a static Morris-Thorne wormhole of throat radius s_0 , an exotic matter distribution must supply a specific negative energy density:

$$\rho = -\frac{c^4}{8\pi G} \frac{s_0^2}{(s_0^2 + l^2)^2}$$

Equating this geometric requirement with the Ford-Roman QEI limit (and setting the sampling time t_0 to the Planck time t_P) yields a lower bound on the throat size of a naturally stable wormhole:

$$s_0^2 \geq \frac{4}{3} l_P^2$$

This calculation demonstrates that while quantum field mechanics allows for the topological folding of spacetime, QEIs strictly constrain naturally stable wormholes to submicroscopic, Planck-scale dimensions

($\sim 10^{-35}$ meters).

Similar constraints plague the Alcubierre warp drive metric. Pfenning and Ford demonstrated that generating an Alcubierre bubble capable of transporting a macroscopic spacecraft at superluminal speeds requires a total negative energy equivalent to roughly 10^{20} solar masses, packed into a bubble wall thinner than 10^{-32} meters, violating all known quantum constraints. While recent theoretical frameworks, such as those proposed by Bobrick and Martire, have identified classes of subluminal, constant-velocity "warp shells" that can be constructed entirely from positive energy—thus bypassing the need to violate energy conditions—these solutions do not enable faster-than-light travel or true topological shortcuts. Any drive that effectively moves superluminally must inevitably break the null energy condition, returning the problem to the insurmountable QEI bounds.

To bypass these restrictions for wormholes, theoretical models suggest utilizing "operational windows" in dynamically expanding backgrounds, where a microscopic wormhole undergoes rapid local inflation. During this brief inflationary phase, the QEI limits are stretched, temporarily enlarging the topological defect to macroscopic scales before the constraints enforce a collapse.

Spacetime Engineering Model	Energy Requirement	QEI Compatibility	Viability Status
Morris-Thorne Wormhole	Sustained negative energy density.	Extremely restricted; naturally limited to Planck-scale throats ($s_0 \sim l_P$).	Submicroscopic only, barring local inflation mechanisms.
Alcubierre Warp Drive (Superluminal)	Astronomical negative mass (10^{20} solar masses).	Grossly violates Ford-Roman bounds; wall thickness $\sim 10^{-32}$ m.	Currently unphysical under standard QFT.
Bobrick-Martire Warp	Positive energy density.	Fully compliant; does	Theoretically

Spacetime Engineering Model	Energy Requirement	QEI Compatibility	Viability Status
Shell (Subluminal)		not violate Null Energy Condition.	permissible, though propulsion mechanics remain unsolved.
GJW Holographic Wormhole	Transient negative energy shockwave via double-trace deformation.	Compliant within localized scattering windows.	Demonstrable in quantum simulations (e.g., SYK model).

Institutional Research Capabilities: Eugene, Oregon

Pioneering research into the stabilization and control of these exotic quantum states, as well as the theoretical modeling of metric backreaction, is deeply embedded in the academic infrastructure of Eugene, Oregon, specifically at the University of Oregon (UO). The integration of applied optical devices with theoretical semiclassical physics makes the UO Physics Department uniquely suited to address the engineering of squeezed vacuums for zero-point energy manipulation and topological investigation.

The Oregon Center for Optical, Molecular, and Quantum Science (OMQ), housed in Willamette Hall, provides an exceptional concentration of expertise across quantum optics, nonlinear optics, and theoretical quantum physics. Within this ecosystem, several specific research groups map directly onto the requirements for realizing the quantum field mechanics of spacetime folding:

- **Raymer Research Group:** Focuses heavily on the quantum mechanics of light, nonlinear optics, and quantum communications technology. Their foundational work on the interaction of light and matter at the quantum level is highly relevant to creating the specific multi-mode squeezed states capable of suppressing vacuum energy over macroscopic volumes.
- **Smith Research Group:** Probes fundamental quantum physics using quantum-enhanced applications to push performance beyond classical limits. The generation of highly non-classical optical states is a prerequisite for maintaining the negative energy wells required by the GJW protocol.
- **Steck Research Group:** Specializes in the physics of ultracold atoms and atom optics. Measuring the minute topological shifts and metric perturbations caused by localized negative energy requires ultra-stable, noise-isolated environments. Ultracold atomic interferometry provides the extreme precision necessary to detect gravitational and topological variations at the micro-scale.
- **Van Enk Research Group:** Provides the theoretical backbone for the study of quantum-information-processing (QIP) protocols. As traversable wormholes are fundamentally quantum channels (where traversal capacity is bounded by OTOCs), Van Enk's research maps directly to the holographic teleportation and entanglement structures (ER=EPR) necessary for computing wormhole channel capacities.
- **Isenberg and Theoretical Relativity:** Professor Jim Isenberg's research into classical and quantum gravity, specifically mathematical relativity and Ricci flow, provides the differential geometry expertise required to model the backreaction of squeezed light on the local metric, moving the research from quantum optics to actual spacetime curvature.
- **Allcock and Wineland Research Group:** Develops hardware and techniques for trapped-ion quantum computing. Simulating the SYK (Sachdev-Ye-Kitaev) model to test

holographic teleportation protocols relies heavily on highly coherent quantum computational systems, which trapped-ion platforms can facilitate.

- **Alemán and McMorran Research Groups:** Provide capabilities in nanoscale systems, defect qubits, and advanced electron microscopy, allowing for the potential fabrication of nano-layered Casimir cavities designed to geometrically enforce negative energy states.

Furthermore, the University of Oregon is a foundational member of the Northwest Quantum Nexus (NQN), a coalition driving scalable quantum computing and Quantum Information Sciences (QIS) across the Pacific Northwest. The cross-pollination between UO's advanced projects laboratories—utilizing state-of-the-art equipment in CAMCOR and the Lokey Laboratories—and quantum information theory makes Eugene a primary node for the experimental verification of topological spacetime engineering.

Research Protocol and Heuristic Prompt

Based on the ontological reframing of mathematical topology into quantum field mechanics, and leveraging the extensive institutional capabilities identified above, the following advanced research prompt is constructed. It is designed to guide computational models, AI reasoning engines, or institutional research teams (such as those integrating UO/OMQ infrastructure) in exploring the stabilization of zero-point energy and the engineering of spacetime.

[BEGIN DEEP RESEARCH PROMPT]

Objective: Synthesize advanced quantum optics (squeezed vacuum states) with holographic quantum gravity (the Gao-Jafferis-Wall protocol) to engineer localized traversable topologies, while strictly optimizing against the boundary constraints of Quantum Energy Inequalities (QEIs) and the Microlocal Spectrum Condition.

Phase 1: Ontological Mapping of Tensor Networks to Physical Systems

1. **AMERA to Holographic Kinematics:** Map the discrete topological tensors of the Multi-Scale Entanglement Renormalization Ansatz (AMERA) to continuous variable (CV) quantum optics. Model the disentanglers and isometries of the AMERA network as cascades of optical parametric amplifiers (OPAs) and highly tuned beam splitters.
2. **ER=EPR Instantiation:** Define a state-preparation protocol to generate a highly entangled macroscopic continuous-variable state, mimicking the Thermofield Double (TFD) state. Establish physically isolated left (L) and right (R) optical cavities to represent the dual boundary CFTs.

Phase 2: Stabilizing Negative Energy via OMQ Architectures

1. **Squeezed Vacuum Generation:** Utilizing the theoretical and experimental frameworks established by the Raymer and Smith research groups, design a highly non-linear pulsed laser configuration to generate deep, multi-mode squeezed vacuum states.
2. **Targeted Energy Suppression:** Direct the squeezed quadrature to interfere with the local zero-point fluctuation background within a micro-fabricated Casimir cavity (integrating nanofabrication techniques from the Alemán and McMorran Labs).
3. **QEI Optimization:** Compute the smeared stress-energy tensor $\langle T_{\mu\nu} \rangle$: $\langle T_{\mu\nu} \rangle = \int g(\tau)^2 d\tau$. Algorithmically optimize the pulse duration (sampling function $g(\tau)$) and the degree of squeezing (r) to maximize the depth of the negative energy density well while remaining strictly compliant with the Fewster-Verch microlocal spectrum condition for Hadamard states, avoiding uncontrolled thermodynamic passivity violations.

Phase 3: The Holographic Spacetime Folding Protocol

1. **Double-Trace Deformation Engineering:** Introduce a non-local coupling function δ

$H = \int h(t, \vec{x}) O_L O_R$ between the L and R isolated optical cavities. Measure the transmission fidelity of a quantum state injected into cavity L, verifying the effective Kruskal coordinate shift ($\Delta U < 0$).

2. **ANEC Violation Detection:** Monitor the system for effective Averaged Null Energy Condition (ANEC) violations. Utilize Out-of-Time-Ordered Correlators (OTOCs) to measure information scrambling and determine the quantum channel capacity of the resultant effective geometry, referencing the theoretical models of the Van Enk group.
3. **Hydrodynamic Channel Extension:** Modulate the non-local coupling to excite effective gravitational shear and sound channels, analyzing the traversal capacity for power-law decay signatures predicted by hydrodynamic GJW models.

Phase 4: Scaling to Macroscopic Topologies

1. **Local Inflation Protocols:** Evaluate mathematical mechanisms (utilizing Jim Isenberg's mathematical relativity frameworks) to bypass static QEI throat-size limitations. Introduce a time-dependent, rapidly expanding background field (an effective local de Sitter phase) to stretch the microscopic negative energy distribution into macroscopic viability.
2. **Stability Analysis:** Assess the temporal duration of the "operational window" before QEI limits force the topological bridge to undergo a pinch-off singularity. Quantify the exact energy input-to-output ratios required to maintain the stability of the zero-point energy suppression during the expansion phase, using ultracold atomic interferometry (Steck Lab) to measure resultant metric fluctuations.

[END DEEP RESEARCH PROMPT]

Conclusion

The transition from classical differential geometry to quantum field mechanics provides the necessary mathematical and physical language to address spacetime engineering. Topologies are no longer static, continuous stages but dynamic entanglement networks governed by holographic dualities and continuous renormalization flows. The Gao-Jafferis-Wall protocol successfully demonstrates that, under precise entanglement architectures, the folding of spacetime to create traversable wormholes is theoretically permissible without requiring exotic, unphysical matter.

However, translating this holographic mathematics into physical reality requires the harnessing of zero-point energy via squeezed vacuum states, a technological frontier heavily reliant on quantum optical research, such as that conducted at the University of Oregon's OMQ. As these technologies mature, they will invariably collide with the absolute limits of nature—Quantum Energy Inequalities and the microlocal spectrum condition—which serve as the thermodynamic guardians of macroscopic causality. The future of spacetime engineering relies entirely on finding transient, dynamic loopholes within these bounds, utilizing rapid local inflation and non-local double-trace deformations to systematically push the boundaries of quantum field mechanics.

Works cited

1. Quantum graphity: A model of emergent locality - ResearchGate, https://www.researchgate.net/publication/1905944_Quantum_Graphity_a_model_of_emergent_locality
2. [PDF] Quantum Graphity - Semantic Scholar, <https://www.semanticscholar.org/paper/7392e5092d2e3fd263cc31b5d06ef842ba729c56>
- 3.

arXiv:hep-th/0611197v1 17 Nov 2006,
<http://alpha.sinp.msu.ru/~panov/Cosmology/Referats/0611197v1.pdf> 4. NewScientist -
[homepages.ucl.ac.uk, http://www.homepages.ucl.ac.uk/~ucapsse/NewScientist08.pdf](http://www.homepages.ucl.ac.uk/~ucapsse/NewScientist08.pdf) 5. (PDF)
 Domain structures in quantum graphity - ResearchGate,
https://www.researchgate.net/publication/221941008_Domain_structures_in_quantum_graphity
 6. Tensor Networks from Kinematic Space - arXiv, <https://arxiv.org/html/1512.01548v1> 7.
 arXiv:1209.3304v1 [hep-th] 14 Sep 2012, <https://arxiv.org/pdf/1209.3304> 8. On the
 Entanglement Entropy in Gaussian cMERA - arXiv, <https://arxiv.org/html/2104.01551v2> 9.
 Connecting entanglement renormalization and gauge/gravity dualities,
<https://arxiv.org/pdf/1109.5592> 10. Tensor networks for dynamic spacetimes (DRAFT - arXiv,
<https://arxiv.org/pdf/1709.09787> 11. arXiv:1808.09072v3 [hep-th] 23 Nov 2018,
<https://arxiv.org/pdf/1808.09072> 12. Traversable wormhole with double trace deformations via ...
 - arXiv, <https://arxiv.org/html/2511.09815v2> 13. Wormhole and Entanglement (Non-)Detection in
 the ER=EPR,
https://www.researchgate.net/publication/282000209_Wormhole_and_Entanglement_Non-Detection_in_the_ER=EPR_Correspondence 14. Quantum Channel Capacity of Traversable
 Wormhole - arXiv, <https://arxiv.org/pdf/2603.26051> 15. Gravitational Wave-Induced Scrambling
 Delay in SYK Wormhole, <https://arxiv.org/pdf/2603.18509> 16. JHEP06(2021)134,
<https://d-nb.info/1247603016/34> 17. JHEP01(2020)050, <https://d-nb.info/1208298054/34> 18.
 Worldsheet traversable wormholes - ProQuest,
<https://search.proquest.com/openview/1a3f051326546fb570e579473d078876/1?pq-origsite=gsc>
[holar&cbl=2034718](https://search.proquest.com/openview/1a3f051326546fb570e579473d078876/1?pq-origsite=gsc&cbl=2034718) 19. A traversable wormhole teleportation protocol in the SYK model,
https://www.researchgate.net/publication/353277129_A_traversable_wormhole_teleportation_protocol_in_the_SYK_model 20. Traversable wormhole with double trace deformations via ... -
 CERN,
https://scoap3-prod-backend.s3.cern.ch/media/harvested_files/10.1103/4rqm-nmx9/4rqm-nmx9.pdf 21. (PDF) Traversable wormhole with double trace deformations via,
https://www.researchgate.net/publication/397595466_Traversable_wormhole_with_double_trace_deformations_via_gravitational_shear_and_sound_channels 22. JHEP04(2021)083,
<https://d-nb.info/1244098779/34> 23. Lectures on quantum energy inequalities
 arXiv:1208.5399v1 [gr-qc, <https://arxiv.org/pdf/1208.5399> 24. Tractor Beams, Pressor Beams
 and Stressor Beams in General, <https://www.mdpi.com/2218-1997/7/8/271> 25. Negative Energy
 Densities in Quantum Field Theory - Academia.edu,
https://www.academia.edu/91588968/Negative_Energy_Densities_in_Quantum_Field_Theory
 26. Extraction of Thrust from Quantum Vacuum Using Squeezed Light,
https://pubs.aip.org/aip/acp/article-pdf/880/1/1034/11386575/1034_1_online.pdf 27. Quantum
 Inequalities from the Second Law - arXiv, <https://arxiv.org/html/2608.11817v1> 28. Squeezed
 Light and Negative Energy Density Quantum States,
<https://www.hathawayresearch.com/portfolios/squeezed-light-negative-energy-density-quantum-states/> 29. (PDF) Experimental Concepts for Generating Negative Energy in the,
https://www.researchgate.net/publication/228623774_Experimental_Concepts_for_Generating_Negative_Energy_in_the_Laboratory 30. Experimental Concepts for Generating Negative
 Energy in the,
https://wiki.naturalphilosophy.org/index.php/Experimental_Concepts_for_Generating_Negative_Energy_in_the_Laboratory 31. Squeezed coherent state - Wikipedia,
https://en.wikipedia.org/wiki/Squeezed_coherent_state 32. A bridge between the single-photon
 and squeezed-vacuum states, <https://opg.optica.org/abstract.cfm?uri=oe-18-17-18254> 33.
 arXiv:0705.3003v1 [quant-ph] 21 May 2007, <https://arxiv.org/pdf/0705.3003> 34. Engineering

stress-energy signs in a subluminal Alcubierre analog,
https://www.researchgate.net/publication/404295430_Engineering_stress-energy_signs_in_a_subluminal_Alcubierre_analog_using_Casimir_cavities_and_magnetized_plasma 35. Parameter estimation of wormholes beyond the Heisenberg limit, <https://arxiv.org/html/1811.02395v2> 36. On wormholes with arbitrarily small quantities of exotic matter,
https://www.researchgate.net/publication/1969769_On_wormholes_with_arbitrarily_small_quantities_of_exotic_matter 37. quantum energy inequalities in two-dimensional conformal field theory, <https://www.worldscientific.com/doi/10.1142/S0129055X05002406> 38. Wormhole Restrictions from Quantum Energy Inequalities - MDPI,
<https://www.mdpi.com/2218-1997/10/7/291> 39. Energy inequalities in quantum field theory,
https://www.worldscientific.com/doi/pdf/10.1142/9789812704016_0056?download=true 40. Local Expansion Mechanisms for Quantum-Scale Wormholes - arXiv,
<https://arxiv.org/pdf/2509.10137> 41. Semiclassical energy conditions for quantum vacuum states - arXiv, <https://arxiv.org/pdf/1306.2076> 42. Quantum Inequalities From the Second Law - Semantic Scholar,
<https://www.semanticscholar.org/paper/Quantum-Inequalities-From-the-Second-Law-Palessandro/b7b9dba6cddb85852bd334cfbdd2ead2fe58fd6a> 43. The necessity of the Hadamard Condition, <https://eprints.whiterose.ac.uk/id/eprint/84763/1/Necessity.pdf> 44. arXiv:math-ph/0502002v1 1 Feb 2005, <https://arxiv.org/pdf/math-ph/0502002> 45. Quantum Energy Inequalities, Part I - Non Sequitur,
<https://nonsequitur.blog/2020/12/07/quantum-energy-inequalities-part-i/> 46. A singularity theorem for evaporating black holes,
https://www.worldscientific.com/doi/pdf/10.1142/9789811269776_0064 47. The double smeared null energy condition - SciPost.org, https://www.scipost.org/preprints/scipost_202202_00025/ 48. A Quantum Energy Inequality for a Non-commutative QFT - arXiv,
<https://arxiv.org/pdf/2503.24068> 49. (PDF) A Quantum Energy Inequality for a Non-commutative QFT,
https://www.researchgate.net/publication/390404760_A_Quantum_Energy_Inequality_for_a_Non-commutative_QFT 50. Findings - Jump Drive, <https://jump-drive.com/findings/> 51. The Metric Tensor, Warp Drives, and Wormholes, <https://spacetime-metric.com/chapters/ch04> 52. Warp drives and the negative-energy problem, <https://warpdrives.org/negative-energy/> 53. [PDF] Introducing physical warp drives - Semantic Scholar,
<https://www.semanticscholar.org/paper/Introducing-physical-warp-drives-Bohrick-Martire/62a78c249435213216fea046dead74dcb5c7a525> 54. About Us - Optical Molecular & Quantum Science, <https://omq.uoregon.edu/about-us> 55. Research Physics | Natural Sciences - University of Oregon, <https://naturalsciences.uoregon.edu/physics/research> 56. Research Groups | Optical Molecular & Quantum Science, <https://omq.uoregon.edu/research-groups> 57. Academic - Daniel A. Steck, <https://steck.us/academic.html> 58. S. VAN ENK | Professor | PhD | University of Oregon, Eugene | UO, <https://www.researchgate.net/profile/S-Van-Enk/2> 59. Jim ISENBERG | University of Oregon, Eugene | UO | Research profile,
<https://www.researchgate.net/profile/Jim-Isenberg-2> 60. PCGM 31 - CCoM,
<https://ccom.ucsd.edu/~pcgm31/> 61. Physics Faculty | College of Arts and Sciences - University of Oregon, <https://cas.uoregon.edu/directory/physics-faculty> 62. Research Opportunities Physics | Natural Sciences,
<https://naturalsciences.uoregon.edu/physics/experiential-learning/research>