

# Deep holes of projective Reed–Solomon codes beyond redundancy four: exact classifications at redundancies five through seven

Tavis Rudd

Unrefereed preprint, Version 1  
July 2026

## Abstract

We classify, up to projective-semilinear equivalence, the deep-hole syndrome directions of projective Reed–Solomon codes at redundancies five and six over every prime power  $q \geq 7$ , with exact projective counts. At redundancy seven we give the complete split-free classification over the same range; it is a deep-hole classification whenever the covering radius is six, in particular for  $q \geq 11$ . At redundancy five the list contains, besides the tangent and conjugate-secant families, an osculating-pair family selected by  $q \bmod 3$ , a characteristic-three nucleus point and wild Artin–Schreier family, and sporadic orbits exactly for  $q \in \{7, 8, 9, 11, 13, 17, 19\}$ . A syndrome determines a Hankel linear system of binary forms and is split-free precisely when the system contains no completely split squarefree form. For cubic pencils, cyclic descent produces the arithmetic families, a genus-one fiber square controls the  $S_3$  case, and exact certificates settle the remaining fields. The balanced  $q = 8$  row gives 1116  $[10, 5, 6]_8$  MDS extensions and hence minimum-support  $\text{AME}(10, 8)$  states and  $[[9, 1, 5]]_8$  quantum MDS codes. For redundancies six and seven we introduce coherent polar flags. A contraction removes a prospective root; retaining it as a forbidden marker is exactly what prevents a lower squarefree witness from lifting with a double root. Complete contained-component calculations and one- and two-marker point counts give the uniform ranges  $q \geq 29$  and  $q \geq 37$ , respectively. Finite certificates close the remaining fields and record the exceptional orbits, stabilizers, and Frobenius fusions. At redundancy seven a second direct-locus enumerator independently reconstructs the split-free set and orbit partition. No contained-component claim is made at arbitrary redundancy.

**Index Terms**—Absolutely maximally entangled states, algebraic coding theory, deep holes, finite fields, maximum-distance-separable extensions, normal rational curves, projective Reed–Solomon codes, quantum MDS codes.

## 1 Introduction

This paper gives two complete deep-hole classifications over all prime powers  $q \geq 7$ , beyond the known cases through redundancy four. At redundancies five and six we classify the projective deep-hole directions for every prime power  $q \geq 7$ , with exact projective counts and semilinear orbit laws. At redundancy seven we classify every split-free syndrome over the same range; this is a deep-hole classification whenever the covering radius is six, in particular for  $q \geq 11$ . The common mechanism is a coherent polar flag, a marked contraction that propagates splitting information without forgetting the factor needed for a squarefree lift.

Let  $\text{PRS}(k)$  denote the projective Reed–Solomon code of length  $q+1$  obtained by evaluating homogeneous binary forms of degree  $k-1$  on  $\mathbb{P}^1(\mathbb{F}_q)$ . If the redundancy is

$$r = q + 1 - k,$$

a parity-check matrix has as columns the  $\mathbb{F}_q$ -points of the degree- $(r-1)$  normal rational curve

$$\mathcal{R}_{r-1} = \{(1, t, \dots, t^{r-1}) : t \in \mathbb{F}_q\} \cup \{(0, \dots, 0, 1)\} \subset \mathbb{P}^{r-1}.$$

Call a projective syndrome direction *split-free* if it is not contained in the span of  $r-2$  distinct columns. When  $\rho(\text{PRS}(q+1-r)) = r-1$ , the split-free directions are precisely the projective syndrome directions of deep-hole cosets. The logical separation used throughout the paper is

|                                                                                                              |
|--------------------------------------------------------------------------------------------------------------|
| $\text{split-free classification} + \rho(\text{PRS}(q+1-r)) = r-1 \implies \text{deep-hole classification.}$ |
|--------------------------------------------------------------------------------------------------------------|

The first term is proved geometrically; the covering-radius equality is a separate coding-theoretic input and is never inferred from a finite split-free census.

The correspondence between deep holes, one-symbol MDS extensions, and uncovered points of a normal rational curve was developed in [Kai17, Prop. 1] and [ZWK20, Lem. II.4 and Thms. I.4–I.6]. Zhang, Wan, and Kaipa classified projective Reed–Solomon deep-hole directions through redundancy four and identified redundancy five as the next case [ZWK20]. Theorem 1.1 closes that case; Theorem 1.2 gives the complete redundancy-six classification and the radius-delimited redundancy-seven result.

**Theorem 1.1** (Complete redundancy-five classification). Let  $q \geq 7$  be a prime power. The covering radius of  $\text{PRS}(q-4)$  is 4. Its projective deep-hole syndromes are exactly the following disjoint families:

- (i) the tangent family  $T$ , of size  $q(q+1)$ ;
- (ii) the conjugate-secant family  $S$ , of size  $q(q^2-1)/2$ ;
- (iii) if  $\text{char } \mathbb{F}_q \neq 3$ , the rational osculating-pair family  $O^+$  when  $q \equiv 2 \pmod{3}$ , of size  $q(q+1)/2$ , or the conjugate osculating-pair family  $O^-$  when  $q \equiv 1 \pmod{3}$ , of size  $q(q-1)/2$ ;
- (iv) in characteristic three, one  $\text{PGL}_2(q)$ -fixed nucleus point and one wild Artin–Schreier orbit  $W$  of size  $(q^2-1)/2$ ;
- (v) the sporadic  $\text{PGL}_2(q)$ -orbits in Table 4, occurring exactly for  $q \in \{7, 8, 9, 11, 13, 17, 19\}$ .

There are no other deep-hole syndromes. The three corresponding nonsporadic totals are

$$\frac{q^2(q+3)}{2}, \quad \frac{q(q+1)(q+2)}{2}, \quad \frac{q^3+3q^2+q+1}{2},$$

according as  $q \equiv 1 \pmod{3}$  with characteristic not three,  $q \equiv 2 \pmod{3}$ , or  $\text{char } \mathbb{F}_q = 3$ .

The Hankel dictionary turns redundancy five into a pencil of cubics. Cyclic descent gives the arithmetic families in Theorem 1.1; the  $S_3$  case is controlled by the off-diagonal fiber square and a genus-one point bound. At higher redundancy, contraction lowers the degree, but the removed root must remain forbidden: restoring a lower witness through that root creates a double factor. Coherent polar flags retain precisely this datum. The three-row reading map in Section 2 identifies the load-bearing argument for each redundancy.

The resulting structural theorem has two outputs.

**Theorem 1.2** (Redundancy-six deep holes and redundancy-seven split-free syndromes). The following statements hold.

- (a) For every prime power  $q \geq 7$ , the projective deep-hole directions of  $\text{PRS}(q-5)$  are exactly the persistent tangent and conjugate-secant families, the exceptional orbits listed for  $q = 7, 8, 9, 11, 13$ , and, when  $q = 2^m$  with odd  $m \geq 5$ , one characteristic-two nucleus orbit. Here  $m \geq 5$  avoids double counting:  $m = 3$ , or  $q = 8$ , is already the  $|\mathcal{O}| = 9$  row of the exceptional table.

- (b) The displayed finite domains through  $q = 32$  have the certified split-free classification stated in Section 6. Together with the structural theorem, these finite classifications give the all-prime-power result: for every  $q \geq 13$  the split-free locus consists of the persistent families and, when  $q = 2^m$  with  $m$  odd, one central nucleus point. This is a deep-hole classification only when  $\rho(\text{PRS}(q - 6)) = 6$ , in particular for  $q \geq 11$ .

At redundancy  $r$ , the semilinear orbits in the persistent conjugate-secant family are parameterized by

$$T/T^{r-1} \quad \text{modulo inversion and Frobenius,}$$

where  $T = \{z \in \mathbb{F}_{q^2}^\times : z^{q+1} = 1\}$  is the norm-one torus and Frobenius is coefficient Frobenius. On the tangent family the unipotent coordinate transforms as  $z \mapsto z + (r - 1)u$ ; consequently its orbit decomposition changes precisely when  $\text{char } \mathbb{F}_q$  divides  $r - 1$ .

The two parts have different scopes. Both are unconditional syndrome classifications over every prime power  $q \geq 7$ , but part (b) becomes a deep-hole classification only when  $\rho(\text{PRS}(q - 6)) = 6$ . Table 2 places the split-free result, radius gate, deep-hole conclusion, and remaining field gap in separate columns for each redundancy.

**Relation to previous work.** The tangent and conjugate-secant families and the coding/normal-rational curve dictionary come from [Kai17, Prop. 1] and [ZWK20, Lem. II.4 and Thms. I.4–I.6]; the covering-radius input in the high-rate range combines the Seroussi–Roth extension theorem [SR86, Thm. 1] with Dür’s completeness–covering-radius equivalence [Dür94], as stated by Kaipa [Kai17, Sec. IV and Prop. 4(1)]. The broader arc/MDS and code-automorphism background is surveyed or established in [BL19, Dür87]. Earlier explicit-family results for projective and generalized projective Reed–Solomon codes include [ZW17, XHX17, Xu23]; the recent extension framework of [WDC23] further clarifies the deep-hole/MDS-extension interface. These works construct families, treat punctured or lower-redundancy regimes, or analyze the extension criterion; they do not supply the complete redundancy-five cubic-pencil exhaustion proved here. Classical apolarity, catalecticants, and binary-form rank strata are treated in [IK99, CS11]; our use of divided powers keeps the small-characteristic contraction formulas explicit. Kaipa, Patanker, and Pradhan classify the  $\text{PGL}_2(q)$ -orbits of binary quartic forms and of lines relative to the twisted cubic [KPP25]. Their orbit classification is adjacent normal-form geometry for the redundancy-six problem; it does not impose the Hankel split-free condition or determine the corresponding deep-hole locus. Three companion papers provide comparisons with relative conic completion, inverse rigidity, and conic-ideal factorization memory [Rud26a, Rud26d, Rud26c]; none is an input here. The MDS–CSS companion supplies the quantum consequences of the balanced  $q = 8$  row stated in Section 4 [Rud26b]. From the viewpoint of binary forms, the question is which linear systems contain a totally split squarefree member. Factorization statistics in linear families [CMP17] and normal-rational-curve nuclei [GH00, Lem. 1 and Thm. 1] are inputs. For splitting families, we use Wang’s Theorem 3.6 [Wan26]: on the étale locus, the factorization type of a specialized polynomial is the cycle type of its Frobenius conjugacy class. The coding-theory references above identify deep syndromes with uncovered points of a normal rational curve and settle the lower-redundancy families; they do not classify the redundancy-five cubic pencils. The orbit-theoretic and factorization references supply normal forms and distribution tools, but not the coherent marked contraction used here. The nucleus results locate characteristic-dependent kernels, while the present work computes which of their coherent lifts meet the PRS splitting incidence. Accordingly, this paper’s contributions are the redundancy-five through seven classifications and the coherent marked contraction that proves the uniform ranges at redundancies six and seven.

## 2 Reading map

The proof can be read at two speeds. For the classification spine, follow the three rows of Table 1; the intervening definitions and lemmas discharge the cited carrier, collision, and point-count steps. Table 2 separately records where a split-free classification needs a covering-radius input before it becomes a deep-hole classification.

Table 1: Fast path through the classification proofs.

| Result | Load-bearing printed argument                                                                                                                       | Imported input                                                                |
|--------|-----------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------|
| R5     | Hankel dictionary; gcd strata; cyclic descent; integral (2, 2) fiber square and point bound                                                         | lower tangent/secant families; high-rate radius theorem; singular-curve bound |
| R6     | one-marker polar line; ordinary and binary contained components; transverse escape                                                                  | R5 lower package and radius gate                                              |
| R7     | two-marker contraction; complete second-marker bad scheme, including cyclic and fixed-gcd collisions; contained rank-two theorem; transverse escape | R5/R6 lower packages; separate covering-radius result                         |

Table 2: Classification status by redundancy. “Unconditional” refers to the displayed field range. A split-free classification becomes a deep-hole classification only after the separate radius gate.

| Red. | Field range | Split-free geometry     | Deep holes/MDS<br>sions                                               | exten-<br>Remaining gap |
|------|-------------|-------------------------|-----------------------------------------------------------------------|-------------------------|
| 5    | $q \geq 7$  | Complete, unconditional | Complete, $\rho = 4$                                                  | unconditional; None     |
| 6    | $q \geq 7$  | Complete, unconditional | Complete, $\rho = 5$                                                  | unconditional; None     |
| 7    | $q \geq 7$  | Complete, unconditional | Complete for $q \geq 11$ ; conditional on $\rho = 6$ at $q = 7, 8, 9$ | Radius at $q = 7, 8, 9$ |

## 3 The syndrome–Hankel dictionary

Let  $E$  be a two-dimensional vector space. We use divided powers for syndromes and ordinary symmetric powers for root forms. This avoids invalid binomial rescalings in small characteristic. For  $f = (a_0 : \dots : a_{r-1}) \in \mathbb{P}(\Gamma^{r-1}E)$ , write  $e_0, \dots, e_{r-1}$  for the divided-power coordinate basis and  $\nu(t) = (1, t, \dots, t^{r-1})$ , with  $\nu(\infty) = e_{r-1}$ . If  $x, y$  is a basis of  $E$  with dual basis  $X, Y$ , our perfect divided-power/symmetric-power pairing is

$$\langle x^{[n-i]}y^{[i]}, X^{n-j}Y^j \rangle = \delta_{ij}, \quad \left\langle \sum_{i=0}^n a_i x^{[n-i]}y^{[i]}, \sum_{i=0}^n b_i X^{n-i}Y^i \right\rangle = \sum_{i=0}^n a_i b_i.$$

Thus the pairing is coefficient extraction, with no binomial factors in any characteristic. Define

$$H_f = \begin{pmatrix} a_0 & a_1 & \cdots & a_{r-2} \\ a_1 & a_2 & \cdots & a_{r-1} \end{pmatrix}, \quad W_f = \ker H_f \subset \text{Sym}^{r-2} E^\vee.$$

Read  $f = (a_0, \dots, a_{r-1})$  as a finite sequence. A form  $g \in W_f$  is the characteristic polynomial of a recurrence satisfied by that sequence, while a squarefree split  $g$  has a basis of geometric-sequence solutions. Thus split-free means that the syndrome sequence satisfies no order- $(r-2)$  recurrence with squarefree completely split characteristic polynomial.

Table 3: Notation used from the Hankel dictionary onward.

| Symbol                                | Meaning                                                             |
|---------------------------------------|---------------------------------------------------------------------|
| $q$                                   | field size; all numerical thresholds are thresholds on prime powers |
| $k$                                   | code dimension in PRS( $k$ )                                        |
| $r = q + 1 - k$                       | redundancy                                                          |
| $n = r - 1$                           | divided-power degree of a syndrome representative                   |
| $E$                                   | a two-dimensional vector space over the current base field          |
| $f \in \Gamma^n E$                    | divided-power representative of a projective syndrome               |
| $W_f \subset \text{Sym}^{n-1} E^\vee$ | Hankel/apolar witness system annihilated by $f$                     |
| $g, \delta, \kappa$                   | genus, deletion degree, and point-bound correction                  |
| $d$                                   | degree of the unavailable marker scheme                             |

**Lemma 3.1** (Hankel criterion). Let  $\lambda_1, \dots, \lambda_{r-2}$  be distinct points of  $\mathbb{P}(E^\vee)(\mathbb{F}_q)$  and put  $g = \prod_i \lambda_i$ . Then

$$f \in \text{span}(\nu(\lambda_1), \dots, \nu(\lambda_{r-2})) \iff g \in W_f.$$

The same statement with repeated factors and Hasse derivatives gives the corresponding osculating spans. Both statements commute with  $\text{PTL}_2(q)$ .

*Proof.* On an affine chart, the geometric sequences  $\nu(\lambda_i)$  solve the recurrence with characteristic polynomial  $g$ ; their independence is the Vandermonde determinant. A root at infinity gives the reversed-chart recurrence. For repeated roots, the confluent Vandermonde basis is formed by Hasse derivatives, so the same argument works in every characteristic. Equivariance follows because substitution transports both the normal rational curve and the recurrence.  $\square$

**Corollary 3.2** (Split-free criterion). Assume  $\rho(\text{PRS}(q+1-r)) = r-1$ . A projective syndrome  $f$  is deep if and only if  $W_f$  contains no completely split squarefree form of degree  $r-2$  over  $\mathbb{F}_q$ .

We use *split-free* without a covering-radius hypothesis and call a direction *shallow* when  $W_f$  contains a completely split squarefree form.

The first locus that persists under contraction is already visible here. If every member of  $W_f$  has a common quadratic factor, then that factor is either a rational square, an irreducible quadratic, or a product of two distinct rational factors. These are respectively the tangent deep family, the conjugate-secant deep family, and a shallow rational secant. The common factor cannot grow with redundancy. If  $H_f$  has rank one, then  $\dim W_f = r-2$ , which is incompatible with a common quadratic factor because the space of degree- $(r-2)$  forms divisible by that factor has dimension  $r-3$ . Otherwise  $\dim W_f = r-3$ , and

$$r-3 \leq r-1 - \deg \gcd(W_f),$$

so its degree is at most two.

### 3.1 Coding consequences and recognition

The dictionary has two immediate coding consequences. First, a split-free direction is an uncovered point of the normal rational curve and hence gives a one-column MDS extension whenever the stated covering-radius gate applies. The classification theorems therefore give exact projective deep-hole counts, not only normal forms. Second, they give recognition procedures. From a syndrome one forms  $H_f$ , computes  $W_f$ , and tests the listed geometric families: a common quadratic factor detects the persistent tangent and conjugate-secant families; the relevant contraction kernel detects the modular locus; and the finite exceptional fields are

decided by the canonical orbit representatives printed in Table 4 and recorded in the supplement.

Recognition at the fixed redundancies proved here begins with row-reduction of  $H_f$  and the listed family-membership tests. Coherent polar induction replaces a direct search through the projective members of  $W_f$ .

## 4 Redundancy five: exceptional cubic covers

Here  $f = (a_0 : \cdots : a_4)$  and  $W_f$  is the pencil annihilated by

$$\begin{pmatrix} a_0 & a_1 & a_2 & a_3 \\ a_1 & a_2 & a_3 & a_4 \end{pmatrix}.$$

These are divided-power coordinates; the coefficients of a cubic in  $W_f$  are ordinary symmetric-power coordinates. Factoring the formal quartic  $\sum a_i T^i U^{4-i}$  would therefore be noninvariant in characteristics two and three.

**Proposition 4.1** (Radius gate). For every prime power  $q \geq 7$ ,

$$\rho(\text{PRS}(q-4)) = 4.$$

*Proof.* The coding/arc criterion identifies radius four with completeness of the quartic normal rational curve. The Seroussi–Roth extension theorem [SR86, Thm. 1], applied to the dual five-dimensional GRS code, gives nonextendability when

$$5 \leq \left\lfloor \frac{q}{2} \right\rfloor + 2,$$

apart from its even-field dimension-three exception. The inequality holds for  $q \geq 7$ , and the exception does not occur. Dür’s completeness–covering-radius equivalence [Dür94], as stated by Kaipa [Kai17, Sec. IV and Prop. 4(1)], now gives radius four. Thus Corollary 3.2 applies. The finite certificate independently checks the same conclusion only in its declared finite domain; it is not the all-field radius proof.  $\square$

### 4.1 Gcd strata

**Proposition 4.2** (Quadratic gcd). If  $\deg \gcd(W_f) = 2$ , then

$$W_f = Q\langle T, U \rangle.$$

If  $Q$  is split squarefree,  $f$  is shallow. If  $Q$  is irreducible,  $f$  belongs to the conjugate-secant family; if  $Q = \lambda^2$ , it belongs to the tangent family. Conversely every point in those families has the displayed gcd. Their deep-family sizes are

$$|T| = q(q+1), \quad |S| = \frac{q(q^2-1)}{2}.$$

*Proof.* The equality follows by dimension. A split  $Q$  and a linear cofactor avoiding its two roots give a split squarefree cubic. An irreducible  $Q$  survives in every member, while a square  $Q$  forces a repeated root in every member. Lemma 3.1, including its confluent and conjugate-root forms, identifies the corresponding syndromes with the secant and tangent spans. The sizes are the established tangent/conjugate-secant counts from the lower-redundancy classification [Kai17, Prop. 1] and [ZWK20, Thms. I.4–I.6].  $\square$

**Proposition 4.3** (Linear gcd). Suppose  $\deg \gcd(W_f) = 1$ . Then  $f$  is shallow for  $q \geq 8$ . At  $q = 7$  the same conclusion is part of Certificate R5.

*Proof.* Write  $W_f = \lambda_0 V$  with  $V$  a basepoint-free quadratic pencil. If its degree-two map  $\psi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$  is separable, its rational deck involution  $\iota$  gives the graph

$$\Gamma_\iota(\mathbb{F}_q) = \{(t, \iota(t)) : t \in \mathbb{P}^1(\mathbb{F}_q)\},$$

which has exactly  $q + 1$  rational points. A safe union bound deletes at most two fixed graph points, at most four graph points over the branch fibers, and at most the two graph points  $(\lambda_0, \iota(\lambda_0))$  and  $(\iota(\lambda_0), \lambda_0)$ . Thus fewer than  $q + 1$  graph points are deleted when  $q \geq 8$ . A surviving point gives a split quadratic with two distinct roots avoiding  $\lambda_0$ , and hence a split squarefree cubic in  $W_f$ . In characteristic two an inseparable quadratic pencil is equivalent to  $\langle T^2, U^2 \rangle$ ; the two overlapping Hankel equations then force  $f$  onto the normal rational curve, contrary to the rank-two pencil hypothesis.  $\square$

## 4.2 The trivial-gcd cubic cover

Choose a basis  $c_1, c_2$  of a trivial-gcd pencil and put

$$X_f = \{([u : v], t) : uc_1(t) + vc_2(t) = 0\} \subset \mathbb{P}^1 \times \mathbb{P}^1.$$

**Proposition 4.4** (Cubic incidence and off-diagonal fiber square). The curve  $X_f$  is geometrically integral of bidegree  $(1, 3)$  and arithmetic genus zero. Projection to the root coordinate is birational, while projection to the pencil coordinate gives a degree-three morphism

$$\phi_f : \mathbb{P}^1 \longrightarrow \mathbb{P}^1.$$

If  $\phi_f$  is separable, division of

$$c_1(t_1)c_2(t_2) - c_2(t_1)c_1(t_2)$$

by the diagonal bracket defines a residual curve  $Y_f \subset \mathbb{P}^1 \times \mathbb{P}^1$  of bidegree  $(2, 2)$  and arithmetic genus one. Its geometric components are the geometric-monodromy orbits on ordered pairs of distinct roots.

*Proof.* A component of  $X_f$  of bidegree  $(0, e)$  would be a common factor of  $c_1, c_2$ ; hence none exists. The bidegree genus formula gives genus zero. A root determines the unique pencil member containing it, so the root projection is birational. For a separable map the diagonal factor in the fiber-square determinant has multiplicity one. Removing its bidegree  $(1, 1)$  from the fiber square leaves bidegree  $(2, 2)$ , whose arithmetic genus is one. The standard fiber-square/monodromy correspondence identifies its components with orbits on distinct ordered sheets.  $\square$

**Lemma 4.5** (Cyclic stratum). Suppose  $\phi_f$  is geometrically cyclic.

- (i) In characteristic different from three there are two ramification points. If they are rational, the corresponding osculating-plane intersection is deep exactly when  $q \equiv 2 \pmod{3}$ ; if they are conjugate, it is deep exactly when  $q \equiv 1 \pmod{3}$ .
- (ii) In characteristic three all osculating planes meet in  $n = (0 : 0 : 1 : 0 : 0)$ , whose pencil is  $\langle T^3, U^3 \rangle$ ; this fixed point is deep.
- (iii) The remaining wild characteristic-three pencils have the form  $\langle U^3, T^3 + aTU^2 \rangle$ . They are deep exactly when  $-a$  is nonsquare, forming one orbit of size  $(q^2 - 1)/2$ .

*Proof. Tame characteristic.* Assume that the characteristic is not three. Riemann–Hurwitz gives two totally ramified points  $t_1, t_2$ . The confluent Hankel criterion places  $f$  in  $\text{Osc}(t_1) \cap \text{Osc}(t_2)$ ; after transporting the pair to  $\{0, \infty\}$ , the pencil is  $\langle T^3, U^3 \rangle$  and a geometric deck generator is  $t \mapsto \zeta t$ .

For a rational ramification pair, Frobenius fixes the points and the deck map descends exactly when  $\zeta^q = \zeta$ , namely  $q \equiv 1 \pmod{3}$ . Thus its complement is deep exactly for  $q \equiv 2 \pmod{3}$ . The setwise stabilizer of the ordered normal form is the split dihedral group of order  $2(q-1)$ , so the orbit size is

$$\frac{q(q^2-1)}{2(q-1)} = \frac{q(q+1)}{2}.$$

For a conjugate pair Frobenius interchanges the fixed points and inverts the multiplier; descent is instead  $q \equiv 2 \pmod{3}$ . The nonsplit dihedral stabilizer has order  $2(q+1)$  and the orbit size is  $q(q-1)/2$ .

*The fixed wild point.* Suppose the characteristic is three. The second Hasse derivative of

$$\nu(t) = (1, t, t^2, t^3, t^4)$$

is  $(0, 0, 1, 3t, 6t^2) = e_2$ , including in the reversed infinity chart. Thus every osculating plane contains  $n = [e_2]$ . Substitution shows that  $n$  is  $\mathrm{PGL}_2(q)$ -fixed and  $W_n = \langle T^3, U^3 \rangle$ , so it is deep.

*The remaining wild orbit.* A remaining separable cyclic cover has one wild totally ramified point. Sending it to infinity and comparing coefficients under a nontrivial translation gives

$$W_f = \langle U^3, T^3 + \alpha T^2 U + \beta T U^2 \rangle, \quad \alpha = 0, \quad \kappa^2 = -\beta.$$

It is therefore represented by

$$f_a = e_2 - a e_4, \quad W_{f_a} = \langle U^3, T^3 + a T U^2 \rangle.$$

The affine members are  $z^3 + az + s$ . Such a member has three rational roots exactly when the additive map  $z \mapsto z^3 + az$  has a nonzero rational kernel, equivalently when  $-a$  is a square. Hence the nonsquare parameters form the deep orbit. Its stabilizer is  $t \mapsto \pm t + \beta$ , of order  $2q$ , and orbit-stabilizer gives  $(q^2-1)/2$ .  $\square$

**Lemma 4.6** ( $S_3$  stratum). If  $\phi_f$  has geometric monodromy  $S_3$  and  $q \geq 23$ , then the pencil contains a completely split squarefree cubic.

*Proof.* By Proposition 4.4, components of  $Y_f$  correspond to monodromy orbits on distinct ordered roots. The  $S_3$  action is two-transitive, so  $Y_f$  is geometrically integral and has arithmetic genus one. The Aubry–Perret singular-curve bound, in the form

$$|\#C(\mathbb{F}_q) - (q+1)| \leq 2\pi_C \sqrt{q}$$

for a reduced geometrically integral projective curve of arithmetic genus  $\pi_C$ , gives

$$\#Y_f(\mathbb{F}_q) \geq q+1 - 2\sqrt{q}$$

[AP95, p. 468]. An integral curve of arithmetic genus one has at most one geometric singular point: if it is singular, its normalization has genus zero and the total delta invariant is one. We therefore delete at most one rational singular point. The diagonal costs

$$Y_f \cdot \Delta = (2, 2) \cdot (1, 1) = 4.$$

Riemann–Hurwitz gives different degree four for the separable cubic map, hence at most four branch values. A singular cubic has at most two distinct roots and at most two ordered off-diagonal pairs, so the branch deletion costs at most eight. If  $q+1 - 2\sqrt{q} > 13$ , a smooth rational point remains outside the diagonal and branch fibers. It gives two distinct rational roots of a squarefree cubic, whose third root is rational. This inequality is equivalent to  $q - 2\sqrt{q} > 12$  and holds for every prime power  $q \geq 23$ .  $\square$



**Proposition 4.7** (Exhaustion and finite bridge). The preceding cases exhaust every redundancy-five pencil for  $q \geq 23$ . Certificate R5 closes exactly

$$q \in \{7, 8, 9, 11, 13, 16, 17, 19\}.$$

It finds sporadic  $S_3$ -stratum orbits exactly at  $q = 7, 8, 9, 11, 13, 17, 19$  and none at  $q = 16$ .

*Proof.* The gcd degree is zero, one, or two. Propositions 4.2 and 4.3 settle the positive-gcd cases. In the trivial-gcd case the cubic map is inseparable or separable. The inseparable case occurs only in characteristic three and the overlap equations give the all-cube nucleus pencil. A separable transitive cubic has geometric monodromy  $C_3$  or  $S_3$ , settled by Lemmas 4.5 and 4.6. The stated finite residue, canonical representatives, stabilizers, histograms, and Frobenius links are exactly the domain of Certificate R5.  $\square$

Table 4: Complete sporadic orbit inventory at redundancy five. A representative  $[a_0, \dots, a_4]$  is in the divided-power syndrome coordinates of Section 3. The last column records the nontrivial Frobenius fusion; a dash means the orbit is fixed.

| $q$ | representative     | orbit size | stabilizer | Frobenius                         |
|-----|--------------------|------------|------------|-----------------------------------|
| 7   | $[0, 1, 0, 0, 3]$  | 28         | 12         | –                                 |
| 7   | $[0, 1, 0, 0, 2]$  | 112        | 3          | –                                 |
| 7   | $[0, 0, 1, 0, 3]$  | 168        | 2          | –                                 |
| 7   | $[0, 1, 0, 3, 2]$  | 168        | 2          | –                                 |
| 7   | $[1, 0, 0, 2, 1]$  | 168        | 2          | –                                 |
| 8   | $[0, 1, 1, 0, 2]$  | 252        | 2          | $\rightarrow [0, 1, 1, 0, 4]$     |
| 8   | $[0, 1, 1, 0, 4]$  | 252        | 2          | $\rightarrow [0, 1, 1, 0, 6]$     |
| 8   | $[0, 1, 1, 0, 6]$  | 252        | 2          | $\rightarrow [0, 1, 1, 0, 2]$     |
| 9   | $[1, 0, 0, 1, 2]$  | 180        | 4          | –                                 |
| 9   | $[0, 1, 0, 4, 1]$  | 360        | 2          | $\leftrightarrow [0, 1, 0, 4, 4]$ |
| 9   | $[0, 1, 0, 4, 4]$  | 360        | 2          | $\leftrightarrow [0, 1, 0, 4, 1]$ |
| 11  | $[1, 0, 0, 1, 10]$ | 330        | 4          | –                                 |
| 11  | $[1, 0, 0, 1, 1]$  | 660        | 2          | –                                 |
| 13  | $[0, 1, 0, 0, 4]$  | 182        | 12         | –                                 |
| 13  | $[1, 0, 0, 4, 7]$  | 546        | 4          | –                                 |
| 17  | $[1, 0, 0, 1, 5]$  | 1224       | 4          | –                                 |
| 19  | $[0, 1, 0, 0, 4]$  | 570        | 12         | –                                 |

The certificate additionally records the complete cubic-pencil member histogram of each representative. Repeated orbit sizes in Table 4 are therefore visibly distinguished without requiring a working-directory file.

Table 5: Independent R5 completeness totals. The first equality is the direct syndrome enumeration versus the orbit–stabilizer sum; the second decomposes the same total into the formula in Theorem 1.1 and the sporadic rows above.

| $q$ | direct = orbit sum | nonsporadic | sporadic |
|-----|--------------------|-------------|----------|
| 7   | $889 = 889$        | 245         | 644      |
| 8   | $1116 = 1116$      | 360         | 756      |
| 9   | $1391 = 1391$      | 491         | 900      |
| 11  | $1848 = 1848$      | 858         | 990      |
| 13  | $2080 = 2080$      | 1352        | 728      |
| 16  | $2432 = 2432$      | 2432        | 0        |
| 17  | $4131 = 4131$      | 2907        | 1224     |
| 19  | $4541 = 4541$      | 3971        | 570      |

**Corollary 4.8** (The balanced quantum extension at  $q = 8$ ). The 1116 projective deep-hole directions of  $\text{PRS}_{\mathbb{F}_8}(4)$  give 1116 relative projective one-column  $[10, 5, 6]_8$  MDS extensions. For each extension  $D$ ,

$$|\Psi_D\rangle = 8^{-5/2} \sum_{c \in D} |c\rangle$$

is a minimum-support  $\text{AME}(10, 8)$  stabilizer state and determines an associated  $[[9, 1, 5]]_8$  quantum MDS code. The directions split as the 360 nonsporadic and 756 sporadic entries of Table 5.

Any product-unitary equivalence between two such equal-phase states is local Clifford, and every transversal conversion between the associated quantum encoders is Clifford factor by factor. In particular, none of the associated quantum MDS codes admits a transversal non-Clifford logical unitary.

*Proof.* A projective deep-hole direction gives a one-column MDS extension. At  $q = 8$ , redundancy five turns the  $[9, 4, 6]_8$  projective Reed–Solomon code into a  $[10, 5, 6]_8$  MDS code, and Table 5 gives the count and its decomposition. The linear MDS–AME construction and the one-party Choi correspondence give the state and the  $[[9, 1, 5]]_8$  code [RGRA18]. The local-unitary and transversal claims are the MDS–CSS rigidity and conversion theorems of the companion paper [Rud26b].  $\square$

*Remark 4.9.* The balance condition for a one-column redundancy- $r$  extension is  $q + 2 = 2r$ . Among the exact fixed-level classifications at redundancies five through seven, it selects only  $(r, q) = (5, 8)$ : the formal alternatives  $q = 10, 12$  are not prime powers. The projective-semilinear orbit table classifies extensions relative to the distinguished punctured Reed–Solomon code; it is not asserted to be a complete table of local-Clifford or local-unitary state orbits.

## 5 Coherent polar induction

### 5.1 Spaces, markers, and bad schemes

Fix a field  $k$ , a two-dimensional  $k$ -space  $E$ , and

$$S_n = \mathbb{P}(\Gamma^n E).$$

For  $\lambda \in E^\vee$ , define divided-power contraction by

$$\langle \iota_\lambda f, g \rangle = \langle f, \lambda g \rangle.$$

Then

$$g \in W_{\iota_\lambda f} \iff \lambda g \in W_f. \quad (1)$$

The lower witness lifts squarefreely only when  $\lambda \nmid g$ . Thus the factor selected for contraction must remain as a marked forbidden root.

**Definition 5.1** (Persistent and modular loci). At redundancy  $j$ , put  $n = j - 1$ . The *persistent scheme*  $\mathcal{P}_j \subset S_n$  is the determinantal scheme

$$\mathcal{P}_j = \{[f] : \text{rank } C_f^{(2)} \leq 2\},$$

cut out by the maximal minors of the three-row catalecticant  $C_f^{(2)} : \text{Sym}^{n-2} E^\vee \rightarrow \Gamma^2 E$ . For every positive-characteristic normal-rational-curve nucleus  $\mathcal{N} \subset \mathbb{P}(\Gamma^{n-1} E)$ , that is, an intersection of the relevant osculating subspaces, let  $\tilde{\mathcal{N}}$  be its affine cone. Its coherent lift is the linear scheme

$$\mathcal{M}_n(\mathcal{N}) = \mathbb{P} \ker \left( \Gamma^n E \longrightarrow \text{Hom}(E^\vee, \Gamma^{n-1} E / \tilde{\mathcal{N}}) \right).$$

The *modular locus*  $\mathcal{M}_j \subset S_n$  is the reduced union of these coherent lifts; Lucas' binomial criterion determines the nucleus coordinate spans at each level. A *collision divisor* records marker parameters at which the marked factor becomes a repeated root.

**Definition 5.2** (Contraction and marker spaces). Let  $C_f : E^\vee \rightarrow \Gamma^{n-1} E$  be  $\lambda \mapsto \iota_\lambda f$ , and put  $U_f = \mathbb{P}(E^\vee) \setminus \mathbb{P}(\ker C_f)$ . The first-polar rational map is

$$\Phi_f : U_f \longrightarrow S_{n-1}, \quad [\lambda] \longmapsto [\iota_\lambda f].$$

Its image is a point or a line. For  $j \geq 1$ , let

$$\text{Conf}_j(\mathbb{P}(E^\vee)) = \{(\lambda_1, \dots, \lambda_j) : \lambda_i \neq \lambda_\ell \text{ for } i \neq \ell\}.$$

The ordered marker space  $U_{f,j}$  is the open subset on which every successive contraction is nonzero. The universal  $j$ -step polar flag is the map

$$(\lambda_1, \dots, \lambda_j) \longmapsto ([\iota_{\lambda_1} f], \dots, [\iota_{\lambda_j} \cdots \iota_{\lambda_1} f]).$$

In an affine coordinate  $r$  the first map is

$$\Phi_f(r) = (a_1 - ra_0, \dots, a_n - ra_{n-1}), \quad \Phi_f(\infty) = (a_0, \dots, a_{n-1}).$$

**Example 5.3** (The complete R5-to-R6 propagation at  $q = 29$ ). Work over  $\mathbb{F}_{29}$  at redundancy six and take

$$f = (6 : 1 : 0 : 0 : 5 : 1).$$

The Hankel kernel is the net of quartics

$$W_f = \ker \begin{pmatrix} 6 & 1 & 0 & 0 & 5 \\ 1 & 0 & 0 & 5 & 1 \end{pmatrix} = \langle (0, 0, 1, 0, 0), (24, 1, 0, 1, 0), (28, 1, 0, 0, 1) \rangle.$$

The displayed basis has no common factor, so the persistent test does not decide the syndrome. Choose the first marker  $r = 0$ . Then

$$\iota_0 f = (1 : 0 : 0 : 5 : 1).$$

This is now a redundancy-five syndrome, and its full cubic pencil is

$$W_{\iota_0 f} = \langle (0, 1, 0, 0), (25, 0, 1, 24) \rangle.$$

The member with pencil parameter  $[9 : 28]$  is

$$h(t) = 4 + 9t - t^2 + 5t^3 \in W_{\iota_0 f},$$

because the two Hankel products are  $4 + 5 \cdot 5 = 0$  and  $5(-1) + 5 = 0$  in  $\mathbb{F}_{29}$ . Direct factorization gives

$$h(t) = 5(t - 2)(t - 6)(t - 27).$$

None of these roots is the marked root 0. Equation (1) therefore lifts  $h$  to

$$t h(t) = (0, 4, 9, 28, 5) \in W_f,$$

a squarefree quartic with roots 0, 2, 6, 27. The Hankel criterion now places  $f$  in the span of the corresponding four normal-rational-curve columns, so  $f$  is shallow. This one calculation exhibits every interface in the R5-to-R6 propagation: the upper Hankel net, the ordinary common-factor test, contraction to the complete R5 cubic pencil, selection on its splitting cover, marker avoidance, and squarefree lifting. In the general proof the R5 curve argument supplies such a pencil parameter after deleting the marked fibers; the displayed parameter  $[9 : 28]$  is one complete instance of that step.

**Definition 5.4** (One-step lower package). Fix  $f \in S_n(\mathbb{F}_q)$  with injective contraction map, so that  $\Phi_f : \mathbb{P}(E^\vee) \rightarrow S_{n-1}$  is defined everywhere. A one-step lower package abstracts the R6 argument: a lower bad locus, the R5 ordered-root curve with its deletions, and the finite set of markers where that construction is unavailable. It consists of:

- (i) a closed bad scheme  $B_{n-1} \subset S_{n-1}$  and a finite stratification  $S_{n-1} \setminus B_{n-1} = \coprod_\alpha X_\alpha$ ;
- (ii) for every  $\lambda \in \mathbb{P}(E^\vee)(\mathbb{F}_q)$  with  $b = \iota_\lambda f \in X_\alpha(\mathbb{F}_q)$ , a specified proper geometrically integral curve  $Y_{\alpha,b}$ , defined over  $\mathbb{F}_q$ , with genus bound  $g_\alpha$  and point bound

$$\#Y_{\alpha,b}(\mathbb{F}_q) \geq q + \kappa_\alpha - 2g_\alpha\sqrt{q};$$

together with a zero-dimensional closed deletion scheme  $D_{\alpha,b,\lambda} \subset Y_{\alpha,b}$ , of degree at most  $\delta_\alpha$ , containing the singular, branch, diagonal, marker, and collision points, such that every rational point outside  $D_{\alpha,b,\lambda}$  determines an ordered rational split squarefree  $g \in W_b$  with  $\lambda \nmid g$ ;

- (iii) a finite parameter scheme  $Z_f \subset \mathbb{P}(E^\vee)$ , of degree at most  $d$ , containing  $\Phi_f^{-1}(B_{n-1})$  and every parameter at which the preceding construction is unavailable, including parameters where a fixed factor of the lower system equals a retained marker.

The index  $\alpha$  includes any constant-field or Frobenius twist; the assertion in (iii) is the required descent statement. All degrees are geometric degrees on the displayed curve or parameter line.

The construction is summarized in Figure 1. The upper branch is the level-specific component calculation; the lower branch is the general transverse argument.




Figure 1: Coherent polar induction. The contained branch is a level-specific component theorem; only the transverse branch is controlled by the general point-count argument.

**Theorem 5.5** (Polar-flag construction). The maps of Definition 5.2 commute with field extension and the natural  $\mathrm{GL}(E)$  action. They include the infinity chart displayed above. If  $g \in W_{\iota_{\lambda_j} \cdots \iota_{\lambda_1} f}$ , then  $\lambda_1 \cdots \lambda_j g \in W_f$ ; the lift is squarefree exactly when  $g$  is squarefree and avoids every marker  $\lambda_i$ .

*Proof.* Contraction is defined by the perfect divided-power/symmetric-power pairing, so it commutes with base change and change of basis. Repeated application of (1) gives the membership assertion. Since the  $\lambda_i$  are distinct on the configuration space, their product is squarefree; the product with  $g$  is squarefree precisely when  $g$  is squarefree and none of the  $\lambda_i$  divides it. The coordinate formula at infinity follows by evaluating  $C_f$  on the second basis covector.  $\square$

**Lemma 5.6** (Three readings of the collision divisor). Assume  $W_f$  has trivial gcd, and let  $\lambda \in \mathbb{P}(E^\vee)$ . The following are equivalent:

- (i) every member of  $W_f$  divisible by  $\lambda$  is divisible by  $\lambda^2$ ;
- (ii)  $\lambda$  is a fixed factor of  $W_{\iota_\lambda f}$ ;
- (iii)  $\lambda$  lies on the collision divisor of the linear series  $W_f$ .

*Proof.* Equation (1) gives the equality of subspaces

$$\lambda W_{\iota_\lambda f} = W_f \cap \lambda \mathrm{Sym}^{n-2} E^\vee.$$

Thus (i) and (ii) are the same statement after dividing by  $\lambda$ . Since  $W_f$  is base-point-free, (i) says exactly that the morphism defined by  $W_f$  has zero differential at  $\lambda$ , which is (iii).  $\square$

**Lemma 5.7** (Uniform collision alternatives). Let  $j \geq 5$ , and let a base-point-free  $g_{j-2}^{j-4}$  on  $\mathbb{P}^1$  define an inseparable morphism in characteristic  $p$ . Then

$$p \mid (j-2), \quad p(j-4) \leq j-2.$$

The only possible inseparable cases are  $(j, p) = (5, 3)$  and  $(6, 2)$ . In the separable case the collision divisor has degree at most  $2j-6$ ; in particular this holds unconditionally for  $j \geq 7$ .

*Proof.* An inseparable morphism factors through absolute Frobenius. Hence  $p \mid (j-2)$ , and its  $j-3$  sections lie in the  $(j-2)/p + 1$ -dimensional space of  $p$ -th powers. This gives the inequality. Since  $(j-2)/(j-4) < 2$  for  $j \geq 7$ , no prime remains; the two smaller cases follow directly. In the separable case some value-derivative minor is nonzero; as a binary Wronskian it has degree  $2(j-2) - 2 = 2j-6$  and contains the collision scheme.  $\square$

**Lemma 5.8** (Linear modular pullbacks). Let  $\mathcal{N} \subset S_{j-2}$  be a projective linear modular nucleus. Then  $\Phi_f^{-1}(\mathcal{N})$  is empty, one point, or the whole parameter line. In the transverse case its degree is at most one.

*Proof.* The polar map has linear coordinates in the parameter, so the pullback ideal is generated by linear forms on  $\mathbb{P}^1$ .  $\square$

**Lemma 5.9** (Old-marker fixed-factor divisor). Assume  $W_f$  has trivial gcd, and fix a retained marker  $x \in \mathbb{P}(E^\vee)$ . The parameters  $\lambda$  for which  $x$  divides every member of  $W_{\iota_\lambda f}$  form a proper closed subscheme of  $\mathbb{P}(E^\vee)$  of degree at most three.

*Proof.* Append the evaluation row at  $x$  to the two consecutive Hankel rows of  $\iota_\lambda f$ . The maximal minors cut out the stated fixed-factor condition. Their entries are linear in  $\lambda$ , so their common zero scheme has degree at most three.

The minors cannot vanish identically. By Equation (1), identical vanishing would say

$$W_f \cap \lambda \operatorname{Sym}^{n-2} E^\vee \subset x \operatorname{Sym}^{n-2} E^\vee \quad \text{for every geometric } \lambda.$$

Given a nonzero  $g \in W_f$ , choose a geometric linear factor  $\lambda \mid g$ . The displayed containment then gives  $x \mid g$ . Thus  $x$  would divide every member of  $W_f$ , contrary to the trivial-gcd hypothesis.  $\square$

**Lemma 5.10** (Exact-linear-gcd transport). Suppose  $W_f = LV$  has exact linear gcd  $L$ , and let  $\lambda \neq L$ . Then every member of  $W_{\iota_\lambda f}$  is divisible by  $L$ . Along a coherent flag chosen outside the rank-at-most-two terminal carrier, the gcd remains exactly  $L$ ; thus the flag reaches the R5 quadratic-graph package with the same fixed factor.

*Proof.* The lifting identity gives

$$\lambda W_{\iota_\lambda f} = W_f \cap \lambda \operatorname{Sym}^{n-2} E^\vee \subset L \operatorname{Sym}^{n-2} E^\vee.$$

Since  $L$  and  $\lambda$  are coprime, division by  $\lambda$  shows that  $L$  divides every lower member. If the lower gcd grows to degree two, the lower syndrome lies in the rank-at-most-two terminal carrier and that parameter is excluded. The dimension bound rules out a larger common factor away from the rank-one boundary. Induction over the marker list proves the assertion. Excluding  $\lambda = L$  is exactly the fixed-factor collision needed to keep the final lifted product squarefree.  $\square$

**Lemma 5.11** (Identically colliding stages). If the collision divisor of a base-point-free stage is the whole parameter line, then the stage is inseparable. The only possibilities are the characteristic-three R5 component and the characteristic-two R6 component, and their coherent lifts lie in the declared persistent or modular locus.

*Proof.* Lemma 5.6 identifies whole-line collision with zero differential of the moving linear series. The numerical argument in Lemma 5.7 leaves only  $(j, p) = (5, 3)$  and  $(6, 2)$ . Proposition 4.7 identifies the first with the wild rank/fixed-factor boundary, while Propositions 6.5 and 6.6 identify the second with the binary modular component. Proposition 5.15 and Lemma 5.8 give their coherent lifts.  $\square$

**Definition 5.12** (Contained pullback and recursive bad scheme). Let  $B \subset S_{n-1}$  be a closed scheme with homogeneous ideal  $I_B$ . On the open  $S_n^\circ \subset S_n$  where  $C_f$  is injective, pull each element of  $I_B$  back along the universal polar map

$$\mathbb{P}(E^\vee) \times S_n^\circ \longrightarrow S_{n-1}.$$

Expand the resulting bihomogeneous forms in the marker coordinates. The vanishing of all their coefficients defines the *contained-pullback scheme*  $\operatorname{Cont}_n(B)$ : its geometric points are exactly the  $f$  for which the whole polar line lies scheme-theoretically in  $B$ . This coefficient ideal is independent of the chosen generators of  $I_B$ .

Let  $\mathcal{B}_5^{\operatorname{red}} \subset S_4$  be the reduced terminal carrier consisting of the rank-at-most-two Hankel locus, the closure of the cyclic carrier on the rank-two open, its true characteristic-two cyclic plane and characteristic-three wild cone, the rank-one contraction boundary, and the two identically colliding inseparable components. The exact-linear-gcd stratum is not part of this carrier: Proposition 4.3 supplies its separate quadratic-graph lower package. Ordinary collision points are deletion divisors, not components of  $\mathcal{B}_5^{\operatorname{red}}$ .

Inductively, let  $\mathcal{B}_{j+1}^{\text{red}} \subset S_j$  be the reduced union of the closure of  $\text{Cont}_j(\mathcal{B}_j^{\text{red}})$ , the noninjective-contraction locus, and the locus on which the universal marker-collision divisor is the whole parameter line. We call this the *reduced recursively pointed contained bad locus* at redundancy  $j + 1$ .

**Definition 5.13** (One-step contained-component assertion). For a specified bad scheme  $B_{n-1}$ , collision divisor, and explicit closed list  $\mathcal{C}_n \subset S_n$ , the assertion  $\text{CC}(n, 1; \mathcal{C}_n)$  says that every  $f$  for which the contraction map is noninjective and hence supplies no polar line,  $\Phi_f^* B_{n-1}$  is the whole parameter line and hence supplies no transverse marker, or the collision divisor is the whole line and hence every lift repeats its marked root, belongs to  $\mathcal{C}_n$ . When the list is clear we write  $\text{CC}(n, 1)$ .

**Theorem 5.14** (One-step polar escape). Let  $f \in S_n(\mathbb{F}_q)$  have a one-step lower package as in Definition 5.4. Put

$$\mathcal{H}_\kappa(g, \delta) = 1 + \left\lfloor \left( g + \sqrt{g^2 + \delta - \kappa} \right)^2 \right\rfloor$$

and assume  $\delta_\alpha \geq \kappa_\alpha - g_\alpha^2$  for every  $\alpha$ . If

$$q \geq \max_\alpha \mathcal{H}_{\kappa_\alpha}(g_\alpha, \delta_\alpha) \quad \text{and} \quad q + 1 > d,$$

then  $W_f$  contains a split squarefree member.

*Proof.* Since  $\#\mathbb{P}^1(\mathbb{F}_q) = q + 1 > d$ , choose  $\lambda \notin Z_f(\mathbb{F}_q)$ . On the corresponding lower stratum the recorded point bound gives

$$q + \kappa_\alpha - 2g_\alpha\sqrt{q} > \delta_\alpha,$$

so there is a rational point outside all deletions. It yields a split squarefree lower member avoiding  $\lambda$ . Equation (1) lifts it to a split squarefree member upstairs, so the original system is shallow.  $\square$

The theorem isolates the numerical escape and one-step lifting argument. Iterating it requires the contained-component assertion at every intermediate level, not only at the top. This paper applies the argument only in the direct redundancy-six and redundancy-seven calculations below and makes no arbitrary-level iteration claim.

## 5.2 Contained-component calculations

The ordinary persistent locus admits a uniform contained-line calculation. This does not classify other lower bad schemes, but it closes the persistent part of every level-specific component gate.

**Proposition 5.15** (Contained rank-two polar lines). Let  $n \geq 5$ , let  $f \in \mathbb{P}(\Gamma^n E)$ , and let

$$A = C_f^{(2)} : \text{Sym}^{n-2} E^\vee \longrightarrow \Gamma^2 E$$

be the three-row catalecticant. Assume the contraction map  $E^\vee \rightarrow \Gamma^{n-1} E$  is injective, so the first-polar map is defined on all of  $\mathbb{P}(E^\vee)$ . This is exactly the complement of the rank-one boundary: after a change of coordinates, a nonzero kernel vector is  $\partial_X$ , and  $\partial_X f = 0$  forces  $f$  to be a scalar multiple of  $Y^{[n]}$ . Such a point lies on the normal rational curve and is already excluded from the split-free locus. Then

$$\Phi_f(\mathbb{P}(E^\vee)) \subset \{b : \text{rank } C_b^{(2)} \leq 2\} \iff \text{rank } A \leq 2$$

scheme-theoretically and in every characteristic. If  $\text{rank } A = 3$ , the rank-drop parameters form a subscheme of  $\mathbb{P}(E^\vee)$  of degree at most three.

*Proof.* For a nonzero  $\lambda \in E^\vee$ , multiplication identifies  $\mathrm{Sym}^{n-3} E^\vee$  with the hyperplane

$$H_\lambda = \lambda \mathrm{Sym}^{n-3} E^\vee \subset \mathrm{Sym}^{n-2} E^\vee,$$

and functoriality gives

$$C_{\iota_\lambda f}^{(2)} = A|_{H_\lambda}.$$

The forward implication is immediate. Conversely, suppose  $\mathrm{rank} A = 3$  and put  $K = \ker A$ , so  $\dim K = n - 4$ . If the lower rank is at most two, rank-nullity on the  $(n - 2)$ -dimensional space  $H_\lambda$  gives

$$\dim(K \cap H_\lambda) \geq (n - 2) - 2 = n - 4 = \dim K.$$

Thus  $K \subset H_\lambda$ . Containment for every geometric  $[\lambda]$  would imply

$$K \subset \bigcap_{[\lambda] \in \mathbb{P}(E_k^\vee)} \lambda \mathrm{Sym}^{n-3} E_k^\vee = 0,$$

because a nonzero binary form cannot be divisible by every linear form. This contradicts  $\dim K = n - 4 > 0$ .

The maximal minors of  $C_{\iota_\lambda f}^{(2)}$  are homogeneous cubics in  $\lambda$ . Vanishing at every geometric point is therefore identical vanishing, proving the scheme-theoretic assertion. In the non-contained case at least one cubic is nonzero, and the common rank-drop scheme lies in its degree-three zero divisor.  $\square$

Thus Proposition 5.15 covers the entire non-rank-boundary locus on which the first-polar line is defined; the excluded noninjective locus is the rank-one normal-rational-curve locus just identified. After rank one and rational split secants are removed as shallow, the upper rank-two locus consists of the persistent tangent and conjugate-secant families. On a persistent fiber the nonsplit torus acts by  $z \mapsto z^n$ , giving  $T/T^n$  modulo inversion and Frobenius, while the tangent unipotent cocycle is  $z \mapsto z + nu$ .

Modular loci are equally explicit. If  $\mathcal{N}$  is a nucleus in the lower divided-power module, Definition 5.1 identifies the complete space whose first-polar line lies in  $\mathcal{N}$  with the linear kernel scheme  $\mathcal{M}_n(\mathcal{N})$ . Lucas' theorem computes the nucleus coordinates, and this kernel computes every coherent lift without an ambient syndrome census. At redundancy  $r$ , if  $p > r - 1$ , Lucas' theorem leaves no nucleus coordinate, so  $\mathcal{M}_r$  is empty. This removes the modular term only; the R5 cyclic carrier survives in large characteristic.

## 6 Redundancies six and seven: complete and gated results

### 6.1 Redundancy six

For

$$f = (a_0 : \cdots : a_5) \in \mathbb{P}(\Gamma^5 E), \quad W_f = \ker \begin{pmatrix} a_0 & a_1 & a_2 & a_3 & a_4 \\ a_1 & a_2 & a_3 & a_4 & a_5 \end{pmatrix},$$

the system  $W_f$  is a net of quartics off the quintic normal rational curve. It is split-free exactly when it has no completely split squarefree quartic.

**Proposition 6.1** (Persistent net and orbit law). If  $\gcd(W_f) = Q$  has degree two, then

$$W_f = Q \mathrm{Sym}^2 E^\vee.$$

The split-free persistent locus has size  $q(q + 1)^2/2$ . On the tangent fiber the unipotent coordinate is  $x \mapsto x + 5u$ ; on the irreducible quadratic fiber the norm-one torus acts through  $z \mapsto z^5$ , with inversion and coefficient Frobenius acting on  $T/T^5$ .



*Proof.* For fixed  $Q$ , the syndromes annihilating  $Q \operatorname{Sym}^3 E^\vee$  form a projective line. If  $Q$  is irreducible, all  $q + 1$  points remain; if  $Q = \lambda^2$ , one endpoint has dependent Hankel rows, leaving  $q$  points. Hence

$$q(q+1) + \frac{q(q-1)}{2}(q+1) = \frac{q(q+1)^2}{2}.$$

Normalize a square to  $U^2$ . The nondegenerate fiber is  $[e_4 + xe_5]$ , and direct degree-five substitution gives  $x \mapsto x + 5u$ . For an irreducible  $Q$ , its stabilizer identifies the  $q + 1$ -point fiber with  $T = \{z \in \mathbb{F}_{q^2}^\times : z^{q+1} = 1\}$  and acts by fifth powers; the normalizer inverts  $z$ , while coefficient Frobenius multiplies  $T/T^5$  by the characteristic. This proves the stated orbit law and the characteristic-five tangent splitting.  $\square$

**Lemma 6.2** (Exact linear gcd is shallow). Let  $q \geq 7$ . If  $\gcd(W_f)$  has degree one, then  $W_f$  contains a split squarefree quartic.

*Proof.* Write  $W_f = LV$ , where  $L$  is rational and  $V \subset \operatorname{Sym}^3 E^\vee$  has vector dimension three. It suffices to find in  $V$  a product of three distinct rational factors, none equal to  $L$ . Move the root of  $L$  to infinity and let  $\ell$  span  $V^\perp$ . On the cubic with distinct finite roots  $x, y, z$ , write

$$\ell((T - xU)(T - yU)(T - zU)) = c_0 + c_1(x + y + z) + c_2(xy + xz + yz) + c_3xyz =: P(x, y, z).$$

Suppose that this value is nonzero for every distinct triple. For fixed distinct  $x, y$ , write  $P = A(x, y) + zB(x, y)$ , where

$$B(x, y) = c_1 + c_2(x + y) + c_3xy.$$

If  $B(x, y) \neq 0$ , the unique zero in  $z$  must be  $x$  or  $y$ . Consequently

$$B(x, y)P(x, y, x)P(x, y, y) = 0$$

for every  $x \neq y$ ; it also vanishes when  $B(x, y) = 0$ . This polynomial has degree at most four in each variable. Since  $q - 1 > 4$ , first varying  $y \neq x$  and then  $x$  shows that it is the zero polynomial. The polynomial ring is a domain. Neither  $P(x, y, x)$  nor  $P(x, y, y)$  vanishes identically unless  $\ell = 0$ , as comparison of the coefficients of  $1, x, y, x^2, xy, x^2y$  shows. Hence  $B = 0$ , so  $c_1 = c_2 = c_3 = 0$ .

Thus  $V$  is the hyperplane of cubics vanishing at infinity, and every member of  $V$  is divisible by  $L$ . This would make  $\gcd(W_f)$  divisible by  $L^2$ , contrary to its exact degree. Therefore a required cubic exists, and multiplying it by  $L$  gives a split squarefree member of  $W_f$ .  $\square$

**Proposition 6.3** (First-polar line and secant degree). For  $r \in \mathbb{F}_q$  put

$$b(r) = (a_1 - ra_0, \dots, a_5 - ra_4), \quad b(\infty) = (a_0, \dots, a_4).$$

Then

$$(T - rU)g \in W_f \iff g \in W_{b(r)}.$$

If  $W_f$  has trivial gcd, its polar line meets the lower secant cubic in degree at most three.

*Proof.* The first equivalence follows by expanding the two Hankel equations of  $(T - rU)g$ . Let

$$C_f = \begin{pmatrix} a_0 & a_1 & a_2 & a_3 \\ a_1 & a_2 & a_3 & a_4 \\ a_2 & a_3 & a_4 & a_5 \end{pmatrix}.$$

The lower catalecticant equals

$$C_f \begin{pmatrix} -r & 0 & 0 \\ 1 & -r & 0 \\ 0 & 1 & -r \\ 0 & 0 & 1 \end{pmatrix}.$$

Cauchy–Binet gives the secant determinant

$$D_f(r) = -\Delta_{012}r^3 + \Delta_{013}r^2 - \Delta_{023}r + \Delta_{123}.$$

It vanishes identically exactly when  $\text{rank } C_f \leq 2$ , equivalently when  $W_f$  has quadratic gcd. In the trivial-gcd case it therefore has at most three zeros.  $\square$

**Proposition 6.4** (Complete lower trivial-gcd carrier). Let  $c \in \mathbb{P}(\Gamma^4 E)$  and suppose that the cubic pencil  $W_c$  has trivial gcd. One of the following applies over an algebraic closure, and the last three alternatives exhaust the cases in which the  $S_3$  package in (i) is unavailable:

- (i) the cubic map is separable with geometric monodromy  $S_3$ , and its off-diagonal fiber square is geometrically integral;
- (ii) the characteristic is neither two nor three and the pencil is cyclic, in which case  $c$  lies on the closure of

$$\{\text{diag}(1, 4, 6, 4, 1)^{-1}[Q^2] : [Q] \in \mathbb{P}(\text{Sym}^2 E^\vee)\};$$

- (iii) the characteristic is two and  $c$  lies on  $\Pi_{\text{cyc}} = \mathbb{P}\langle e_1, e_2, e_3 \rangle$ ;
- (iv) the characteristic is three and  $c$  lies on  $\text{Join}(e_2, C_4)$ , the closure of the wild cyclic pencils together with the inseparable nucleus point.

Thus these displayed tame, binary, and ternary loci exhaust the trivial-gcd lower pencils for which the  $S_3$  package is unavailable.

*Proof.* Proposition 4.4 associates to  $W_c$  a degree-three map. If it is separable, its transitive geometric monodromy is  $C_3$  or  $S_3$ . In the  $S_3$  case two-transitivity makes the off-diagonal fiber square geometrically integral, giving (i). In the cyclic case, Lemma 4.5 identifies the two ramification points with a binary quadratic  $Q$ . Substitution in the two Hankel equations gives the divided-power syndrome  $\text{diag}(1, 4, 6, 4, 1)^{-1}[Q^2]$  whenever the displayed diagonal is invertible. This proves (ii) in characteristics other than two and three.

The same coefficient calculation can be made without division. In characteristic two it is

$$g \cdot e_2 = [0 : \delta\gamma : \alpha\delta + \beta\gamma : \alpha\beta : 0], \quad g = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}.$$

Its image is dense in the plane  $\mathbb{P}\langle e_1, e_2, e_3 \rangle$ , so its closure is exactly the binary locus in (iii). In characteristic three every separable cyclic pencil is wild. Lemma 4.5 puts it in the  $\text{PGL}_2$ -orbit of  $e_2 - ae_4$ . Here  $e_2$  is fixed and the orbit closure of  $e_4$  is the quartic normal rational curve  $C_4$ ; hence the complete wild closure is  $\text{Join}(e_2, C_4)$ . The only inseparable trivial-gcd pencil is the all-cube nucleus pencil, namely its vertex  $e_2$ , by Proposition 4.7. This proves (iv) and exhausts the inseparable case. The three alternatives are therefore exhaustive.  $\square$

**Proposition 6.5** (Cyclic and collision degrees). For a trivial-gcd net:

- (i) in characteristic different from two and three, the lower tame cyclic carrier is a binomially rescaled projected quadratic Veronese surface of degree four and contains no polar line;

- (ii) the pointed ramification condition has degree at most six and does not vanish identically;
- (iii) an  $S_3$  lower pencil with the marked root deleted has total deletion degree at most nineteen.

*Proof.* In syndrome coordinates the tame cyclic surface is

$$\{\text{diag}(1, 4, 6, 4, 1)^{-1}[Q^2] : [Q] \in \mathbb{P}(\text{Sym}^2 E^\vee)\}.$$

When 2 and 3 are invertible this is the projected quadratic Veronese surface, of degree four, and contains no line. Characteristic three has a different wild closure, treated in Proposition 6.6. For (ii), Lemma 5.7 with  $j = 6$  gives the degree-six bound except possibly in characteristic two. In that characteristic, inseparability would identify  $W_f$ , after a change of coordinates, with the full pure-square space  $\langle T^4, T^2U^2, U^4 \rangle$ . Its annihilator is  $\langle e_1^\vee, e_3^\vee \rangle$ . For the two consecutive Hankel rows this would require simultaneously

$$a_0 = a_2 = a_4 = 0, \quad a_1 = a_3 = a_5 = 0,$$

contrary to  $f \neq 0$ . Thus the bound holds in every characteristic.

Finally the R5 off-diagonal curve has deletion degree thirteen, including its possible rational singular point. The unique cubic through a specified marker contributes at most six ordered pairs, giving  $(g, \delta, \kappa) = (1, 19, 1)$  and

$$q + 1 - 2\sqrt{q} > 19.$$

Its first prime-power solution is 29. □

**Proposition 6.6** (Modular contained components). In characteristic three the lower wild carrier is the degree-four cone

$$\text{Join}(e_2, C_4),$$

and no polar line of a trivial-gcd net is contained in it. In characteristic two the cyclic carrier is the plane

$$\Pi_{\text{cyc}} = \mathbb{P}\langle e_1, e_2, e_3 \rangle,$$

whose unique contained polar-line component is

$$\mathcal{N}_3 = \mathbb{P}\langle e_2, e_3 \rangle.$$

*Proof.* Every line on the wild cone is a ruling. By equivariance it suffices to compare the two consecutive rows on one ruling; in the frame  $\langle e_2, e_4 \rangle$  the conditions

$$(a_0, \dots, a_4), (a_1, \dots, a_5) \in \langle e_2, e_4 \rangle$$

force the overlapping coordinates successively to vanish. Undoing the frame gives  $f = \mu\nu(t)$ , including  $f = \mu e_5$  at infinity. These are fixed-factor/rank-boundary syndromes, so no trivial-gcd polar line is contained in the wild cone.

In characteristic two, direct substitution gives

$$g \cdot e_2 = [0 : \delta\gamma : \alpha\delta + \beta\gamma : \alpha\beta : 0],$$

whose closure is  $\Pi_{\text{cyc}}$ . The consecutive-row overlap is now literal: the first row lies in the plane exactly when  $a_0 = a_4 = 0$ , and the second exactly when  $a_1 = a_5 = 0$ . Hence a polar line lies in the plane exactly when

$$a_0 = a_1 = a_4 = a_5 = 0,$$

namely when  $f \in \mathbb{P}\langle e_2, e_3 \rangle$ . This proves both existence and uniqueness of the contained binary component. □

**Proposition 6.7** (Binary nucleus arithmetic). The scheme  $\mathcal{N}_3(\mathbb{F}_q)$  is one projective-semilinear orbit of  $q + 1$  points. It is split-free over  $q = 2^m$  exactly when  $m$  is odd.

*Proof.* For the representative  $e_3$ ,

$$W_{e_3} = \langle 1, t, t^4 \rangle.$$

If  $C = 0$ , the member  $A + Bt + Ct^4$  has degree at most one and cannot have four distinct roots. Assume  $C \neq 0$ . The root differences of  $A + Bt + Ct^4$  then lie in the kernel of  $u \mapsto u^4 + (B/C)u$ . Four distinct rational roots exist exactly when all roots of  $u^3 = B/C$  are rational, equivalently when  $3 \mid q - 1$ , which for  $q = 2^m$  means  $m$  is even. The stabilizer of  $e_3$  is the Borel of order  $q(q - 1)$ , so the orbit has  $q + 1$  points; its lower-unipotent transforms and endpoint are precisely  $\mathcal{N}_3(\mathbb{F}_q)$ .  $\square$

**Proposition 6.8** (Exhaustive R6 contained-component classification). For a redundancy-six syndrome, the alternatives in Definition 5.13 are exhausted as follows:

- (i) a noninjective contraction map is the rank-one normal-rational- curve boundary and is shallow;
- (ii) a polar line contained in the lower secant cubic comes from the upper rank-two catalecticant locus; after its shallow rational secants are removed, this is the persistent tangent and conjugate-secant locus;
- (iii) no trivial-gcd polar line is contained in the tame cyclic surface or the characteristic-three wild cone;
- (iv) in characteristic two, containment in the cyclic plane occurs exactly for  $f \in \mathcal{N}_3 = \mathbb{P}\langle e_2, e_3 \rangle$ ;
- (v) on the remaining trivial-gcd locus the collision divisor is not identically zero.

The exact-linear-gcd lower stratum is not a contained component: it is the separate quadratic-graph package of Proposition 4.3. Consequently the complete nonshallow contained list is the persistent locus together with  $\mathcal{N}_3$  in characteristic two. There is no unlisted contained component.

*Proof.* The noninjective and secant-cubic alternatives are Proposition 5.15 and Proposition 6.3. The tame cyclic and collision assertions are Proposition 6.5; the wild and binary specializations are Proposition 6.6. Proposition 6.4 proves that these are the complete trivial-gcd lower failure loci, rather than a selected list of examples. Together they are all the clauses of the terminal carrier in Definition 5.13: contraction indeterminacy, containment in each component of the lower bad scheme, and identical collision. Hence the list is exhaustive.  $\square$

**Proposition 6.9** (R6 one-step exhaustion). Assume  $q \geq 29$ . Then every split-free redundancy-six syndrome lies in the persistent quadratic-gcd locus or, when  $q = 2^m$  with odd  $m$ , in  $\mathcal{N}_3(\mathbb{F}_q)$ .

*Proof. Contained nets.* The complete contained branch is Proposition 6.8. The common-factor trichotomy is exhaustive. A quadratic gcd gives the persistent tangent and conjugate-secant loci, together with the shallow rational secant; an exact linear gcd is shallow by Lemma 6.2. It remains to treat a trivial-gcd net.

*Transverse exclusions.* Its first-polar line is not contained in the lower secant cubic, by Proposition 6.3. The pullback of that cubic has degree at most three. The tame cyclic surface has degree four and contains no polar line. In characteristic three its wild replacement has the same degree and no contained trivial-gcd polar line; in characteristic two the replacement is

a plane, whose only contained component is  $\mathcal{N}_3$ , by Proposition 6.6. Finally, Proposition 6.5 gives a nonzero collision divisor of degree at most six. Outside the contained binary component, the excluded parameter scheme therefore has degree

$$d \leq 3 + 4 + 6 = 13.$$

*Bottom pencils.* For every remaining marker, the lower cubic pencil has exact linear gcd or geometric monodromy  $S_3$ . In the first case the graph argument of Proposition 4.3 is pointed as follows: besides its eight original deletions, remove the at most two graph points whose split quadratic contains the new marker. Since  $q + 1 > 10$ , a split squarefree cubic avoiding that marker remains. Here the fixed gcd  $\lambda_0$  differs from the marker  $r$ . Indeed,  $\lambda_0 = r$  says that every cubic in  $W_{b(r)}$  vanishes at  $r$ , so every lifted quartic has a double root there; by Lemma 5.6, this is exactly a point of the degree-six collision divisor already placed in  $Z_f$ , so no additional deletion budget is needed. In the  $S_3$  case, Lemma 4.6 supplies the geometrically integral off-diagonal curve. Its singular, diagonal, and branch deletions have degree at most thirteen, and deleting the unique cubic through the marker removes at most six ordered pairs. Thus

$$(g, \delta, \kappa) = (1, 19, 1), \quad q + 1 - 2\sqrt{q} > 19$$

for every  $q \geq 29$ . Explicitly,

$$\mathcal{H}_1(1, 19) = 1 + \left\lfloor (1 + \sqrt{19})^2 \right\rfloor = 29, \quad 30 - 2\sqrt{29} > 19.$$

These data form the one-step package of Definition 5.4, with  $d = 13$ , so Theorem 5.14 produces a split squarefree quartic.

*The modular residue.* This is one field-order threshold, not three characteristic-dependent estimates: the first characteristic-three and characteristic-two prime powers above it happen to be 81 and 32, respectively. On  $\mathcal{N}_3$ , Proposition 6.7 gives the asserted odd-degree split-free orbit and proves shallowness in even extension degree.  $\square$

The covering-radius gate is complete. Its exact high-rate endpoint is  $q = 8$ , since  $6 = \lfloor 8/2 \rfloor + 2$ . Thus Seroussi–Roth nonextendability and Dür’s equivalence give  $\rho(\text{PRS}(q-5)) = 5$  for  $q \geq 8$ , while Certificate R6 checks it directly at  $q = 7$  and independently confirms  $q = 8$ .

**Theorem 6.10** (Redundancy six). For every prime power  $q \geq 7$ , the deep syndromes of  $\text{PRS}(q-5)$  are the persistent locus of size  $q(q+1)^2/2$ , together with:

- (i) respectively 18, 11, 4, 2, 1 exceptional  $\text{PGL}_2(q)$  orbits at  $q = 7, 8, 9, 11, 13$ ;
- (ii) one characteristic-two nucleus orbit of size  $q + 1$  when  $q = 2^m$  and odd  $m \geq 5$ .

The lower bound  $m \geq 5$  avoids double counting: the remaining odd case  $m = 3$ , namely  $q = 8$ , is already the  $|\mathcal{O}| = 9$  row of the exceptional table. There are no other exceptional orbits. The complete persistent orbit law is  $T/T^5$  modulo inversion and Frobenius, with a tangent fixed/nonzero split exactly in characteristic five. At  $q = 7, 8, 9, 11, 13$ , the exceptional direction totals are

$$5152, \quad 4713, \quad 1800, \quad 792, \quad 546,$$

and the corresponding total numbers of deep syndrome directions are

$$5376, \quad 5037, \quad 2250, \quad 1584, \quad 1820.$$

*Proof.* Propositions 6.8 and 6.9 give the complete conceptual exhaustion for every prime power  $q \geq 29$ . Certificate R6 exhausts

$$q \in \{7, 8, 9, 11, 13, 16, 17, 19, 23, 25, 27\},$$

finds exceptional projective-orbit counts 18, 11, 4, 2, 1 only at the five displayed fields, and no further orbit at the remaining fields. Together with the separate radius gate, this covers every prime power  $q \geq 7$ .  $\square$

The listed sources are disjoint. The persistent locus has quadratic gcd, every finite exceptional row has trivial gcd, and the separate nucleus term occurs only for binary fields beyond the finite table.

The five small tables are exhaustive computations. For an exceptional net  $W_f$ , let  $I_f(r)$  be the number of irreducible cubic members of the quotient pencil  $W_{b(r)}$ , including the infinity chart, and define its shared-root pair count by

$$E_f = \sum_{r \in \mathbb{P}^1(\mathbb{F}_q)} \binom{I_f(r)}{2}.$$

In odd characteristic let

$$F_f(T, U) = \sum_{i=0}^5 a_i T^{5-i} U^i$$

be the binary quintic attached to  $f = (a_0 : \cdots : a_5)$ , and let  $\tau_5(f)$  be its rational factorization type; for example,  $1^3 + 2$  denotes a triple rational factor and an irreducible quadratic factor. Their semilinear orbit counts are 18, 5, 2, 2, 1;  $E_f$ , together in odd characteristic with  $\tau_5(f)$ , separates the classes up to precisely Frobenius fusion. Table 6 prints the canonical representatives, stabilizers, orbit sizes, and fusion lengths.

Table 6: Exceptional redundancy-six classes. Each row is one projective-semilinear class. The columns give a canonical syndrome representative, the size and stabilizer order of each constituent  $\mathrm{PGL}_2(q)$ -orbit, the number  $\phi$  of such orbits fused by coefficient Frobenius, the shared-root pair count  $E$ , and the binary-quintic factor type  $\tau_5$  in odd characteristic.

| $q$ | representative            | $ \mathcal{O} $ | $ G_f $ | $\phi$ | $E$ | $\tau_5$        |
|-----|---------------------------|-----------------|---------|--------|-----|-----------------|
| 7   | $[0 : 0 : 0 : 1 : 0 : 1]$ | 168             | 2       | 1      | 7   | $1^3 + 2$       |
| 7   | $[0 : 0 : 1 : 0 : 3 : 1]$ | 336             | 1       | 1      | 19  | $1^2 + 1 + 2$   |
| 7   | $[0 : 0 : 1 : 0 : 3 : 2]$ | 336             | 1       | 1      | 15  | $1^2 + 3$       |
| 7   | $[0 : 1 : 0 : 0 : 0 : 1]$ | 168             | 2       | 1      | 10  | $1 + 2 + 2$     |
| 7   | $[0 : 1 : 0 : 0 : 2 : 0]$ | 112             | 3       | 1      | 33  | $1 + 1 + 3$     |
| 7   | $[0 : 1 : 0 : 1 : 1 : 2]$ | 336             | 1       | 1      | 9   | $1 + 4$         |
| 7   | $[0 : 1 : 0 : 1 : 1 : 5]$ | 336             | 1       | 1      | 9   | $1 + 1 + 3$     |
| 7   | $[0 : 1 : 0 : 1 : 3 : 0]$ | 336             | 1       | 1      | 21  | $1 + 1 + 1 + 2$ |
| 7   | $[0 : 1 : 0 : 1 : 3 : 3]$ | 336             | 1       | 1      | 7   | $1 + 1 + 3$     |
| 7   | $[0 : 1 : 0 : 1 : 3 : 6]$ | 336             | 1       | 1      | 14  | $1 + 1 + 3$     |
| 7   | $[0 : 1 : 0 : 3 : 0 : 5]$ | 168             | 2       | 1      | 11  | $1 + 4$         |
| 7   | $[0 : 1 : 0 : 3 : 0 : 6]$ | 168             | 2       | 1      | 20  | $1 + 4$         |
| 7   | $[0 : 1 : 0 : 3 : 2 : 1]$ | 336             | 1       | 1      | 13  | $1 + 1 + 3$     |
| 7   | $[1 : 0 : 0 : 0 : 1 : 2]$ | 336             | 1       | 1      | 7   | $1 + 4$         |
| 7   | $[1 : 0 : 0 : 0 : 1 : 3]$ | 336             | 1       | 1      | 23  | 5               |
| 7   | $[1 : 0 : 0 : 3 : 1 : 3]$ | 336             | 1       | 1      | 13  | $1 + 4$         |
| 7   | $[1 : 0 : 1 : 0 : 5 : 2]$ | 336             | 1       | 1      | 8   | $1 + 4$         |
| 7   | $[1 : 0 : 1 : 0 : 6 : 2]$ | 336             | 1       | 1      | 8   | 5               |
| 8   | $[0 : 0 : 0 : 1 : 0 : 0]$ | 9               | 56      | 1      | 0   | —               |

| $q$ | representative             | $ \mathcal{O} $ | $ G_f $ | $\phi$ | $E$ | $\tau_5$    |
|-----|----------------------------|-----------------|---------|--------|-----|-------------|
| 8   | $[0 : 1 : 1 : 0 : 2 : 1]$  | 504             | 1       | 3      | 20  | –           |
| 8   | $[0 : 1 : 1 : 0 : 2 : 5]$  | 504             | 1       | 3      | 19  | –           |
| 8   | $[1 : 0 : 0 : 1 : 3 : 6]$  | 504             | 1       | 3      | 14  | –           |
| 8   | $[1 : 0 : 1 : 3 : 5 : 2]$  | 168             | 3       | 1      | 54  | –           |
| 9   | $[0 : 1 : 0 : 0 : 0 : 4]$  | 180             | 4       | 2      | 28  | $1 + 4$     |
| 9   | $[0 : 1 : 0 : 1 : 4 : 2]$  | 720             | 1       | 2      | 29  | $1 + 4$     |
| 11  | $[0 : 0 : 0 : 1 : 0 : 0]$  | 132             | 10      | 1      | 60  | $1^3 + 1^2$ |
| 11  | $[0 : 1 : 0 : 2 : 0 : 10]$ | 660             | 2       | 1      | 30  | $1 + 4$     |
| 13  | $[0 : 1 : 0 : 0 : 0 : 2]$  | 546             | 4       | 1      | 60  | $1 + 4$     |

For  $q = 8$ , integers are binary polynomial-basis codes with  $\alpha^3 = \alpha + 1$ ; for  $q = 9$ , they are ternary polynomial-basis codes with  $\alpha^2 = -1$ . Thus  $\sum \phi = (18, 11, 4, 2, 1)$  and  $\sum \phi|\mathcal{O}| = (5152, 4713, 1800, 792, 546)$ , in the field order 7, 8, 9, 11, 13.

## 6.2 Redundancy seven

For

$$f = (a_0 : \dots : a_6) \in \mathbb{P}(\Gamma^6 E), \quad W_f = \ker \begin{pmatrix} a_0 & a_1 & a_2 & a_3 & a_4 & a_5 \\ a_1 & a_2 & a_3 & a_4 & a_5 & a_6 \end{pmatrix},$$

the system  $W_f$  is a four-dimensional web of quintics. For an affine marker  $r$  the contractions are

$$b(r) = (a_1 - ra_0, \dots, a_6 - ra_5).$$

Coefficient collection gives

$$(t - r)g \in W_f \iff g \in W_{b(r)};$$

the lift is squarefree exactly when  $g(r) \neq 0$ .

The contracted net  $W_{b(r)}$  has quadratic gcd exactly when its three-row catalecticant has rank at most two. Unless the first-polar line is contained in that rank-two locus, these parameters form the degree-at-most-three intersection of Proposition 6.3; the contained case is Corollary 6.15.

**Proposition 6.11** (Complete two-marker lower package). Let  $W_h$  be a trivial-gcd quartic net and let  $x$  be a prescribed forbidden root. Assume in characteristic two that  $h \notin \mathcal{N}_3$ . For every prime power  $q \geq 37$ , the net contains a split squarefree quartic avoiding  $x$ .

*Proof. Choosing the second marker.* By equivariance choose coordinates in which  $x$  is finite; all parameter equations below are then homogenized in  $s$ , so the infinity chart is included. Choose the second contraction marker  $s$ . The unavailable values of  $s$  comprise:  $s = x$ , of degree one; at most three intersections with the lower secant cubic, by Proposition 6.3; at most four intersections with the cyclic surface or its characteristic-three wild replacement; and at most six points of the collision divisor, by Propositions 6.5 and 6.6. In characteristic two the cyclic replacement is a plane, so its pullback has degree at most one unless the polar line is contained in it, the excluded case  $h \in \mathcal{N}_3$ , by Proposition 6.6. We must also exclude the parameters for which the lower pencil has fixed factor  $x$ . They are cut out by the 3-minors obtained by appending the fixed evaluation row  $(1, x, x^2, x^3)$  to the two Hankel rows of  $b(s)$ , and hence have degree at most two in  $s$ . These minors are not all zero. If they vanished identically, then

$$W_h \cap \{g : g(s) = 0\} \subseteq \{g : g(x) = 0\}$$

for every geometric  $s$ . Given  $g \in W_h$ , choose a geometric root  $s$  of  $g$ ; then  $g(x) = 0$ . Thus  $x$  would divide every member of  $W_h$ , contradicting the trivial-gcd hypothesis. Hence the complete second-step parameter scheme has degree at most

$$1 + 3 + 4 + 6 + 2 = 16 < q + 1.$$

Choose  $s$  outside it.

*The linear-gcd branch.* The resulting cubic pencil is neither persistent nor cyclic. If it has exact linear gcd  $\lambda_0$ , then  $\lambda_0 \notin \{x, s\}$ : equality with  $x$  is excluded by the fixed-factor minors above, while equality with  $s$  is the pointed collision excluded by Proposition 6.5. The graph proof of Proposition 4.3 has eight original deletions; avoiding the two additional markers costs at most two graph points each. Since  $q + 1 > 12$ , a split squarefree cubic avoiding  $x$  and  $s$  remains.

*The  $S_3$  branch.* Otherwise the bottom cover has geometric monodromy  $S_3$ . Its off-diagonal curve is geometrically integral of arithmetic genus one. The singular, diagonal, and branch deletions cost at most thirteen, and the fibers through  $x$  and  $s$  cost at most six ordered pairs each. Hence

$$(g, \delta, \kappa) = (1, 25, 1), \quad q + 1 - 2\sqrt{q} > 25,$$

whose first prime-power solution is 37. In either case the cubic lifts through  $s$  to a split squarefree quartic in  $W_h$  avoiding  $x$ .  $\square$

**Proposition 6.12** (Exact-gcd-one avoidance). Suppose the lower net is  $\ell_x U$ , where  $U = \ker \lambda \subset \text{Sym}^3 E^\vee$  has dimension three and  $\gcd(U) = 1$ , so that  $\ell_x$  is the exact gcd of the lower net. If  $x = r$ , the parameter belongs to the collision divisor. If  $x \neq r$  and  $q \geq 16$ , then  $U$  contains a split cubic avoiding both  $x$  and  $r$ .

*Proof.* If  $x = r$ , every lifted quintic has the repeated factor  $(t - r)^2$ . Lemma 5.6 identifies this with the degree-eight collision divisor of Proposition 6.13. This is the unavailable branch, so assume  $x \neq r$ . Put  $S = \mathbb{P}^1(\mathbb{F}_q) \setminus \{x, r\}$ . For ordered distinct  $u, v \in S$ , the map  $w \mapsto \lambda(\ell_u \ell_v \ell_w)$  is linear. If it vanishes identically, choose any remaining  $w$ ; otherwise its homogenization is a nonzero linear form on  $\mathbb{P}^1$ , so it has exactly one rational projective zero, including the point at infinity when appropriate. The equations  $w = x, r, u, v$  have respective bidegrees  $(1, 1), (1, 1), (2, 1), (1, 2)$  in  $(u, v)$ . None is identically zero: the first two would give a common factor of  $U$ , contrary to  $\gcd(U) = 1$ , and the last two are excluded because double-root cubics span  $\text{Sym}^3 E^\vee$ . A nonzero bidegree- $(a, b)$  divisor on  $\mathbb{P}^1 \times \mathbb{P}^1$  has at most  $(a + b)(q + 1)$  rational points. Thus the bad pairs number at most  $10(q + 1)$ , whereas  $(q - 1)(q - 2) > 10(q + 1)$  for  $q \geq 16$ .  $\square$

**Proposition 6.13** (Collision divisor). After fixed factors are removed, the collision divisor of the moving  $g_5^3$  has degree at most eight.

*Proof.* This is Lemma 5.7 with  $j = 7$ : the series is separable and  $2j - 6 = 8$ .  $\square$

The characteristic-two nucleus line at redundancy six has exactly one coherent lift, the central point  $e_3$ , with

$$W_{e_3} = \langle 1, t, t^4, t^5 \rangle.$$

It is split-free for odd  $m$ , and hence deep whenever the separate radius-six gate holds. For even  $m$ , the homogenization of  $t^4 + t$  has the five roots  $\mathbb{P}^1(\mathbb{F}_4)$ .

**Proposition 6.14** (Central binary lift). The complete inverse image of the R6 nucleus line under consecutive sextic contraction is  $[e_3]$ . It is split-free over  $\mathbb{F}_{2^m}$  exactly when  $m$  is odd.



*Proof.* Putting both consecutive rows in  $\mathbb{P}\langle e_2, e_3 \rangle$  gives  $a_0 = a_1 = a_2 = a_4 = a_5 = a_6 = 0$ . Thus only  $e_3$  remains, with  $W_{e_3} = \langle 1, t, t^4, t^5 \rangle$ . For odd  $m$ , contraction at a root of a hypothetical split quintic would give a split quartic on the R6 nucleus line, contradicting Proposition 6.7. For even  $m$ ,  $\mathbb{F}_4 \subset \mathbb{F}_q$  and the homogenization of  $t^4 + t$  has its four affine  $\mathbb{F}_4$  roots and the simple root at infinity.  $\square$

**Corollary 6.15** (Exhaustive R7 contained-component classification). The assertion CC(6, 1) is exhaustive in every characteristic. A noninjective contraction is the shallow rank-one boundary. Containment in the lower persistent locus gives upper catalecticant rank at most two; after the shallow rational secants are removed, these are exactly the persistent tangent and conjugate-secant families. The complete lift of the only additional lower component, the characteristic-two nucleus line, is the central point  $e_3$ . The remaining collision divisor is nonzero. Thus the nonshallow contained list is precisely the persistent locus and, in characteristic two,  $e_3$ ; there is no further contained component.

*Proof.* Apply Proposition 5.15 with  $n = 6$ . The three rational types of the quadratic recurrence are split squarefree, repeated rational, and irreducible. They give respectively the shallow rational secant, tangent, and conjugate-secant loci. The remaining lower binary nucleus line has the separate complete lift computed in Proposition 6.14. Proposition 6.13 shows that the collision divisor is nonzero. These are respectively the contraction-indeterminacy, lower-component-containment, and identical-collision alternatives of Definition 5.13, so the list is exhaustive.  $\square$

**Theorem 6.16** (Redundancy-seven classification). Certificate R7 unconditionally classifies

$$q \in \{7, 8, 9, 11, 13, 16, 17, 19, 23, 25, 27, 29, 31, 32\}.$$

For every prime power  $q \geq 13$ , the split-free syndromes at redundancy seven are the persistent locus of size  $q(q+1)^2/2$ , together with the central nucleus point when  $q = 2^m$  and  $m$  is odd. At  $q = 7, 8, 9, 11$  the additional split-free orbit-size profiles, after removing the persistent locus and also the central nucleus singleton at  $q = 8$ , are:

$$\begin{array}{l|l} 7 & 56, 84^5, 112^2, 168^{45}, 336^{141} \\ 8 & 63, 72, 84^3, 168^4, 252^{24}, 504^{86} \\ 9 & 180^3, 240^6, 360^{18}, 720^{27} \\ 11 & 264^2, 440, 660^2. \end{array}$$

This is the complete split-free classification for all prime powers  $q \geq 7$ . Since Seroussi–Roth nonextendability and Dür’s equivalence give  $\rho(\text{PRS}(q-6)) = 6$  for every prime power  $q \geq 11$ , it is also the complete deep-hole classification in that range. At  $q = 7, 8, 9$  the table is not promoted to a code deep-hole classification without a separate covering-radius calculation. The persistent orbit law is  $T/T^6$  modulo inversion and Frobenius, with tangent splitting in characteristics two and three.

*Proof.* For fixed irreducible quadratic  $Q$ , the persistent fiber has  $q+1$  points, while for  $Q = \lambda^2$  it has  $q$  non-rank-one points. There are  $q(q-1)/2$  irreducible quadratics and  $q+1$  rational squares. Hence the persistent count is

$$\frac{q(q-1)}{2}(q+1) + (q+1)q = \frac{q(q+1)^2}{2}.$$

For  $q \geq 37$ , choose a first marker outside the at most three catalecticant intersections, eight collision parameters, and, in characteristic two, one intersection with the nucleus line. If the contracted quartic net has trivial gcd, Proposition 6.11 supplies the complete second one-step

package: it excludes the cyclic/wild carrier and fixed-gcd/marker coincidences before applying either the pointed linear-gcd graph or the  $S_3$  cover. If the contracted net has exact linear gcd, write it as  $\ell_x U$ . Exactness gives  $\gcd(U) = 1$ , so Proposition 6.12 applies; its excluded  $x = r$  case is already among the eight collision parameters. Corollary 6.15 leaves exactly the persistent locus and Proposition 6.14. Certificate R7 covers

$$q \in \{7, 8, 9, 11, 13, 16, 17, 19, 23, 25, 27, 29, 31, 32\}$$

by the exact primary quotient enumeration. A separate replay checks the recorded representatives and their orbits. An independent direct-locus replay imports no R7 generator or quotient partition: below 16 it constructs the literal four-secant complement, from 16 onward it constructs the classified R6 pointed locus directly, including the transient marked orbit at  $q = 19$ , and in every field it intersects all contraction conditions and rebuilds the complete  $\mathrm{PGL}_2(q)$ -orbit and Frobenius partition. It agrees exactly with the public record in all fourteen fields. The radius theorem promotes the split-free list only from  $q \geq 11$ .  $\square$

For the four finite rows Certificate R7 contains the canonical representative, stabilizer, persistent/central flags, and Frobenius link of every orbit. The displayed size profile is only a compact summary; repeated sizes are distinguished by those canonical records. There are respectively 194, 119, 54, 5 exceptional representatives at  $q = 7, 8, 9, 11$ ; the complete lists are printed in `supplement/CLASSIFICATION-RECORDS.md`, while the matching JSON is the machine-readable record. The following gives one printed canonical representative for every orbit-size block in the displayed profile; repeated orbits within a block are distinguished in the full list:

| $q$ | $ \mathcal{O} $ | representative                  |
|-----|-----------------|---------------------------------|
| 7   | 56              | $[0 : 0 : 0 : 0 : 1 : 0 : 0]$   |
|     | 84              | $[0 : 1 : 0 : 0 : 0 : 1 : 0]$   |
|     | 112             | $[0 : 0 : 1 : 0 : 0 : 2 : 0]$   |
|     | 168             | $[0 : 0 : 0 : 0 : 1 : 0 : 1]$   |
|     | 336             | $[0 : 0 : 0 : 1 : 0 : 1 : 1]$   |
| 8   | 63              | $[0 : 0 : 0 : 1 : 0 : 0 : 1]$   |
|     | 72              | $[0 : 0 : 0 : 0 : 1 : 0 : 0]$   |
|     | 84              | $[0 : 1 : 1 : 1 : 1 : 3 : 6]$   |
|     | 168             | $[1 : 0 : 1 : 1 : 3 : 7 : 7]$   |
|     | 252             | $[0 : 0 : 1 : 0 : 1 : 0 : 0]$   |
| 9   | 504             | $[0 : 0 : 1 : 0 : 1 : 0 : 1]$   |
|     | 180             | $[0 : 1 : 0 : 0 : 0 : 4 : 0]$   |
|     | 240             | $[0 : 0 : 1 : 0 : 1 : 1 : 1]$   |
|     | 360             | $[0 : 1 : 0 : 1 : 3 : 3 : 2]$   |
|     | 720             | $[0 : 0 : 1 : 1 : 0 : 2 : 2]$   |
| 11  | 264             | $[0 : 1 : 0 : 0 : 0 : 0 : 2]$   |
|     | 440             | $[1 : 0 : 1 : 1 : 4 : 3 : 3]$   |
|     | 660             | $[1 : 0 : 1 : 0 : 1 : 4 : 8]$ . |

For  $q = 8, 9$ , the coordinates use the same polynomial-basis encoding specified below Table 6.

The  $q = 19$  lower net

$$W = \langle 1, t^3, t^4 \rangle$$

shows why the parameter must remain marked. Its split squarefree members are exactly

$$U(T^3 - uU^3), \quad u \in (\mathbb{F}_{19}^\times)^3 = \{1, 7, 8, 11, 12, 18\}.$$

All six contain the marked point at infinity. The obstruction is pointed rather than intrinsic to  $W$ : at the marker 0, for example,  $U(T^3 - U^3)$  is an admissible member. No coherent

Table 7: Field-range closure ledger. “Certificate” means a complete finite classification with canonical representatives, not merely an orbit count.

| Result | prime powers                                                         | closure mechanism                                                                                                          |
|--------|----------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------|
| R5     | 7, 8, 9, 11, 13, 16, 17, 19<br>$q \geq 23$                           | direct census and independent replay<br>cubic-cover geometry, Lemma 4.6                                                    |
| R6     | 7, 8, 9, 11, 13, 16<br>17, 19, 23, 25, 27<br>$q \geq 29$             | direct syndrome/radius census<br>finite structural bridge certificate<br>marked-cover point bound and contained components |
| R7     | 7, 8, 9, 11<br>13, 16, 17, 19, 23, 25, 27, 29, 31, 32<br>$q \geq 37$ | direct split-free census<br>coherent-polar finite bridge certificate<br>marked-cover point bound and contained components  |

sextic polar line lies in the infinity-pointed orbit. Thus the R7 proof escapes over the marker parameter; classifying unpointed bad lower fibers would retain a false exception.

The R7 rows classify split-free syndromes. Conversion to code deep-hole directions uses the separate covering-radius gate and is made only for  $q \geq 11$ . The public certificate schema and exact replay boundaries are given in the supplement; the table does not promote orbit-size profiles to identifiers.

There is no prime power in any of the open numerical intervals (19, 23), (27, 29), or (32, 37). Thus adjacent finite and geometric rows in Table 7 leave no unlisted field order.

## 7 Verification and formal boundary

No geometric classification theorem in this paper is claimed to be formalized in Lean. The formal layer checks coordinate identities and conditional synthesis implications with the component geometry, rational-point existence, group actions, and certificate semantics kept as explicit inputs.

The conceptual theorems above are accompanied by deterministic evidence bundles. Table 8 records which claims use exhaustive computation. Each public label resolves, through the supplement schema, to its generator, certificate, replay where stated, and SHA-256 manifest. Table 1 gives the corresponding claim-level mathematical dependencies.

Table 8: Public verification map. Exact filenames, commands, hashes, domains, and stop conditions belong to the electronic supplement.

| Public label      | Verified domain                                                    | Replay and trust boundary                                                                                                                                                                                |
|-------------------|--------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Certificate R5    | nineteen audited field records; eight-field finite bridge below 23 | independent pointwise and orbitwise replay; canonical representatives and exhaustion records                                                                                                             |
| Certificate R6    | direct scan through $q = 16$ ; finite bridge below 29              | independent syndrome replay and separate radius gate                                                                                                                                                     |
| Certificate R6-NF | small exceptional normal forms                                     | deterministic normal-form reconstruction; not a second implementation                                                                                                                                    |
| Certificate R7    | $q = 7, 8, 9, 11$ and finite bridge below 37                       | primary quotient enumeration, representative/orbit replay, and an independent direct-locus enumeration of the complete split-free sets and orbit partitions; split-free and radius flags remain separate |

Table 6 is a deterministic projection of the hash-pinned public R6 and R6-NF records. The top-level verifier regenerates it and checks byte-for-byte equality with the included source.

The public labels are defined by `supplement/CERTIFICATE-SCHEMA.md`. The paper-local bundle, literal replay commands, toolchain locks, hashes, byte counts, and working directories are recorded in `supplement/REPRODUCING.md`, `supplement/EVIDENCE-MANIFEST.json`, and `supplement/RELEASE-MANIFEST.md`. The single entry point `python3 supplement/verify.py` checks the bundle manifest, classification records and their expected hashes, and the local PDF and supplement-artifact rows printed in the release manifest; its `--replay` option runs the paper-local Python replays.

The paper-facing Lean gate checks coordinate algebra and conditional synthesis, not the geometric classifications or rational-point arguments. The exact declarations, hypotheses, axiom audit, and reproduction command are listed in `supplement/LEAN-STATEMENTS.md`. Lean checks the contraction algebra, degree-specific R6/R7 budget arithmetic, finite-record arithmetic, and conditional synthesis interfaces used by the R5–R7 results proved here. Companion stable-component and arbitrary-level budget modules remain regression checked but are not evidence for a theorem in this paper. Lean also checks the R5–R7 threshold-to-range arithmetic and the exact  $(q, r) = (8, 6)$  endpoint used at redundancy six. The formal adapter keeps the dual-GRS identification, Seroussi–Roth nonextendability, Dür equivalence, and the concrete radius proposition as separate hypotheses. The balanced quantum corollary has a separate cross-paper gate. Lean checks its field-eight certificate arithmetic, balanced-row uniqueness, and the length-ten LU and transversal-Clifford specializations; the standard MDS–AME and one-party Choi correspondences remain explicit cited inputs.

## Data, code, and disclosure

The electronic supplement accompanying this manuscript contains the source, canonical classification representatives and stabilizers, certificate schemas, generators, independent replays where available, expected hashes, toolchain locks, and a top-level verification command. Section 7 assigns each computational or formal claim its exact trust route.

## 8 Scope and open problems

Proposition 6.8 and Corollary 6.15 give complete contained-component calculations at redundancies six and seven. Extending those calculations to arbitrary redundancy requires an exact scheme-theoretic ledger for the bottom carrier and a proof that all coherent modular lifts are charged with their full union degree. Those steps are not claimed here.

1. Redundancy five is complete for every  $q \geq 7$ .
2. Redundancy six is a complete deep-hole classification over every prime power  $q \geq 7$ . Redundancy seven has a complete split-free classification over the same range; its  $q = 7, 8, 9$  covering-radius gate remains separate.
3. At higher redundancy, the exact recursively contained components, the union degree of all modular pullbacks, and arithmetic membership in the resulting Lucas carriers remain open.

In particular, the paper does not settle the general Reed–Solomon deep-hole conjecture and makes no arbitrary-redundancy classification.

The last sporadic fields at redundancies five, six, and seven are 19, 13, 11, while the corresponding point-count gates are 23, 29, 37. These opposite trends ask whether sufficiently high redundancy has any sporadic fields outside the persistent and modular loci.

## Acknowledgment

This project was developed with extensive assistance from OpenAI Codex and Anthropic Claude. Under the author's direction, the systems assisted throughout with proof exploration, literature research, symbolic and finite computation, code and formal-proof development, verification, and manuscript drafting and revision. The author checked the resulting arguments, computations, code, and cited sources, reviewed and edited all AI-assisted material, and assumes responsibility for all content.

## References

- [AP95] Yves Aubry and Marc Perret. Coverings of singular curves over finite fields. *Manuscripta Mathematica*, 88:467–478, 1995.
- [BL19] Simeon Ball and Michel Lavrauw. Arcs in finite projective spaces. *EMS Surveys in Mathematical Sciences*, 6(1–2):133–172, 2019.
- [CMP17] Eda Cesaratto, Guillermo Matera, and Mariana Pérez. The distribution of factorization patterns on linear families of polynomials over a finite field. *Combinatorica*, 37(5):805–836, 2017.
- [CS11] Gonzalo Comas and Malena Seiguer. On the rank of a binary form. *Foundations of Computational Mathematics*, 11:65–78, 2011.
- [Dür87] Arne Dür. The automorphism groups of Reed–Solomon codes. *Journal of Combinatorial Theory, Series A*, 44(1):69–82, 1987.
- [Dür94] Arne Dür. On the covering radius of Reed–Solomon codes. *Discrete Mathematics*, 126(1–3):99–105, 1994.
- [GH00] Johannes Gmainer and Hans Havlicek. Nuclei of normal rational curves. *Journal of Geometry*, 69:117–130, 2000.
- [IK99] Anthony Iarrobino and Vassil Kanev. *Power Sums, Gorenstein Algebras, and Determinantal Loci*, volume 1721 of *Lecture Notes in Mathematics*. Springer, Berlin, Heidelberg, 1999.
- [Kai17] Krishna Kaipa. Deep holes and MDS extensions of Reed–Solomon codes. *IEEE Transactions on Information Theory*, 63(8):4940–4948, August 2017.
- [KPP25] Krishna Kaipa, Nupur Patanker, and Puspendu Pradhan. On the  $PGL_2(q)$ -orbits of lines of  $PG(3, q)$  and binary quartic forms. <https://arxiv.org/abs/2312.07118>, 2025. Version 3, 9 August 2025.
- [RGRA18] Zahra Raissi, Christian Gogolin, Arnau Riera, and Antonio Acín. Optimal quantum error correcting codes from absolutely maximally entangled states. *Journal of Physics A: Mathematical and Theoretical*, 51(7):075301, 2018.
- [Rud26a] Tavis Rudd. Arcs complete outside a prescribed conic: An exact defect identity and  $\rho_C(16) = 9$ . Companion paper in this bundle, 2026.
- [Rud26b] Tavis Rudd. Local-unitary rigidity and transversal clifford groups for mds–css ame states. Companion paper in this bundle, 2026.
- [Rud26c] Tavis Rudd. Quadratic recovery and cubic orientation in conic matching quotients. Companion paper in this bundle, 2026.
- [Rud26d] Tavis Rudd. Reconstructing the Clebsch code from its deep-hole syndrome locus. Companion paper in this bundle, 2026.
- [SR86] Gadiel Seroussi and Ron M. Roth. On MDS extensions of generalized Reed–Solomon codes. *IEEE Transactions on Information Theory*, 32(3):349–354, 1986.
- [Wan26] Tianhao Wang. Splitting of polynomial families via galois theory. <https://arxiv.org/abs/2606.12810>, 2026. Version 1, 11 June 2026.
- [WDC23] Yansheng Wu, Cunsheng Ding, and Tingfang Chen. Extended codes and deep holes of MDS codes. <https://arxiv.org/abs/2312.05534>, 2023. Version 1, 9 December 2023.
- [XHX17] Xiaofan Xu, Shaofang Hong, and Yongchao Xu. On deep holes of generalized projective Reed–Solomon codes. <https://arxiv.org/abs/1705.07823>, 2017. Version 1, 22 May 2017.
- [Xu23] Xiaofan Xu. On deep holes of projective Reed–Solomon codes over finite fields with even characteristic. *Wuhan University Journal of Natural Sciences*, 28(1):15–19, 2023.
- [ZW17] Jun Zhang and Daqing Wan. Explicit deep holes of Reed–Solomon codes. <https://arxiv.org/abs/1711.02292>, 2017. Version 1, 7 November 2017.
- [ZWK20] Jun Zhang, Daqing Wan, and Krishna Kaipa. Deep holes of projective Reed–Solomon codes. *IEEE Transactions on Information Theory*, 66(4):2392–2401, 2020.