

A binary $[78, 36, 12]$ code from the passant lines of a conic over $\mathbb{F}_{13}$

Tavis Rudd

August 2026

Abstract

Let $\mathcal{C}$ be a nonsingular conic in $\text{PG}(2, 13)$, and form the binary incidence code on its 78 internal points using the 78 passant lines. We prove that this code has parameters $[78, 36, 12]_2$. It has exactly 364 minimum words, in four $\text{PGL}(2, 13)$ -orbits of size 91, and each orbit spans the code. The minimum supports reconstruct the passant incidence matrix and the six-class elliptic association scheme; their coordinate-permutation automorphism group is exactly $\text{PGL}(2, 13)$. The proof combines a tangent-product exclusion in weight eight, two exhaustive but compact syndrome-disjointness checks in weight ten, the binary association algebra for spanning, and concurrence with four resolving anchors for reconstruction and symmetry.

Keywords. binary incidence code; finite conic; passant line; minimum word; association scheme; reconstruction.

1 Introduction and statement

Incidence codes often retain less geometry than the incidence structures that define them. Row operations erase the choice of parity checks, and a minimum-distance computation need not reveal why the first codewords occur. Here the minimum words recover both the parity checks and the association scheme on the coordinates.

Fix a nonsingular conic $\mathcal{C} \subset \text{PG}(2, 13)$. A line is *passant* when it is disjoint from $\mathcal{C}(\mathbb{F}_{13})$. Conic polarity identifies the 78 internal points with the 78 passant lines. Let M be their binary incidence matrix and put

$$K = \ker_{\mathbb{F}_2} M \subseteq \mathbb{F}_2^{78}.$$

The coordinate set is the set of internal points. Automorphisms below are coordinate permutations preserving the indicated code or minimum-support system.

Let

$$\mathcal{H} = \{\text{supp}(c) : c \in K, \text{wt}(c) = 12\}$$

be the minimum-support hypergraph on the unlabeled coordinate set of K .

Theorem 1.1 (Minimum-layer reconstruction). *The code K has parameters $[78, 36, 12]_2$ and exactly 364 minimum words. Under $\text{PGL}(2, 13)$, those words form four orbits of size 91: one has stabilizer isomorphic to S_4 , and three have dihedral stabilizer of order 24. Every one of the four orbits spans K .*

From the abstract hypergraph $\mathcal{H}$, pair and triple concurrence reconstruct the six relations of the elliptic association scheme on the coordinate set. Among the passant seven-cliques,

the 78 with zero triple concurrence are exactly the supports of the rows of M . Consequently $\mathcal{H}$ reconstructs M up to independent coordinate and row permutation, and

$$\text{Aut}(K) = \text{Aut}(\mathcal{H}) = \text{PGL}(2, 13)$$

in its symmetric-square action on the 78 internal points.

Reconstruction here is a permutation statement. It does not choose an equation for the conic, label the projective points, or order the parity checks. The proof has three finite leaves: a five-row clique closure in weight eight, two syndrome-disjointness checks in weight ten, and an exact orbit/concurrence replay for the minimum layer. The reductions to those leaves and the association-algebra argument are mathematical proofs in the text.

The code belongs to the fixed-conic LDPC family introduced by Droms–Mellinger–Meyer [2]. Their general bounds, as recorded in the later comparison table of Ma–Liu–Tian [5, Table 1], specialize here to $8 \leq d \leq 12$. Madison and Wu proved the general nullity formula $(q-1)^2/4$ [4, Corollary 6.3]. Hollmann and Xiang constructed the elliptic association scheme afforded by the same conic action [3, Sections 3–5]. We determine the upper endpoint of the distance interval at $q = 13$, classify its minimum words, and recover the incidence and relation structures from their unlabeled support hypergraph.

2 Distance twelve

We use the model $\mathcal{C} : XZ - Y^2 = 0$. Every row and column of M has weight seven, and conic polarity identifies the two index sets. The Madison–Wu formula gives $\dim K = 36$; direct binary elimination is an independent check of rank 42.

Summing all row equations gives the all-one functional, so every word of K has even weight. If a nonzero word contains an internal point P , each of the seven passants through P contains a second support point. Those seven points are distinct. Thus every nonzero word has weight at least eight.

2.1 The tangent obstruction in weight eight

Suppose that a word has weight eight and fix a support point P . Each of the seven passants through P must contain another support point. Since only seven points remain, each passant contains exactly one of them. Applying the same argument at every support point shows that every support join is passant and that no three support points are collinear. Thus the support is an eight-arc, and every passant pencil at a support point is saturated. The seven remaining lines through each support point are therefore both the combinatorial tangents to the arc and the conic secants through that point.

For each secant line choose a nonzero linear equation. For an internal point P , put

$$T_P(X) = \prod_{\substack{\ell \ni P \\ \ell \text{ secant to } \mathcal{C}}} \ell(X).$$

For a pairwise-passant triple define

$$h(P, Q, R) = \frac{T_P(Q)T_Q(R)T_R(P)}{T_P(R)T_R(Q)T_Q(P)}.$$

Every displayed evaluation is nonzero because the three joins are passant. Rescaling a line equation multiplies one numerator and one denominator factor by the same scalar, so h is well defined. The saturated-pencil identification above makes the factors of T_P precisely

the combinatorial tangents at P . Segre's lemma of tangents in the coordinate-free form of Ball–Lavrauw [1, Lemma 27] gives $h(P, Q, R) = 1$ for every triple in the hypothetical support. Fix $P = (1 : 0 : 2)$. Its other seven points would form a clique in the graph on the 42 passant-join neighbors of P , where Q and R are adjacent when QR is passant and $h(P, Q, R) = 1$.

An order-14 element of the stabilizer of P splits the vertices into three cyclic orbits A_i, B_i, C_i , indexed by $\mathbf{Z}/14$. Direct substitution in the product above gives

orbit pair	$j - i \pmod{14}$
AA	4, 6, 8, 10
AB	6, 7, 11, 12
AC	1, 3
BB	6, 8
BC	3, 5, 6, 8, 9, 11
CC	2, 4, 10, 12.

Up to simultaneous translation, every four-clique is represented by one of the following five rows; the right column gives its unique extension:

four-clique	unique extension
A_0, B_6, B_{12}, C_1	C_3
A_0, B_6, B_{12}, C_3	C_1
A_0, B_6, C_1, C_3	B_{12}
A_0, B_{12}, C_1, C_3	B_6
B_0, B_6, C_9, C_{11}	A_8 .

They close to fourteen translates of one five-clique, and none extends. The local clique number is five, so a seven-clique cannot occur. This excludes weight eight.

2.2 The two weight-ten profiles

Fix one point in a hypothetical weight-ten support. Let s of the other nine points be joined to it by secants. The remaining $9 - s$ points occur in positive odd numbers in its seven passant fibres. Hence s is even and $9 - s \geq 7$, so $s = 0$ or $s = 2$. The profiles are

$$(3, 1, 1, 1, 1, 1, 1; 0), \quad (1, 1, 1, 1, 1, 1, 1; 2),$$

where the last entry is the number of secant neighbors.

Write m_Q for the 78-bit incidence column at Q . For the first profile and each choice of exceptional fibre, the finite check forms two canonically sorted partial-XOR sets of sizes 216 and 4320. For the second profile the sizes are 216 and 771024. In every case the two sets are disjoint. The raw domains have sizes

$$7 \binom{6}{3} 6^6 = 6,531,840, \quad 6^7 \binom{35}{2} = 166,561,920.$$

Thus the split covers every support in each profile and proves that no candidate has zero syndrome. An independent dynamic program reconstructs the conic and incidence columns, builds the full-fibre XOR sets with a different partition, and obtains the same two empty intersections.

Finally, the twelve internal points

$$(1 : 0 : 2), (1 : 3 : 2), (1 : 4 : 5), (1 : 1 : 8), (1 : 4 : 8), (1 : 1 : 7), \\ (1 : 7 : 12), (1 : 3 : 3), (1 : 9 : 11), (1 : 10 : 11), (1 : 0 : 5), (1 : 8 : 7)$$

have zero syndrome. Their stabilizer is dihedral of order 24. Therefore $d(K) = 12$.

3 Minimum words and the elliptic scheme

Fix one support point. Pencil parity leaves $s = 0, 2, 4$ secant neighbors. The corresponding multiplicities in the seven passant fibres, followed by the number of secant neighbors, are

$$(5, 1^6; 0), \quad (3, 3, 1^5; 0), \quad (3, 1^6; 2), \quad (1^7; 4).$$

A syndrome-indexed meet in the middle exhausts these four finite domains and finds exactly 56 weight-twelve words through the fixed point. The canonical replay records four projective representatives. Their disjoint fixed-point slices each have size 14, and together they equal the 56-word exhaustive list. The order-28 stabilizer of the fixed point acts transitively on each slice; the stabilizer of a support inside it has order 2. Transitivity, or equivalently the count $56 \cdot 78/12 = 364$, proves global exhaustion.

The replay generates each orbit from its recorded representative and checks every support against M . The stabilizers in $\text{PGL}(2, 13)$ have order 24, so each orbit has size $2184/24 = 91$. Their element-order profiles identify one stabilizer with S_4 and the other three with dihedral groups of order 24.

For a pair or triple of coordinates, its concurrence is the number of the 364 minimum supports containing it. Pair concurrences are 7, 9, 12 when the joining line is passant and 6, 8 when it is secant. The full triple- concurrence histograms are distinct on the six $\text{PGL}(2, 13)$ -orbitals. Hence these intrinsic counts recover all six colors of the elliptic association scheme.

After recovering the passant color, form its seven-cliques. Exactly 78 have zero concurrence on every one of their triples. Direct incidence comparison identifies them with the supports of the 78 rows of M . Pair and triple concurrence are intrinsic to the minimum-support hypergraph, so this reconstruction is independent of the original projective labels.

4 Spanning and automorphisms

For internal points $P = (x_P : y_P : z_P)$ and Q , set

$$\Delta(P) = y_P^2 - x_P z_P, \quad \beta(P, Q) = 2y_P y_Q - x_P z_Q - z_P x_Q,$$

and

$$\rho(P, Q) = \frac{\beta(P, Q)^2}{\Delta(P)\Delta(Q)}.$$

On distinct internal points, ρ takes the six values 0, 1, 3, 9, 10, 12. Its level relations are the elliptic scheme; write A_r for their adjacency matrices. Polarity identifies A_0 with M , up to row order.

The needed part of the integral intersection table reduces modulo two to

$$\begin{aligned} A_0^2 &= I + A_9 + A_{10} + A_{12}, \\ A_0 A_r &= 0 \quad (r = 9, 10, 12), \\ A_9^2 &= A_{10}, \quad A_{10}^2 = A_{12}, \quad A_{12}^2 = A_9. \end{aligned} \tag{1}$$

These are parity reductions of intersection numbers. The $\text{PGL}(2, 13)$ -action is transitive on each relation, so fixing one point and substituting one representative of every relation checks all entries.

Put $B = A_9$. Since $A_0 B = 0$, the image of B lies in K . If $x \in K$ and $Bx = 0$, then (1) gives

$$0 = A_0^2 x = (I + B + B^2 + B^4)x = x.$$

Thus B is injective on the 36-dimensional space K , so $\text{im } B = K$, and B, B^2, B^4 all have rank 36. If N_i is the 91-by-78 support matrix of the i -th minimum-word orbit, direct pair counting in one representative gives

$$(N_1^\top N_1, N_2^\top N_2, N_3^\top N_3, N_4^\top N_4) = (A_9, A_9, A_{12}, A_{10}).$$

Every N_i therefore has rank at least 36. Its rows lie in K , so each orbit spans K .

It remains to determine the scheme automorphisms. Use the four anchors

$$P_0 = (1 : 0 : 2), \quad P_1 = (1 : 1 : 7), \quad P_2 = (1 : 0 : 7), \quad P_3 = (1 : 1 : 3).$$

The rigidity mechanism is summarized by

$$\underbrace{(P_0, P_1, P_2)}_{(10,3,9) \text{ regular orbit}} \implies \underbrace{P_3}_{(3,1,9) \text{ is unique}} \implies \underbrace{P \mapsto ([P = P_i], \rho(P, P_i))_{i=0}^3}_{\text{injective on the 78 points}}.$$

The first three satisfy

$$(\rho(P_0, P_1), \rho(P_0, P_2), \rho(P_1, P_2)) = (10, 3, 9).$$

There are $78 \cdot 14 \cdot 2 = 2184$ ordered triples with this pattern: A_{10} has valency 14 and $p_{3,9}^{10} = 2$. Substitution in the symmetric-square action shows that the stabilizer of (P_0, P_1, P_2) in $\text{PGL}(2, 13)$ is trivial. Since the group also has order 2184, it acts simply transitively on these triples.

For a fixed triple, P_3 is the unique point with relation signature $(3, 1, 9)$. The four values

$$(\rho(P, P_0), \rho(P, P_1), \rho(P, P_2), \rho(P, P_3)),$$

with equality to an anchor recorded on the diagonal, distinguish all 78 points. Consequently a scheme automorphism can be composed with a unique element of $\text{PGL}(2, 13)$ to fix the first three anchors; it then fixes the fourth and every point. Thus the reconstructed scheme has automorphism group $\text{PGL}(2, 13)$.

Every coordinate automorphism of K preserves $\mathcal{H}$. Conversely, $\mathcal{H}$ spans K , so every automorphism of $\mathcal{H}$ preserves K . The two coordinate-permutation groups coincide. This completes the proof of Theorem 1.1.

5 Proof and verification boundary

Parity and pencil geometry give the low-weight reductions; the tangent product controls weight eight; the association algebra explains spanning; and concurrence with four anchors gives reconstruction and rigidity. Three steps remain finite: the two weight-ten syndrome exclusions, the fixed-point classification of the minimum words, and the bounded clique and anchor checks.

The accompanying `verification/` directory contains the canonical weight-ten generator, an independent replay with a separately reconstructed incidence matrix, and the main exact replay. The last records the four minimum-word representatives and checks the clique closure, rank, orbit exhaustion, concurrence reconstruction, spanning identities, and anchors. The entry point `verify_evidence.py` checks the recorded byte counts and SHA-256 hashes before running all three programs. Every computation is deterministic and uses exact finite-field or binary arithmetic.

The shared Lean library formalizes the normalized conic and incidence code, the two parity-profile reductions, the tangent graph, the binary association argument, and the reconstruction and anchor interfaces. The paper-owned package checks the finite leaves and transports them to the following manuscript-facing conclusions:

- rank 42 and code dimension 36;
- exclusion of the normalized weight-eight case and reduction of every weight-ten word to the two profiles in Section 2;
- the 56-word fixed-point exhaustion and the four rank-36 orbit spans;
- uniqueness of the 78 reconstructed passant rows; and
- every automorphism of the six-valued relation scheme is projective.

The paper-owned aggregate namespace is `PassantCodeQ13.Gates.Main`; its terminals are

```
incidenceRankAndCodeDimension
fixedPointWeightTwelveExhaustion
minimumOrbitsSpanRhoZeroKernel
recoveredRowFamilyIsUnique
ellipticSchemeAutomorphismsAreProjective.
```

The shared low-weight namespace is `RelativeConicArcs.Gates.PassantCodeQ13`; its terminals are

```
weightEight_semantic_transport
arbitrary_weightTen_profile_transport.
```

Native evaluation discharges the finite leaves. Under Lean 4.32.0-rc1, the tracked audit reports the axioms of each terminal. The formal scope stops at two human transports. Projective transitivity and the double count carry the fixed-point exhaustion to all 364 minimum words. Concurrence carries minimum-support automorphisms to automorphisms of the polar-relation scheme. Segre’s lemma of tangents is the cited classical input to the formal weight-eight transport.

6 Conclusion

The code determines a row space, but its minimum words recover more: they distinguish the original parity-check rows inside that space. At $q = 13$, concurrence recovers the elliptic relations, the binary association algebra explains why every minimum orbit spans, and four anchors force the full projective symmetry.

References

- [1] S. Ball and M. Lavrauw, Arcs in finite projective spaces, *EMS Surveys in Mathematical Sciences* **6** (2019), 133–172. arXiv:1908.10772.
- [2] S. V. Droms, K. E. Mellinger, and C. Meyer, LDPC codes generated by conics in the classical projective plane, *Designs, Codes and Cryptography* **40** (2006), 343–356. doi:10.1007/s10623-006-0022-6.
- [3] H. D. L. Hollmann and Q. Xiang, Association schemes from the action of $\text{PGL}(2, q)$ fixing a nonsingular conic in $\text{PG}(2, q)$, *Journal of Algebraic Combinatorics* **24** (2006), 157–193. doi:10.1007/s10801-006-0005-8; arXiv:math/0503573.
- [4] A. L. Madison and J. Wu, On binary codes from conics in $\text{PG}(2, q)$, *European Journal of Combinatorics* **33** (2012), 33–48. doi:10.1016/j.ejc.2011.08.001; arXiv:1104.0324.
- [5] L. Ma, S. Liu, and Z. Tian, The binary codes generated from quadrics in projective spaces, *AIMS Mathematics* **9** (2024), 29333–29345. doi:10.3934/math.20241421.

AI assistance disclosure

This project was developed with extensive assistance from OpenAI Codex and Anthropic Claude. Under the author's direction, the systems assisted with proof exploration, literature research, symbolic and finite computation, code and formal-proof development, verification, and manuscript drafting and revision. The author checked the resulting arguments, computations, code, and cited sources, reviewed and edited all AI-assisted material, and assumes responsibility for all content.