

# Complete Bounded Repair Ports: Transfer, Reliability, and Geometric Structure

Tavis Rudd

July 2026

## Abstract

For a linear code and a distinguished coordinate, the complete bounded repair port records every dual-support recovery using at most a prescribed number of helpers. We keep three layers of this object visible: its support clutter, its scalar recovery coefficients, and its reliability under helper failures.

We prove an exact weighted-functional transfer theorem for concatenation and identify the persistent pointed obstruction to replicating a prescribed port. Every fixed represented port satisfying this obstruction bound occurs with positive density in an asymptotically good fixed-alphabet family. The same object has an exact deletion-contraction reliability calculus and a radius-truncated BEC EXIT hierarchy whose successive differences give the cheapest available repair radius. At full radius, repair reliability is a specialization of the Las Vergnas polynomial of the elementary perspective  $M \setminus x \rightarrow M/x$ ; the bounded-radius filtration is extra information.

Two characteristic-three families illustrate the distinction. A twisted-cubic-axis code has exact coordinatewise matching and transversal rows and transfers strictly beyond the ordinary support-distance gate. A quartic normal rational curve with its nucleus has harmonic-quadruple circuits, a Steiner  $S(3, 4, q + 1)$  nucleus port, and parallel repairs at the nucleus, whereas every curve-target repair has the nucleus as a compulsory helper. These parallel and series geometries separate bounded-radius from full-port information.

## 1 Introduction

Locality asks how many surviving symbols suffice to recover one unavailable coordinate. Availability asks for many disjoint choices of helpers. Both are standard in locally repairable codes [7, 13, 9], but neither describes the whole local failure event: two codes can have the same locality and matching number while their permitted repair sets overlap very differently.

The natural pointed object is the family of *all* bounded dual-support repairs. We call it a complete bounded repair port. Its matching number is disjoint availability, its transversal number minus one is worst-case helper failure tolerance, and its monotone Boolean function gives exact stochastic reliability. The coefficient layer matters as well: a repair support comes from a scalar equation, and concatenation is controlled by the minimum cost of realizing the corresponding block functional rather than by support alone.

The paper has six parts. Section 2 fixes the complete pointed object and separates support, coefficient, and probability layers. Section 3 gives exact and usable concatenation gates. Section 4 turns the pointed gate into a prescribed positive-density realization theorem. Section 5 develops reliability and bounded EXIT. Section 6 identifies the full-port polynomial as standard pointed-Tutte structure and isolates what the radius filtration adds. Section 7 compares cubic-axis and quartic-nucleus flagships.

The asymptotic coding ingredients, reliability identities, and pointed-Tutte polynomial are classical. We develop their exact organization around the complete bounded pointed object, together with the transfer boundary, prescribed-port consequence, and the two exact geometric port inventories. We make no minimum-bandwidth claim, no full-MAP claim for a radius cutoff, and no threshold claim for harmonic closure. In the literature reviewed, we did not locate the exact complete-port confinement criterion or the two coordinatewise port inventories [13, 10, 8, 5]; this is a bounded comparison, not a priority claim.

## 2 Complete bounded repair ports

Let  $C \leq \mathbb{F}_q^X$  be a linear code, let  $x \in X$ , and let  $r \geq 0$ .

**Definition 2.1** (Support layer). The complete radius- $r$  repair hypergraph at  $x$  is

$$\mathcal{H}_x^{(\leq r)}(C) = \left\{ R \subseteq X \setminus \{x\} : |R| \leq r, \exists w \in C^\perp, w_x \neq 0, \text{supp}(w) \setminus \{x\} = R \right\}.$$

Its minimal members form the repair clutter  $\min \mathcal{H}_x^{(\leq r)}(C)$ . Write  $\nu_x^{(\leq r)}$  and  $\tau_x^{(\leq r)}$  for its matching and transversal numbers.

The distinction between the complete hypergraph and its minimal clutter is useful: the former records every bounded witness, while every monotone support invariant below depends only on the latter.

**Proposition 2.2** (Basic invariants). *Assume the empty helper set is not a repair. Then*

$$\nu_x^{(\leq r)} \leq \tau_x^{(\leq r)}.$$

*Both numbers are unchanged after deleting nonminimal repair sets, so either the complete hypergraph or its minimal clutter may be used for these optima.*

*Proof.* A matching of  $m$  nonempty edges forces every transversal to select at least one point from each edge. Minimalization preserves the upward closure of the success event, hence preserves matching and transversal optima.  $\square$

**Definition 2.3** (Coefficient layer). Write

$$\mathcal{P}_x^{(\leq r)}(C) = \{w \in C^\perp : w_x = 1, |\text{supp}(w) \setminus \{x\}| \leq r\}$$

for the normalized coefficient port. For  $R \in \mathcal{H}_x^{(\leq r)}(C)$ , its coefficient fiber is the set of normalized coefficient vectors

$$\left( -\frac{w_y}{w_x} \right)_{y \in R} \quad (w \in C^\perp, \text{supp}(w) = R \cup \{x\}).$$

Each vector gives the exact recovery equation  $c_x = \sum_{y \in R} (-w_y/w_x) c_y$ . The direct protocol reads one complete  $\mathbb{F}_q$  symbol from every helper; no subpacketized access or bandwidth optimum is asserted.

**Definition 2.4** (Probability layer). Let  $A \subseteq X \setminus \{x\}$  be the surviving helpers and put

$$f_x^{(\leq r)}(A) = 1 \iff \exists R \in \mathcal{H}_x^{(\leq r)}(C) \text{ with } R \subseteq A.$$

For independent survival probabilities  $(s_y)$ , the reliability is  $R_x^{(\leq r)}(s) = \mathbb{E}[f_x^{(\leq r)}]$ .

Exact equality of complete ports preserves all support-derived invariants: matching, transversals, blocker clutters, multivariate reliability, and every correlated-law reliability after transporting the helper labels. Coefficient fibers require the finer functional data used next.

### 3 Exact weighted-functional transfer

Let  $I \leq \mathbb{F}_q^m$  be an inner  $[m, k, d]_q$  code and identify its message space with  $L = \mathbb{F}_{q^k}$  through an  $\mathbb{F}_q$ -linear encoder  $e : L \rightarrow I$ . For  $w \in \mathbb{F}_q^m$ , define the induced block functional

$$\Phi_I(w) : L \longrightarrow \mathbb{F}_q, \quad a \longmapsto \langle w, e(a) \rangle.$$

The zero fiber is  $I^\perp$ . For  $\beta \in \text{Hom}_{\mathbb{F}_q}(L, \mathbb{F}_q)$  put

$$\lambda(\beta) = \min_{\Phi_I(w)=\beta} \text{wt}(w), \quad \mu_x(\beta) = \min_{\substack{\Phi_I(w)=\beta \\ w_x \neq 0}} \text{wt}(w),$$

with value  $\infty$  for an empty constrained fiber.

Let  $O \leq L^J$  be an outer code. Its functional dual consists of tuples  $(\beta_j)_{j \in J}$  satisfying

$$\sum_{j \in J} \beta_j(a_j) = 0 \quad \text{for every } a = (a_j) \in O.$$

The weighted cost of such a tuple is  $\sum_j \lambda(\beta_j)$ . This is the ordinary quotient/coset-leader weight attached to the inner encoder; the pointed use below is the repair-specific part. Write

$$\text{fwt}(\beta) = |\{j : \beta_j \neq 0\}|.$$

Thus the functional-dual tuples have three disjoint strata:  $\text{fwt}(\beta) = 0$ ,  $\text{fwt}(\beta) = 1$ , and  $\text{fwt}(\beta) \geq 2$ . Keeping the singleton stratum is essential when an outer coordinate projection is not surjective. If every coordinate projection of  $O$  is surjective, a functional-dual tuple cannot have functional weight one.

The all-zero functional sector has the fixed pointed cost

$$z_x(I) = \begin{cases} \mu_x(0) + d(I^\perp), & I^\perp \neq 0, \\ \infty, & I^\perp = 0. \end{cases}$$

**Theorem 3.1** (Exact pointed confinement and transfer). *Assume  $|J| \geq 2$ , and fix an outer block  $j$  and an inner coordinate  $x$ . The minimum weight of a concatenated dual witness through  $(j, x)$  that is not an embedded inner-dual witness is*

$$\min \left\{ z_x(I), \min_{\substack{(\beta_\ell) \text{ in the functional dual of } O \\ (\beta_\ell) \neq 0}} \left( \mu_x(\beta_j) + \sum_{\ell \neq j} \lambda(\beta_\ell) \right) \right\},$$

where a minimum over an empty set is  $\infty$ . Consequently, if this pointed cost is greater than  $r + 1$ , then

$$\mathcal{H}_{(j,x)}^{(\leq r)}(O \circ I) = \text{embed}_j(\mathcal{H}_x^{(\leq r)}(I)).$$

Under the same hypothesis, zero-extension is a bijection

$$\mathcal{P}_x^{(\leq r)}(I) \xrightarrow{\sim} \mathcal{P}_{(j,x)}^{(\leq r)}(O \circ I).$$

In particular it suffices that

$$r + 1 < 2d(I^\perp) \quad \text{and} \quad \sum_j \lambda(\beta_j) \geq r + 2$$

for every nonzero functional-dual tuple.

*Proof.* Let  $w = (w_\ell)_{\ell \in J}$  be a block word and set  $\beta_\ell = \Phi_I(w_\ell)$ . For every outer word  $a = (a_\ell) \in O$ ,

$$\langle w, e(a) \rangle = \sum_{\ell \in J} \langle w_\ell, e(a_\ell) \rangle = \sum_{\ell \in J} \beta_\ell(a_\ell).$$

Hence  $w \in (O \circ I)^\perp$  if and only if  $\beta = (\beta_\ell)$  belongs to the functional dual of  $O$ . This calculation is the only concatenated-dual decomposition used below; no fiber enumerator is needed.

Fix such a tuple  $\beta$ . The constraints on the blocks are independent. At the target, the condition  $w_j(x) \neq 0$  gives minimum  $\mu_x(\beta_j)$ . At every other block the minimum is  $\lambda(\beta_\ell)$ . Each ordinary fiber is nonempty because a functional on the encoded inner space extends to the ambient coordinate space, and its minimum is attained because Hamming weight takes values in the well-ordered set  $\mathbb{N}$ . The constrained target fiber is either empty, in which case its cost and the fixed-fiber cost are infinite, or has an attained minimum by the same argument. Therefore the exact pointed minimum in the fixed fiber  $\beta$  is

$$\mu_x(\beta_j) + \sum_{\ell \neq j} \lambda(\beta_\ell).$$

If  $\beta \neq 0$ , a realization cannot be an embedded inner-dual word in block  $j$ , since every embedded inner-dual word induces the zero functional in every block. Taking the minimum over the nonzero functional dual gives the second term in the theorem. This minimum includes both remaining strata: functional weight one and functional weight at least two.

It remains to analyze  $\beta = 0$ . Then every  $w_\ell$  lies in  $I^\perp$ . A pointed realization that is not embedded in block  $j$  consists of a word in  $I^\perp$  that is nonzero at  $x$ , together with a nonzero word of  $I^\perp$  in at least one off-target block. Conversely, placing any such pair in two blocks and putting zero elsewhere gives a pointed nonembedded concatenated-dual word. Since  $|J| \geq 2$ , the two choices can be made independently, and their exact minimum is  $\mu_x(0) + d(I^\perp) = z_x(I)$ . If either required word does not exist, both the constrained minimum and  $z_x(I)$  are infinite. The zero and nonzero functional sectors partition all pointed nonembedded witnesses, proving the displayed minimum formula.

Suppose that this minimum is greater than  $r + 1$ , and take a radius- $r$  pointed repair witness in the concatenated code. Its total weight is at most  $r + 1$ , so it cannot be nonembedded. It is therefore the zero-extension of an inner-dual word in block  $j$ . Deleting the target coordinate identifies its helper support with the image under  $\text{embed}_j$  of a radius- $r$  inner helper support. This proves one inclusion of the asserted support-port equality. If the witness is normalized at the target, its nonzero block is the correspondingly normalized inner word, so this identification also gives the asserted coefficient-port bijection. Conversely, the zero-extension of every radius- $r$  inner repair word is orthogonal to the concatenated code, and its helper support is the corresponding embedded support; normalization is unchanged. This proves the reverse inclusions in both layers.

Finally, whenever the zero-sector cost is finite,  $\mu_x(0) \geq d(I^\perp)$ , so  $z_x(I) \geq 2d(I^\perp)$ . For a nonzero functional tuple,  $\mu_x(\beta_j) \geq \lambda(\beta_j)$ , and hence its pointed cost is at least  $\sum_\ell \lambda(\beta_\ell)$ . The two displayed numerical hypotheses therefore put every pointed nonembedded witness above weight  $r + 1$ , and the transfer conclusion follows.  $\square$

The functional-dual decomposition is classical concatenation machinery [2, Theorem 2.3][3, Theorem 2.1]. The weighted statement matters because ordinary functional support distance can be strictly weaker. For the completed field-nine cubic-axis inner code, a Singer-shifted generalized single-parity-check  $[5, 4, 2]_{6561}$  outer code has functional distance five but weighted distance at least six. Its exact nonembedded threshold is six, so radius-four transfer holds although the support-distance gate requiring functional distance six fails. Here Singer-shifted means that the five parity-

check multipliers are chosen through a regular Singer action so that the inner code's twenty unit-cost projective functional classes have disjoint translates in the five blocks. This finite calculation belongs to the checked transfer chain described in Section 8; Singer regularity is its only classical group input [15].

## 4 Prescribed positive-density realization

The pointed zero-functional sector persists even when outer dual distance grows.

**Theorem 4.1** (Prescribed represented ports). *Let  $O_N \leq L^N$  be  $L$ -linear outer codes with  $d(O_N^\perp) \rightarrow \infty$ , and let  $C_N = O_N \circ I$ . For all sufficiently large  $N$ , every pointed dual witness through  $(j, x)$  of weight at most  $r + 1$  is confined to block  $j$  if and only if*

$$r + 1 < z_x(I).$$

*Under this condition the radius- $r$  port at  $x$  occurs at the  $N$  targets  $(j, x)$ , hence with density  $1/m$  in  $C_N$ . More precisely, at every block*

$$\mathcal{H}_{(j,x)}^{(\leq r)}(C_N) = \text{embed}_j(\mathcal{H}_x^{(\leq r)}(I)), \quad \mathcal{P}_{(j,x)}^{(\leq r)}(C_N) = \text{embed}_j(\mathcal{P}_x^{(\leq r)}(I)).$$

*Proof.* Every  $\mathbb{F}_q$ -linear functional on  $L$  has a unique form  $a \mapsto \text{Tr}_{L/\mathbb{F}_q}(ba)$ , by nondegeneracy of the trace pairing. Let  $\beta = (\beta_j)$  lie in the functional dual of the restricted outer code, and write  $\beta_j(a) = \text{Tr}(b_j a)$ . Since  $O_N$  is  $L$ -linear, applying  $\beta$  to every scalar multiple  $ca$  of every  $a \in O_N$  shows

$$\text{Tr}\left(c \sum_j b_j a_j\right) = 0 \quad \text{for every } c \in L.$$

Trace nondegeneracy gives  $(b_j) \in O_N^\perp$ . Conversely, every outer dual word gives such a functional tuple. The trace representation is coordinatewise injective, so it preserves support exactly. Thus the functional-dual support distance is  $d(O_N^\perp)$ .

Fix  $r$ . For all sufficiently large  $N$ ,  $d(O_N^\perp) \geq r + 2$ . Every realization of a nonzero functional tuple then has weight at least  $r + 2$ : each nonzero functional requires a nonzero inner block. The exact pointed decomposition of Theorem 3.1 consequently leaves only the zero-functional sector in the bounded range. For  $N \geq 2$ , its exact cost is the block-independent quantity  $z_x(I)$ . Hence confinement through every  $(j, x)$  holds exactly when  $r + 1 < z_x(I)$ .

If  $r + 1 < z_x(I)$ , Theorem 3.1 identifies both support and normalized coefficient ports by zero-extension. If  $r + 1 \geq z_x(I)$ , the attained pointed inner-dual minimum at  $x$ , together with a minimum nonzero inner-dual word in a second block, gives a zero-functional pointed nonembedded witness of weight at most  $r + 1$ . This proves necessity as well as sufficiency. Finally, the  $N$  targets  $(j, x)$  among the  $mN$  concatenated coordinates give density  $1/m$ .  $\square$

**Proposition 4.2** (Concatenated parameters). *If  $I$  has parameters  $[m, k, d]_q$  and  $O_N \leq L^N$  has  $L$ -dimension  $K_N$  and symbol distance  $D_N$ , then  $C_N = O_N \circ I$  has length  $mN$ ,  $\mathbb{F}_q$ -dimension  $kK_N$ , and minimum distance at least  $dD_N$ . In particular, if*

$$\frac{K_N}{N} \rightarrow R > 0, \quad \liminf_N \frac{D_N}{N} \geq \delta > 0,$$

*then the concatenated family has asymptotic rate  $kR/m$  and relative distance at least  $d\delta/m$ .*

*Proof.* The blockwise inner encoder is injective, so restriction of scalars multiplies the outer  $L$ -dimension by  $[L : \mathbb{F}_q] = k$ . A nonzero outer word has at least  $D_N$  nonzero symbols, and each nonzero symbol encodes to an inner word of weight at least  $d$ . The finite parameter statements follow; division by  $mN$  gives the asymptotic claims.  $\square$

**Corollary 4.3** (Positive-density MDS coefficient fingerprints). *Let  $I$  be a positive-dimensional  $[m, k]_q$  MDS code and fix  $x$ . Its radius- $k$  support port is the complete  $k$ -uniform clutter on the other coordinates, while its radius- $k$  coefficient port reconstructs  $I$ , and  $z_x(I) = 2(k + 1)$ . Both ports occur at density  $1/m$  in every sufficiently long concatenation by an outer family with dual distance tending to infinity.*

*Proof.* The minimum-port assertions are the MDS reconstruction theorem. Every minimum dual support through  $x$  has size  $k + 1$ , so  $\mu_x(0) = d(I^\perp) = k + 1$  and

$$z_x(I) = 2(k + 1) > k + 1.$$

Apply Theorem 4.1 with  $r = k$ . The support port depends only on the MDS parameters; the transferred normalized coefficients retain the represented inner code.  $\square$

**Corollary 4.4** (Positive-density Clebsch repair ports). *Let  $I$  be the  $[6, 3, 4]_{11}$  Clebsch hexagon code [14], and fix a coordinate  $x$ . The minimal radius-three repair clutter at  $x$  is the complete three-uniform hypergraph on the other five coordinates. Hence it has ten minimal repair supports,*

$$\nu_x^{(\leq 3)} = 1, \quad \tau_x^{(\leq 3)} = 3,$$

*and homogeneous helper-survival reliability*

$$10s^3(1 - s)^2 + 5s^4(1 - s) + s^5.$$

*Already the radius-three coefficient port at  $x$  determines  $I$ . Moreover  $z_x(I) = 8$ , so both the minimum radius-three port and the full radius-five coefficient port occur with density  $1/6$  in an asymptotically good fixed- $\mathbb{F}_{11}$  family.*

*Proof.* The dual  $I^\perp$  is again an  $[6, 3, 4]_{11}$  MDS code. Every four-set containing  $x$  is therefore the support of a dual word, unique up to scalar. The minimal helper sets are consequently all  $\binom{5}{3} = 10$  triples on the other coordinates. Their matching and transversal numbers are one and three, and recovery succeeds exactly when at least three of the five helpers survive, giving the displayed polynomial.

At radius three the common-core star of minimum normalized dual words is already a basis of  $I^\perp$ , by Corollary 4.3. Thus the minimum coefficient port, and hence also the full radius-five port, recovers  $I^\perp$  and  $I$ .

Finally, the minimum dual-word weight through  $x$  is four, so

$$z_x(I) = \mu_x(0) + d(I^\perp) = 4 + 4 = 8.$$

Taking  $r = 5$  gives  $r + 1 = 6 < 8$ . Theorem 4.1 therefore confines every full pointed witness to its inner block and reproduces the coefficient fibers at one of the six coordinate types in every block. Choosing asymptotically good outer codes over  $\mathbb{F}_{11^3}$  gives the claimed fixed- $\mathbb{F}_{11}$  family and density.  $\square$

**Remark 4.5.** The support clutter and reliability polynomial in Corollary 4.4 are shared by every  $[6, 3, 4]$  MDS code. The Clebsch-specific information lies in the scalar coefficient fibers: they recover the inner code that is characterized geometrically in [14].

**Remark 4.6.** Corollary 4.3 applies without a new transfer argument to every represented MDS inner code. This includes the usual arc and projective Reed–Solomon presentations, and the classical MDS code presentations underlying AME constructions, whenever their displayed field and dimension hypotheses hold. Their minimum support ports are generic MDS data; only their normalized coefficient ports distinguish the represented geometric or quantum-code realization.

This is a represented-port theorem: it does not say that every abstract hypergraph is linearly representable. It yields standard scaled asymptotic regions. If  $Q = q^k$ , random  $L$ -linear outer codes of rate  $R$  can have relative distances  $\delta, \delta^\perp$  whenever

$$H_Q(\delta) < 1 - R, \quad H_Q(\delta^\perp) < R.$$

For an inner  $[m, k, d]_q$  code the concatenated parameters satisfy

$$R_{\text{concat}} \rightarrow \frac{k}{m}R, \quad \delta_{\text{concat}} \geq \frac{d}{m}\delta.$$

Indeed, for a random  $L$ -linear outer code the expected numbers of nonzero words and dual words below the two cutoffs have exponential rates  $R + H_Q(\delta) - 1$  and  $H_Q(\delta^\perp) - R$ , respectively, so both vanish simultaneously; this is the random-linear-code form of the Gilbert–Varshamov argument [11, Chapter 17]. When  $Q$  is square, algebraic-geometry outer codes similarly give, for  $\gamma = 1/A(Q)$  and  $\gamma < R < 1 - \gamma$ ,

$$\delta(O_N) \geq 1 - R - \gamma, \quad \delta(O_N^\perp) \geq R - \gamma, \quad \delta_{\text{concat}} \geq \frac{d}{m}(1 - R - \gamma).$$

These are the classical GV and TVZ ingredients. For the completed  $[20, 4, 9]_9$  seed,  $Q = 6561$  and  $\gamma \leq 1/80$ , recovering the concrete rate-1/10 family as one instance [17, Chapters 2 and 7]. The stronger self-dual family used for that concrete instance is precisely Stichtenoth’s theorem over  $\mathbb{F}_{6561}$  [16, Theorem 1.6(ii)].

## 5 Reliability and bounded EXIT

Let  $\mathcal{H}$  be a repair clutter on helper set  $V$ . For  $v \in V$ , define

$$\mathcal{H} - v = \min\{E \in \mathcal{H} : v \notin E\}, \quad \mathcal{H}/v = \min\{E \setminus \{v\} : E \in \mathcal{H}\}.$$

**Theorem 5.1** (Reliability calculus). *For independent helper survival probabilities  $(s_v)$ ,*

$$R_{\mathcal{H}}(s) = (1 - s_v)R_{\mathcal{H}-v}(s) + s_v R_{\mathcal{H}/v}(s),$$

and

$$\frac{\partial R_{\mathcal{H}}}{\partial s_v} = R_{\mathcal{H}/v} - R_{\mathcal{H}-v} = \Pr(v \text{ is pivotal}).$$

*Under homogeneous survival  $s$ ,  $R'_{\mathcal{H}}(s) = \sum_v \Pr_s(v \text{ is pivotal})$ . If the minimum blocker size is  $\tau$  and there are  $b_\tau$  minimum blockers, then for helper failure probability  $p$ ,*

$$1 - R_{\mathcal{H}}(1 - p) = b_\tau p^\tau + O(p^{\tau+1}).$$

*Proof.* Write  $f_{\mathcal{H}}(A) = 1$  when  $A$  contains an edge of  $\mathcal{H}$ , and write it as zero otherwise. The reliability is the finite multilinear sum

$$R_{\mathcal{H}}(s) = \sum_{A \subseteq V} f_{\mathcal{H}}(A) \prod_{y \in A} s_y \prod_{y \notin A} (1 - s_y).$$

Partition the subsets according to whether they contain  $v$ . Erasing  $v$  from those that do contain it identifies the two parts with the Boolean systems  $\mathcal{H} - v$  and  $\mathcal{H}/v$ , respectively. This gives

$$R_{\mathcal{H}} = (1 - s_v)R_{\mathcal{H}-v} + s_v R_{\mathcal{H}/v}.$$

The two conditional reliabilities do not involve  $s_v$ , so differentiation gives  $R_{\mathcal{H}/v} - R_{\mathcal{H}-v}$ . Monotonicity makes this difference the conditional probability that adjoining  $v$  changes failure into success, which is precisely pivotality. On the diagonal  $s_v = s$ , the chain rule sums these coordinate derivatives and gives the Russo–Margulis identity.

Now expose the failed set  $F$ . Repair fails exactly when  $F$  is a transversal of  $\mathcal{H}$ , equivalently when it contains a minimal blocker. Thus

$$1 - R_{\mathcal{H}}(1 - p) = \sum_{\substack{F \subseteq V \\ F \text{ is a transversal}}} p^{|F|} (1 - p)^{|V| - |F|}.$$

There are no summands below degree  $\tau$ , and the transversals of size  $\tau$  are exactly the  $b_{\tau}$  minimum blockers. Their contribution is  $b_{\tau} p^{\tau} (1 - p)^{|V| - \tau} = b_{\tau} p^{\tau} + O(p^{\tau+1})$ . Every remaining term contains at least  $p^{\tau+1}$ ; the sum is finite, proving the expansion.  $\square$

These are the standard deletion–contraction and pivotality identities for a monotone reliability system [4].

For EXIT conventions, condition the target  $x$  unavailable and erase helper  $v$  with probability  $p_v$ . Put

$$h_x^{(\leq r)}(p) = \Pr(f_x^{(\leq r)}(A) = 0), \quad R_x^{(\leq r)} = 1 - h_x^{(\leq r)}.$$

This is an *extrinsic* failure probability. If the target is itself erased with probability  $p_x$ , its residual erasure probability is  $p_x h_x^{(\leq r)}$ . At full radius it is symbol-MAP EXIT; at finite radius it is a bounded-query decoder and carries no capacity claim [1, 12].

**Proposition 5.2** (Bounded-EXIT hierarchy). *The failure form of deletion–contraction is*

$$h_{\mathcal{H}}(p) = p_v h_{\mathcal{H}-v}(p) + (1 - p_v) h_{\mathcal{H}/v}(p), \quad \frac{\partial h_{\mathcal{H}}}{\partial p_v} = \Pr(v \text{ is pivotal}).$$

If  $L_x(A)$  is the size of the cheapest repair contained in the realized survivor set, with  $L_x = \infty$  when none exists, then

$$\Pr(L_x = r) = h_x^{(\leq r-1)} - h_x^{(\leq r)}, \quad \Pr(L_x = \infty) = h_x^{\text{MAP}}.$$

The first identity is for  $r \geq 1$ .

*Proof.* Conditioning on whether  $v$  is erased gives the displayed recurrence: when it is erased, only repairs avoiding  $v$  remain; when it survives, it may be contracted from every repair. Equivalently, substitute  $s_y = 1 - p_y$  in the reliability recurrence and take complements. Differentiation reverses both signs, so the derivative is again the pivotal probability.



Let  $\mathcal{H}_x^{(\leq r)}$  be the full port filtered by edge size at most  $r$ . The event  $L_x \leq r$  is exactly success for this truncated port. Hence

$$\{L_x = r\} = \{L_x > r - 1\} \setminus \{L_x > r\},$$

and taking probabilities gives the difference of failure curves. At full radius, the complementary event is that no repair exists, which is  $L_x = \infty$ .  $\square$

Minimal transversals of  $\mathcal{H}_x^{(\leq r)}$  are exact target-specific, one-shot bounded-repair failure certificates. They are not automatically the representation-dependent stopping sets of a global iterative Tanner decoder.

For a length- $N$  code define the radius- $r$  locality deficit

$$L_r(x) = \int_0^1 (h_x^{(\leq r)}(p) - h_x^{\text{MAP}}(p)) dp.$$

If  $\Delta a_k$  counts size- $k$  survivor sets repaired by full MAP but not at radius  $r$ , beta integration gives

$$L_r(x) = \sum_{k=0}^{N-1} \frac{\Delta a_k}{N \binom{N-1}{k}}.$$

With entropy normalized in  $\log_q$  units, the unrecoverable-symbol indicator is the normalized conditional entropy on the  $q$ -ary erasure channel. Under this convention the symbol-MAP areas sum to the code dimension  $K$  [1], so

$$\sum_x \int_0^1 h_x^{(\leq r)}(p) dp = K + \sum_x L_r(x).$$

## 6 Pointed Tutte structure

Let  $M$  be the column matroid of a projective-system code on  $E = V \sqcup \{x\}$ , with  $x$  neither a loop nor a coloop. Put

$$\epsilon_x(A) = r_M(A \cup \{x\}) - r_M(A) \in \{0, 1\}.$$

The set  $A$  repairs  $x$  exactly when  $\epsilon_x(A) = 0$ .

**Theorem 6.1** (Pointed perspective polynomial). *Let  $D = M \setminus x$  and  $Q = M/x$ . The Las Vergnas polynomial of the elementary perspective  $D \rightarrow Q$  is*

$$T_{D \rightarrow Q}(X, Y, Z) = \sum_{A \subseteq V} (X - 1)^{r(M) - r_M(A \cup \{x\})} (Y - 1)^{|A| - r_M(A)} Z^{\epsilon_x(A)}.$$

If

$$S_x(u) = \sum_{\epsilon_x(A)=0} u^{|A|},$$

then, for  $n = |V|$ ,

$$R_{(M,x)}(s) = (1 - s)^n S_x\left(\frac{s}{1 - s}\right).$$

Writing  $U_K(u, y) = \sum_A u^{|A|} y^{r_K(A)}$ , one also has

$$S_x(u) = \frac{\partial}{\partial y} (U_D(u, y) - U_Q(u, y)) \Big|_{y=1}.$$

*Proof.* Contraction gives  $r_D(A) = r_M(A)$  and  $r_Q(A) = r_M(A \cup \{x\}) - 1$ , so  $r_D(A) - r_Q(A) = 1 - \epsilon_x(A)$ . Substitution in the rank-one perspective expansion gives the displayed polynomial. The  $Z^0$  terms are exactly the successful sets; more explicitly,

$$S_x(u) = u^{r(M)}[Z^0]T_{D \rightarrow Q}(1 + u^{-1}, 1 + u, Z),$$

which yields the reliability formula. Finally,

$$\partial_y(y^{r_D(A)} - y^{r_Q(A)}) \Big|_{y=1} = r_D(A) - r_Q(A),$$

which is the zero–one success indicator.  $\square$

This identification is the singleton specialization of standard set-pointed Tutte theory [19, equation (3.1)]; it is not a new polynomial. Pointed duality gives

$$R_{(M,x)}(s) + R_{(M^*,x)}(1 - s) = 1,$$

and the minimal failure blockers of  $(M, x)$  are exactly the minimal repair sets of  $(M^*, x)$ . For a represented matroid they are the sets  $Q' \setminus \{x\}$ , where  $Q'$  ranges over cocircuits containing  $x$ ; equivalently,  $Q'$  is an inclusion-minimal row-code support through  $x$ . In particular,

$$\tau_{\text{full}}(x) = d_x(C) - 1, \quad d_x(C) = \min\{\text{wt}(c) : c \in C, c_x \neq 0\}.$$

The full pointed polynomial forgets the repair radius. Six-element circuits in a rank-five code create five-helper repairs that are absent from a radius-four port. Thus the hierarchy in Section 5 is a genuine filtered refinement of the standard perspective invariant.

## 7 Geometric flagships

### 7.1 The twisted-cubic–axis family

Let  $q = 3^h$  and work in  $\text{PG}(3, q)$ . Put

$$T_q = \{C(t) = (1 : t : t^2 : t^3) : t \in \mathbb{F}_q\},$$

$$L_q = \{A(u) = (0 : 1 : u : 0) : u \in \mathbb{F}_q\} \cup \{A_\infty = (0 : 0 : 1 : 0)\}.$$

The line  $L_q$  is the characteristic-three axis of the osculating-plane pencil [5]; all statements below follow directly from the displayed coordinates.

**Theorem 7.1** (Cubic–axis port). *The row code on  $T_q \sqcup L_q$  is  $[2q + 1, 4, q - 1]_q$ . Its minimal circuits of size at most four are the triples on  $L_q$  and the sets*

$$\{C(s), C(t), C(u), (0 : e_1 : e_2 : 0)\}$$

for distinct  $s, t, u$ , where  $e_1 = s + t + u$  and  $e_2 = st + su + tu$ . Every cubic coordinate has

$$(\nu, \tau) = \left( \frac{q-1}{2}, q-2 \right).$$

If  $Z_3(q)$  is the maximum size of a three-term-zero-sum-free subset of  $(\mathbb{F}_q, +)$ , then

$$(\nu, \tau)_{A_\infty} = \left( \frac{5q-3}{6}, 2q-1-Z_3(q) \right),$$

$$(\nu, \tau)_{A(a)} = \left( \frac{5q-9}{6}, 2q-2-Z_3(q) \right).$$

In particular every coordinate satisfies  $\tau > \nu$  for  $q \geq 9$ .

*Proof sketch.* A plane containing  $L_q$  contains at most one cubic point by Frobenius injectivity; a plane not containing it meets the line once and the cubic at most three times. This gives the code parameters. Vandermonde determinants classify the displayed small circuits. At an axis target the repair clutter is a disjoint union of  $K_q$  and a zero-sum-triple hypergraph, so matching and transversal numbers add; complements of transversals are precisely zero-sum-free sets. At a cubic target, inversion relabels the other cubic helpers by  $\mathbb{F}_q^*$ . A multiplicative-generator pairing gives  $(q-1)/2$  repairs with distinct axis completions; no larger matching is possible because each repair uses two of the  $q-1$  cubic helpers. The shifted-inverse complement lemma bounds a repair-free helper set by  $q+2$ , attained by all axis helpers together with one cubic helper. Taking complements gives  $\tau = q-2$ . The stated strict inequalities follow from the elementary bound  $Z_3(q) \leq q-2$  and the displayed formulas.  $\square$

At  $q = 9$ ,  $Z_3(9) = 4$ , and the three coordinate types have exact rows

$$(4, 7), \quad (6, 12), \quad (7, 13),$$

with multiplicities 9, 9, 1. Restoring the cubic point at infinity gives a  $[2q+2, 4, q]_q$  code whose full minimal port is visible at radius four and has uniform rows

$$\left(\frac{q-1}{2}, q-1\right) \text{ on the cubic,} \quad \left(\frac{5q-3}{6}, 2q-3\right) \text{ on the axis.}$$

This is the inner code in the strict weighted-transfer example above.

## 7.2 The quartic-nucleus harmonic family

Let

$$V(t) = (1, t, t^2, t^3, t^4), \quad V(\infty) = e_4, \quad N = e_2,$$

and let  $S_q = \{V(t) : t \in \mathbb{P}^1(\mathbb{F}_q)\} \cup \{N\}$ . The classical normal-rational-curve nucleus formula identifies  $N$  as the hyperplane nucleus in characteristic three [6].

**Theorem 7.2** (Quartic-nucleus port). *For  $q = 3^h \geq 9$ , the row code  $Q_q$  on  $S_q$  has parameters*

$$[q+2, 5, q-3]_q, \quad d(Q_q^\perp) = 5.$$

*Its minimal circuits of size at most five are exactly*

$$\{N\} \cup B,$$

*where  $B$  is a harmonic quadruple of  $\mathbb{P}^1(\mathbb{F}_q)$ . These quadruples form a Steiner system  $S(3, 4, q+1)$ , the characteristic-three four-set orbit in the standard  $\text{PGL}(2, q)$  design construction [18, Section 2]. Hence the radius-four repairs are all  $B$  at the nucleus and all  $\{N\} \cup (B \setminus \{x\})$  at a curve target  $x \in B$ . Every curve target has  $(\nu, \tau) = (1, 1)$ . At  $q = 9$  the nucleus has  $(\nu, \tau) = (2, 5)$ , with 30 repairs, while each curve target has 12.*

*Proof.* For distinct finite  $a, b, c, d$ , the hyperplane through their quartic columns contains  $N$  exactly when  $e_2(a, b, c, d) = 0$ ; with  $\infty$  present the condition is  $a + b + c = 0$ . Given three finite points, if  $e_1 = a + b + c \neq 0$ , the unique completion is  $d = -e_2/e_1$ ; if  $e_1 = 0$ , the unique completion is  $\infty$ . The completion is distinct from the original triple. This proves the Steiner property and identifies the blocks as the projective images of  $\{\infty, 0, 1, -1\}$ .

Any five curve points are independent. The determinant criterion shows that  $\{N\} \cup B$  has rank four exactly for a harmonic block, and deleting any point restores independence. Thus there are no smaller circuits and the dual distance is five. A hyperplane contains at most four curve points and possibly  $N$ , with equality on a harmonic block, so the primal distance is  $(q + 2) - 5 = q - 3$ . The repair and row statements follow from the circuit inventory; the field-nine nucleus row is an exact finite enumeration.  $\square$

The inner-dual half of the coarse transfer gate is automatic at radius four because  $5 < 2d(Q_q^\perp) = 10$ . Theorem 4.1 therefore places the entire harmonic port with positive density in asymptotically good fixed- $\mathbb{F}_q$  families.

### 7.3 Reliability, EXIT, and the geometric contrast

The nucleus repairs are the blocks of the Steiner system  $S(3, 4, q + 1)$ . At a curve target, every radius-four repair contains the nucleus. Thus the nucleus port is a parallel family of four-helper repairs, whereas the curve port has a compulsory series helper. The reliability and bounded-EXIT theorem applies to both ports without an exact finite profile. Exact field-nine Bernstein rows, EXIT deficits, and Poisson approximations are computational refinements and are not used in the body theorem chain.

Finally, the harmonic small-circuit closure has a deterministic nucleus gate. Let  $D(S)$  close curve set  $S$  under “three points of a harmonic block add the fourth.” Then

$$\text{cl}_4(S) = S \quad \text{if } S \text{ contains no block,} \quad \text{cl}_4(S) = \{N\} \cup D(S) \quad \text{otherwise,}$$

and  $\text{cl}_4(S \cup \{N\}) = \{N\} \cup D(S)$ . At  $q = 9$  there is a block-free rank-five set of five curve points that linearly spans all eleven points but is inert under  $\text{cl}_4$ ; adjoining  $N$  completes all five missing curve points in one parallel round. This is a finite deterministic separation, not an asymptotic propagation or threshold theorem.

## 8 Verification and provenance

The coefficient-port reconstruction and exact transfer chains are checked in Lean. The import-only gate `RepairPorts.Gates.CompletePorts` audits the paper-facing declarations. Its two transfer terminals are

```
RepairPorts.exactFunctionalStrata,
RepairPorts.exactPointedConfinementAndTransfer.
```

The first gives the exact zero-, singleton-, and multisupport-functional profiles; the second gives the pointed minimum formula and complete-port equality. Both report only `propext`, `Classical.choice`, and `Quot.sound`. The strict field-nine corollary is the conclusion of

```
RepairCodes.projectiveAxisTwistedCubic_strict_weighted_transfer_of_regular_
projective_action.
```

Singer regularity is an explicit hypothesis of that declaration, not a hidden formal axiom. The exact public module closure and artifact provenance are release-gate data, not claims made by this private draft.

Trace duality, scaled concatenated parameters, exact target density, eventual necessity and sufficiency, support/coefficient transfer, and the MDS fingerprint corollary are audited through

```

RepairPorts.concatenatedRestrictedCode_parameters,
RepairPorts.eventually_pointedConfinement_iff_zeroCost,
RepairPorts.eventually_prescribedPorts,
RepairPorts.eventually_mdsMinimumCoefficientFingerprints.

```

They report only `propext`, `Classical.choice`, and `Quot.sound`. Random-GV or AG/TVZ existence of an outer family is the explicit classical input; no such existence theorem occurs in the Lean dependency closure.

The finite reliability and bounded-EXIT calculus is audited through

```

RepairPorts.portReliability_delete_contract,
RepairPorts.hasDerivAt_portReliability_update,
RepairPorts.hasDerivAt_homogeneous_portReliability,
RepairPorts.eraserFailureProbability_delete_contract,
RepairPorts.noRepairProbability_eq_eraserFailure,
RepairPorts.cheapestRepairRadiusProbability_eq_failure_sub,
RepairPorts.blockerFailurePolynomial_eq_minimum_term_add_remainder.

```

These declarations use finite sums and polynomial algebra only. Exact finite profiles and Poisson approximations remain computational refinements outside the body proof chain.

For the concrete completed field-nine family, the external existence input is Stichtenoth’s self-dual TVZ-family theorem over  $\mathbb{F}_{6561}$  [16, Theorem 1.6(ii)]. Classical inputs are labeled as such: outer-family existence, concatenated-dual decomposition, Singer regularity, normal-rational-curve nuclei, and the Las Vergnas perspective polynomial.

## 9 Conclusion

A complete bounded repair port is one pointed object with three compatible views. Its support clutter measures disjoint availability and adversarial tolerance; its coefficient fibers give exact scalar recovery and the correct weighted transfer cost; its Boolean function gives reliability and bounded EXIT. Exact concatenation transfers the whole object, and the persistent pointed obstruction precisely controls positive-density realization.

The cubic-axis family shows rich overlapping ports and strict weighted transfer beyond a support-only gate. The quartic-nucleus family replaces that overlap pattern by a harmonic Steiner design: nucleus repairs form a parallel block family while curve targets share a compulsory series helper. Pointed-Tutte theory explains the full-radius structure; the radius filtration records when each repair first becomes available.

A sharper cubic blocker-stability statement is not needed for the present results. Sequential composition, general service regions, coefficient optimization, log-concavity, and random harmonic-cascade thresholds lie outside this paper.

## References

- [1] Alexei Ashikhmin, Gerhard Kramer, and Stephan ten Brink. Extrinsic information transfer functions: Model and erasure channel properties. *IEEE Transactions on Information Theory*, 50(11):2657–2673, 2004.
- [2] Hao Chen, San Ling, and Chaoping Xing. Asymptotically good quantum codes exceeding the Ashikhmin–Litsyn–Tsfasman bound. *IEEE Transactions on Information Theory*, 47(5):2055–2058, 2001.

- [3] Hao Chen, San Ling, and Chaoping Xing. Quantum codes from concatenated algebraic-geometric codes. *IEEE Transactions on Information Theory*, 51(8):2915–2920, 2005.
- [4] Charles J. Colbourn. *The Combinatorics of Network Reliability*. Oxford University Press, New York, 1987.
- [5] Alexander A. Davydov, Stefano Marcugini, and Fernanda Pambianco. Twisted cubic and orbits of lines in  $\text{PG}(3, q)$ , 2021. arXiv:2103.12655.
- [6] Johannes Gmainer and Hans Havlicek. Nuclei of normal rational curves, 2013.
- [7] Parikshit Gopalan, Cheng Huang, Huseyin Simitci, and Sergey Yekhanin. On the locality of codeword symbols. *IEEE Transactions on Information Theory*, 58(11):6925–6934, 2012.
- [8] Anina Gruica, Benjamin Jany, and Alberto Ravagnani. LRCs: Duality, LP bounds, and field size. *Designs, Codes and Cryptography*, 94:100, 2026.
- [9] Michel Lavrauw and Geertrui Van de Voorde. Locally repairable codes with high availability based on generalised quadrangles, 2019. arXiv:1912.06372.
- [10] Shu Liu, Liming Ma, Tingyi Wu, and Chaoping Xing. Good locally repairable codes via propagation rules, 2022. arXiv:2208.04484.
- [11] Florence Jessie MacWilliams and Neil James Alexander Sloane. *The Theory of Error-Correcting Codes*, volume 16 of *North-Holland Mathematical Library*. North-Holland, Amsterdam, 1977.
- [12] Arya Mazumdar. Capacity of locally recoverable codes. *IEEE Journal on Selected Areas in Information Theory*, 4:276–285, 2023.
- [13] Lluís Parnies-Juarez, Henk D. L. Hollmann, and Frédérique Oggier. Locally repairable codes with multiple repair alternatives. In *2013 IEEE International Symposium on Information Theory*, pages 892–896, 2013.
- [14] Tavis Rudd. Deep-hole rigidity of the Clebsch hexagon code. Working paper, 2026.
- [15] James Singer. A theorem in finite projective geometry and some applications to number theory. *Transactions of the American Mathematical Society*, 43(3):377–385, 1938.
- [16] Henning Stichtenoth. Transitive and self-dual codes attaining the tsfasman–vladut–zink bound. *IEEE Transactions on Information Theory*, 52(5):2218–2224, 2006.
- [17] Henning Stichtenoth. *Algebraic Function Fields and Codes*, volume 254 of *Graduate Texts in Mathematics*. Springer, 2 edition, 2009.
- [18] Paul Tricot. On 3-designs from  $\text{PGL}(2, q)$ , 2024.
- [19] Michel Las Vergnas. The Tutte polynomial of a morphism of matroids I: Set-pointed matroids and matroid perspectives. *Annales de l’Institut Fourier*, 49(3):973–1015, 1999.