

Quadratic recovery and cubic orientation in conic matching quotients

Tavis Rudd

Abstract

Restriction to a conic forgets how its marked points were paired into secants. For the balanced B_3 and H_3 configurations, we show that the quotient by the conic equation nevertheless recovers the two factorization sheets from quadratic products of affine evaluation functions. More generally, if a full $\mathrm{PGL}_2(q)$ -orbit has a unique quadratic trade whose two levels are one-factorizations, then edge counting forces a $q+q$ split, and subgroup classification leaves exactly these two orbits. Within signed tensor moments, the first nonzero term is an anti-invariant cubic line that detects sheet interchange. The resulting 14- and 22-point configurations are self-associated and arithmetically Gorenstein, with that signed cubic as the inverse system of an Artinian reduction. For the $A_3/\mathbb{F}_5$, $B_3/\mathbb{F}_7$, and $H_3/\mathbb{F}_{11}$ configurations, the difference spans have dimensions 3, 6, 10; in type H_3 , the only missing Fischer layer is QH_2 . The base-choice cocycle generates $H^1(\mathrm{PGL}_2(11), M_8)$, so the H_3 affine quotient has no equivariant origin.

Keywords. conic ideal; secant matching; Coxeter configuration; self-associated points; arithmetically Gorenstein; cubic inverse system; modular representation.

2020 Mathematics Subject Classification. 05B25, 51E22, 20C20, 13A02.

1 Introduction

The companion rigidity paper isolates the Clebsch hexagon through its uncovered syndrome locus and reconstructs its projective geometry from decoding data [7]. Here we begin with marked secant geometry and ask a complementary reconstruction question: what does common restriction to a conic forget, and what does the conic-ideal quotient remember?

The answer is governed by one mechanism. A four-endpoint switch shows that competing secant products differ by the conic equation, so their quotients form an affine configuration. In the balanced cases, Hadamard squares recover an unordered bipartition of that configuration; the first signed tensor that survives is cubic and orients the two parts. Thus quadratic data recover the factorization without choosing its orientation, and the first odd covariant restores that choice.

type	linear quotient	sheets/cubic
$A_3/\mathbb{F}_5$	rank 3	not applicable
$B_3/\mathbb{F}_7$	rank 6	recovered/oriented
$H_3/\mathbb{F}_{11}$	rank 10	recovered/oriented

The proof has two layers. Section 2 proves the quotient and switch identities symbolically for arbitrary matchings. Section 3 obtains all three ranks from the affine connecting cocycle and the defining-characteristic SL_2 -modules in the Fischer decomposition. In type B_3 , one transparent outer-sheet quotient supplies the radial line; type H_3 uses the analogous second-Laplacian scalar after cohomology excludes the middle layer. A stable two-level

quadratic-trade line gives the special-projective block system, and the one-factorization property forces both blocks to have size q . A maximal-subgroup argument then leaves only $B_3/\mathbb{F}_7$ and $H_3/\mathbb{F}_{11}$; the apparent $\mathbb{F}_5$ candidate acquires an outer normalizer and has only five points. In Section 4, the defining-characteristic permutation module proves the sheet restriction and moment ranks, and elementary duality forces the nonzero cubic and full Schur cube. Signed Gale duality then gives the Gorenstein conclusions. Appendix F maps each finite input to its primary check and independent replay.

Theorem 1.1 (Quadratic recovery and cubic orientation). *Let the A_3 , B_3 , and H_3 matching configurations be the finite matching orbits defined in Section 3. Then:*

- (i) *matching products scaled as in Section 2 have equal restriction to the conic, and their conic-ideal quotients have difference ranks 3, 6, and 10;*
- (ii) *among all odd prime powers q , the $B_3/\mathbb{F}_7$ and $H_3/\mathbb{F}_{11}$ configurations are the only full $\mathrm{PGL}_2(q)$ -orbits whose quadratic-trade line has two levels, each a one-factorization of $\mathbb{P}^1(\mathbb{F}_q)$;*
- (iii) *in the B_3 and H_3 cases, the two factorization sheets are, up to interchange, the unique complementary halves with equal first and second tensor moments;*
- (iv) *in those two cases, the signed moments vanish through degree two, while a nonzero degree-three tensor transforms by the character that exchanges the sheets;*
- (v) *the homogenized B_3 and H_3 configurations are self-associated with Gorenstein coordinate rings; the signed cubic is the Macaulay inverse system of their Artinian reductions.*

Appendix B gives a modular interpretation of the H_3 affine-origin obstruction and its homogeneous extension. Sections 2–4 prove the quotient, rank, and quadratic/cubic recovery results; Appendices A–E record H_3 refinements, and Appendix F documents verification.

2 Conic matching products

Write the standard conic as

$$\mathcal{C} = V(Q), \quad Q = XZ - Y^2,$$

and parametrize it by $\nu(s : t) = (s^2 : st : t^2)$. For points $a = (a_0 : a_1)$ and $b = (b_0 : b_1)$ of $\mathbb{P}^1$, put $[a, b] = a_0b_1 - a_1b_0$. We scale the secant L_{ab} through $\nu(a)$ and $\nu(b)$ by the condition

$$L_{ab}(\nu(s : t)) = [a, (s : t)] [b, (s : t)].$$

Let Ω be a set of $2m$ distinct marked points of $\mathbb{P}^1$, with fixed vector representatives. For a perfect matching M of Ω , define its matching product

$$P_M = \prod_{\{a,b\} \in M} L_{ab}.$$

The vector representatives are part of the marked-conic data. Their role is to make the products comparable before passing to projective equivalence.

Proposition 2.1 (The matching-secant quotient). *For every $m \geq 2$ and every two perfect matchings M, N of Ω ,*

$$P_M|_{\mathcal{C}} = P_N|_{\mathcal{C}}, \quad P_M - P_N \in (Q).$$

Consequently, after choosing a reference matching M_0 ,

$$\Phi_{M_0}(M) = \frac{P_M - P_{M_0}}{Q} \in H^0(\mathbb{P}^2, \mathcal{O}(m-2)) \quad (2.1)$$

is an affine configuration. Changing the reference matching translates this configuration. If $g \in \mathrm{GL}_2$ preserves Ω , $\rho(g)$ is its action on binary quadratics, and representatives are transported by g , then

$$\Phi_{gM_0}(gM) \circ \rho(g) = \det(g)^{2m-2} \Phi_{M_0}(M). \quad (2.2)$$

Thus the difference space, its rank, and its affine moment conditions do not depend on M_0 and are equivariant under the endpoint stabilizer.

Proof. For $x \in \mathbb{P}^1$,

$$P_M(\nu(x)) = \prod_{a \in \Omega} [a, x],$$

which does not depend on the pairing. The kernel of the Veronese pullback

$$k[X, Y, Z] \longrightarrow k[s, t], \quad (X, Y, Z) \longmapsto (s^2, st, t^2)$$

is the principal ideal (Q) , so Q divides every difference. Changing the reference follows by subtraction. Finally, $L_{ga,gb} \circ \rho(g) = \det(g)^2 L_{ab}$ and $Q \circ \rho(g) = \det(g)^2 Q$, which gives (2.2). $\square$

2.1 The four-endpoint switch

For four distinct endpoints, the Plücker relation gives the local switch identity

$$L_{ab}L_{cd} - L_{ac}L_{bd} = [a, d][b, c] Q. \quad (2.3)$$

Multiplying (2.3) by the secant factors common to two matchings proves divisibility for a single switch. Any two perfect matchings are joined by a sequence of such switches: if M and N differ at a , write $\{a, b\} \in M$ and $\{a, c\} \in N$; switching the two M -edges through b and c creates $\{a, c\}$ and reduces the symmetric difference. This also gives an elementary proof of the divisibility clause in Proposition 2.1.

The switch calculation and the global pullback proof play different roles. The pullback proves the quotient for arbitrary matchings at once. The switch identity records how local changes of factorization move in the quotient space. A selected Coxeter orbit need not contain every perfect matching or every intermediate switch.

3 The rank-three configurations

We now select three finite matching orbits. This selection is discrete input: Proposition 2.1 applies to every matching, but it does not distinguish the Coxeter orbits. Here and below, “rank three” refers to Coxeter rank. It does not refer to the dimensions of the conic-ideal quotient spaces, which are 3, 6, 10.

For $q \in \{5, 7, 11\}$, index

$$\mathbb{P}^1(\mathbb{F}_q) = \{0, 1, \dots, q-1, \infty\},$$

where a denotes $(1 : a)$ and ∞ denotes $(0 : 1)$. Let $G_q = \mathrm{PGL}_2(q)$ act in the usual way. The three base matchings are

T	q	M_T	
A_3	5	$\{0, \infty\}, \{1, 4\}, \{2, 3\},$	(3.1)
B_3	7	$\{0, 2\}, \{1, 4\}, \{3, \infty\}, \{5, 6\},$	
H_3	11	$\{0, 1\}, \{2, 5\}, \{3, 7\}, \{4, 9\}, \{6, 8\}, \{10, \infty\}.$	

Define the *rank-three matching configuration*

$$\Omega_T = G_q \cdot M_T.$$

The stabilizers of the three displayed matchings are respectively S_4, S_4, A_5 , and hence

$$|\Omega_{A_3}| = 5, \quad |\Omega_{B_3}| = 14, \quad |\Omega_{H_3}| = 22. \quad (3.2)$$

These marker spaces and their stabilizers are classical. In particular, Edge and Dye identify the exceptional finite conic configurations and substantial parts of their incidence geometry [2, 3]. We use the Coxeter names for the corresponding tetrahedral, octahedral, and icosahedral data; neither the orbit counts in (3.2) nor the ambiguity of an unmarked conic is asserted here as new.

Theorem 3.1 (Intrinsic completeness of recovered factorization sheets). *Let q be an odd prime power, put*

$$G = \mathrm{PGL}_2(q), \quad G^+ = \mathrm{PSL}_2(q),$$

and let Ω be a G -orbit of perfect matchings of $\mathbb{P}^1(\mathbb{F}_q)$. Form its conic-ideal quotient configuration and let $L \subseteq \mathbb{F}_q^\Omega$ be its affine evaluation space. Suppose $(L^{\circ 2})^\perp$ is a line whose nonzero vectors have exactly two level sets, and that each level set is a one-factorization: its matchings partition the edges of the complete graph on $\mathbb{P}^1(\mathbb{F}_q)$. Then

$$(q, \Omega) = (7, \Omega_{B_3}) \quad \text{or} \quad (q, \Omega) = (11, \Omega_{H_3}). \quad (3.2a)$$

Conversely, both displayed orbits satisfy these intrinsic hypotheses.

Proof. Proposition 2.1 makes L , its square, and the line

$$R = (L^{\circ 2})^\perp$$

G -stable. Choose $0 \neq \tau \in R$. Its zero set is G -invariant, so transitivity gives τ full support. The two level sets of τ depend only on R , hence G permutes them. This action is nontrivial because G is transitive on Ω .

For $q > 3$, the commutator subgroup of G is G^+ , and G/G^+ has order two. The kernel of the action on the two levels is therefore G^+ . The stabilizer of a point fixes its level and so lies in G^+ ; consequently the two levels are exactly the two G^+ -orbits. An outer element acts on R by a scalar and exchanges the two nonzero values of τ , so those values are opposites.

Each one-factorization of the complete graph on $q + 1$ vertices has

$$\frac{\binom{q+1}{2}}{(q+1)/2} = q$$

perfect matchings. Thus the intrinsic levels have size q , and $|\Omega| = 2q$. When $q = 3$, there are only three perfect matchings altogether, so two disjoint one-factorizations cannot occur.

Now fix $M \in \Omega$ and let $K = \text{Stab}_G(M)$. Orbit-stabilizer and the derived two-sheet conclusion give

$$|K| = \frac{q^2 - 1}{2}, \quad K \leq G^+, \quad [G^+ : K] = q. \quad (3.2b)$$

Assume $q \geq 5$, and place K in a maximal subgroup of G^+ . Dickson's maximal-subgroup theorem, in the form recorded by Giudici [4, Theorem 2.2], leaves Borel, dihedral, subfield, and exceptional A_4, S_4, A_5 subgroups. The first two have order smaller than $(q^2 - 1)/2$. A proper subfield subgroup is no larger than $\text{PGL}_2(r)$, where $q = r^e$ and $e \geq 2$, and

$$r(r^2 - 1) < \frac{r^{2e} - 1}{2}.$$

Thus it too is too small. An exceptional maximal subgroup has order at most 60, so (3.2b) leaves $q = 5, 7, 9, 11$. For $q = 9$, the required order 40 divides none of 12, 24, 60. In the other three cases, order and divisibility force

$$(q, K) = (5, A_4), (7, S_4), (11, A_5). \quad (3.2c)$$

It remains to ask which of these subgroup actions preserves a matching. This is a block-system question, not a matching census. For $q = 5$, the A_4 -action on the six endpoints is transitive with point stabilizer C_2 . Its normal Klein four subgroup gives the unique two-point block system. The normalizer of A_4 in $\text{PGL}_2(5)$ is S_4 ; uniqueness makes this whole normalizer preserve the matching. Its full orbit therefore has five points, namely Ω_{A_3} , rather than ten. In particular, the other ten matchings cannot form a balanced $5 + 5$ orbit.

For $q = 7$, the S_4 -action on eight endpoints has point stabilizer C_3 . A two-point block stabilizer must have order six, and the unique possibility containing that C_3 is its normalizer S_3 . Hence the invariant matching is unique. Similarly, for $q = 11$, the A_5 -action on twelve endpoints has point stabilizer C_5 ; its unique possible two-point block stabilizer is $N_{A_5}(C_5) = D_{10}$. The invariant matching is again unique. In each case the two G^+ -classes of the exceptional subgroup are fused by G , and the subgroup has no outer normalizer [4]. Its full matching stabilizer is therefore exactly S_4 or A_5 , so the orbit has size $2q$ and splits into the two claimed G^+ -orbits. The unique block systems are precisely the matchings displayed in (3.1), up to G -conjugacy.

Conversely, $q = 7, 11$ are both $3 \pmod{4}$, so G^+ is transitive on unordered pairs of endpoints. Each q -matching sheet therefore covers every edge equally often; its total of $q(q+1)/2$ edge occurrences equals the number of edges, so it is a one-factorization. Propositions 4.1 and 4.2 identify $(L^{\circ 2})^\perp$ with the sheet-sign line. Hence both displayed orbits satisfy the intrinsic hypotheses. $\square$

Fix M_T as reference and apply (2.1) to every member of Ω_T . Write $R_d = \mathbb{F}_q[X, Y, Z]_d$ and

$$\Delta_Q = 4\partial_X\partial_Z - \partial_Y^2, \quad \mathcal{H}_d = \ker(\Delta_Q : R_d \longrightarrow R_{d-2}). \quad (3.3)$$

The operator Δ_Q is the Laplacian attached to the dual quadratic form. In the degrees used below, $\mathcal{H}_d$ is the top harmonic summand, while powers of Q give radial summands.

Theorem 3.2 (The 3, 6, 10 quotient ranks). *For the matching configurations in (3.1), let*

$$W_T = \text{span}_{\mathbb{F}_q} \{ \Phi_{M_T}(M) : M \in \Omega_T \}.$$

Then

T	$\deg \Phi$	W_T	$\dim W_T$
A_3	1	$R_1 = \mathcal{H}_1$	3,
B_3	2	$R_2 = \mathcal{H}_2 \oplus \mathbb{F}_7 Q$	6,
H_3	4	$\mathcal{H}_4 \oplus \mathbb{F}_{11} Q^2$	10.

(3.4)

If $h_T = 4, 6, 10$ is the Coxeter number and $d = h_T/2 - 1$, the three rows have the uniform form

$$W_T = \begin{cases} \mathcal{H}_d, & d \text{ odd}, \\ \mathcal{H}_d \oplus \mathbb{F}_q Q^{d/2}, & d \text{ even}, \end{cases} \quad \dim W_T = h_T - 1 + \mathbf{1}_{\{d \text{ even}\}}.$$

In particular, the first two configurations fill the complete quotient form space. The H_3 configuration spans a canonical ten-dimensional subspace of the fifteen-dimensional space R_4 .

Proof mode. The quotient construction and all three spanning assertions are proved symbolically. Irreducibility forces the A_3 image and the top B_3 summand; one displayed outer-sheet quotient supplies the B_3 radial line. For H_3 , a Sylow-normalizer cohomology argument excludes the middle Fischer layer and forces the top harmonic summand; one displayed outer-sheet quotient supplies the remaining radial line. A separate implementation reconstructs the three full quotient matrices and the two smaller radial witnesses from the data in (3.1).

Proof. A matching in (3.1) has $m = 3, 4, 6$ edges, respectively, so Proposition 2.1 places its quotient in R_{m-2} , of degrees 1, 2, 4. Direct differentiation gives

$$\dim \mathcal{H}_1 = 3, \quad \dim \mathcal{H}_2 = 5, \quad \dim \mathcal{H}_4 = 9.$$

The radial vector $Q \in R_2$ is not harmonic in characteristic 7.

We first prove the A_3 and B_3 rows without quotient matrices. For $q = 5, 7$, put

$$G = \mathrm{PGL}_2(q), \quad G^+ = \mathrm{PSL}_2(q), \quad \chi(g) = \det(g)^{(q-1)/2}.$$

The Dickson form $s^q t - s t^q$, which is the common conic restriction of the matching products, has contragredient weight $\det^{-1}$. The Clebsch identification and Fischer decomposition therefore give

$$\begin{aligned} q = 5 & \quad \tilde{R}_1 = R_1 \otimes \det^{-1} = M_2, \\ q = 7 & \quad \tilde{R}_2 = R_2 \otimes \det^{-1} = M_4 \oplus M_0, \end{aligned} \tag{3.4a}$$

where, uniformly,

$$M_r = \mathrm{Sym}^r(V^\vee) \otimes \det^{r/2} \otimes \chi, \quad M_0 = \chi.$$

The scalar action is trivial, so these are G -modules. On G^+ , M_2 and M_4 are the absolutely irreducible symmetric powers of highest weights $2 < 5$ and $4 < 7$, respectively [8, Fact 3.1].

For either type let

$$c(g) = \Phi_{M_T}(g M_T).$$

It is an affine cocycle, $c(gh) = gc(h) + c(g)$, and its value span is G -stable because

$$b c(g) = c(bg) - c(b). \tag{3.4b}$$

Distinct matchings have distinct products: unique factorization recovers the normalized secant factors. Hence every nonbase orbit point gives a nonzero cocycle value. In type A_3 , irreducibility of M_2 now forces the value span to be all of R_1 .

In type B_3 , project the cocycle to $M_0 = \chi$. Its restriction to G^+ is a homomorphism from the simple group $\mathrm{PSL}_2(7)$ to the additive group of $\mathbb{F}_7$, and is therefore zero. A nonbase point on the G^+ -sheet consequently gives a nonzero vector in M_4 ; irreducibility and (3.4b) force the whole five-space $M_4 = \mathcal{H}_2$.

It remains to supply the radial line. The nonsquare-determinant matrix

$$u = \begin{pmatrix} 0 & 1 \\ 1 & 2 \end{pmatrix}$$

sends the base matching to

$$M_* = \{\{0, 2\}, \{1, 5\}, \{3, 4\}, \{6, \infty\}\}.$$

Using $L_{ab} = abX - (a+b)Y + Z$ for finite a, b , direct cancellation in the two products gives the two-line quotient

$$\Phi_{M_{B_3}}(M_*) = (Z - 2Y)(3Z + Y) = 3L_{02}^2. \quad (3.4c)$$

Thus the outer quotient is the square of the secant common to the two matchings, up to a nonzero scalar. The dual quadratic norm of a secant line is nonzero, whereas it vanishes for a tangent line. Equivalently, in the displayed normalization,

$$\Delta_Q(3L_{02}^2) = 4 \neq 0 \quad \text{in } \mathbb{F}_7. \quad (3.4d)$$

Since Δ_Q kills $M_4 = \mathcal{H}_2$, this outer-sheet value has nonzero $M_0 = \mathbb{F}_7 Q$ -projection. The span already contains M_4 , so it contains M_0 as well. Hence $W_{B_3} = R_2$, proving the first two rows of (3.4).

We prove the H_3 row equivariantly. Put $G = \mathrm{PGL}_2(11)$, $G^+ = \mathrm{PSL}_2(11)$, and let $\chi(g) = \det(g)^5$ be the square-class character. On the conic, the common restriction of the matching products is the Dickson form $s^{11}t - st^{11}$, of contragredient weight $\det^{-1}$. Let $V = \mathbb{F}_{11}^2$, with binary forms carrying the contragredient $\mathrm{GL}_2(11)$ -action. The equivariant Fischer decomposition obtained from the Clebsch identification $\mathrm{Sym}^2 V \simeq \mathbb{F}_{11}^3$, together with the Dickson weight in the quotient transformation (2.2), gives the linearized module

$$\tilde{R}_4 = R_4 \otimes \det^{-1} = M_8 \oplus M_4 \oplus M_0, \quad (3.5)$$

Explicitly, $g \in \mathrm{GL}_2(11)$ acts on the first expression by

$$g \cdot f = \det(g)^{-1} (f \circ \rho(g)^{-1}).$$

The three summands are

$$\begin{aligned} M_8 &= \mathcal{H}_4 \otimes \det^{-1} \simeq \mathrm{Sym}^8(V^\vee) \otimes \det^{-1}, \\ M_4 &= Q\mathcal{H}_2 \otimes \det^{-1} \simeq \mathrm{Sym}^4(V^\vee) \otimes \det^{-3}, \\ M_0 &= \mathbb{F}_{11} Q^2 \otimes \det^{-1} \simeq \det^{-5} = \chi. \end{aligned} \quad (3.6)$$

Equivalently, all three summands have the uniform form

$$M_r = \mathrm{Sym}^r(V^\vee) \otimes \det^{r/2} \otimes \chi, \quad r = 8, 4, 0.$$

A scalar λI acts here as $\lambda^{-r} \lambda^r (\lambda^2)^5 = 1$, so these modules and their direct sum descend from $\mathrm{GL}_2(11)$ to G . This is the G -action used below. The determinant twist changes only the linearization, not the underlying Fischer subspaces $\mathcal{H}_4, Q\mathcal{H}_2, \mathbb{F}_{11} Q^2$ of R_4 .

We need two elementary cohomology facts. Let $U = \langle u \rangle$ be the upper-unipotent Sylow 11-subgroup and let a generate the diagonal torus of order 10. Since $r < 10$, the cyclic norm $1 + u + \cdots + u^{10} = (u - 1)^{10}$ vanishes on M_r , while $(u - 1)M_r$ has codimension one. Thus $H^1(U, M_r)$ is represented by the class of x^r . Diagonal conjugation acts on that class by

$$a^{4-r/2}. \quad (3.7)$$

Restriction from G to U is injective because $[G : U] = 120$ is invertible in $\mathbb{F}_{11}$, and its image is fixed by the Sylow normalizer. Equation (3.7) therefore gives

$$H^1(G, M_4) = H^1(G, M_0) = 0, \quad \dim H^1(G, M_8) \leq 1. \quad (3.8)$$

The stabilizer $H = A_5$ of M_{H_3} lies in G^+ . The restrictions of M_8, M_4, M_0 to H are respectively $4 \oplus 5, 5, 1$, so

$$M_8^H = M_4^H = 0, \quad \dim M_0^H = 1. \quad (3.9)$$

Equivalently, average the characters $(9, 1, 0, -1, -1)$, $(5, 1, -1, 0, 0)$, and $(1, 1, 1, 1, 1)$ over the A_5 class sizes $(1, 15, 20, 12, 12)$.

Let $c(g) = \Phi_{M_{H_3}}(gM_{H_3})$. The common-restriction identity gives

$$c(gh) = gc(h) + c(g),$$

and c vanishes on H . Project c through (3.5). By (3.8), its M_4 -component is a coboundary; its translating vector lies in $M_4^H = 0$, so that component vanishes. This proves, without quotient coordinates,

$$\Delta_Q \Phi_{M_{H_3}}(M) \in \mathbb{F}_{11}Q \quad \text{for every } M \in \Omega_{H_3}. \quad (3.10)$$

Likewise the M_0 -component is a coboundary and hence vanishes on G^+ . Choose any other matching in the eleven-element G^+ -orbit of M_{H_3} . Its quotient is nonzero: equality of the two secant products would identify their irreducible line factors and hence the matchings. Its M_4 - and M_0 -components vanish, so it gives a nonzero vector in M_8 . The degree-eight restricted $\mathrm{SL}_2(\mathbb{F}_{11})$ -module is irreducible [8, Fact 3.1]. Moreover the span of the cocycle values is G -stable, since

$$bc(g) = c(bg) - c(b) \quad (b, g \in G).$$

Therefore the quotient span contains all of M_8 .

It remains only to see the radial line. With ∞ indexed by 11, take

$$M_- = \{\{0, 1\}, \{2, \infty\}, \{3, 8\}, \{4, 6\}, \{5, 9\}, \{7, 10\}\}.$$

The nonsquare-determinant projectivity represented on V by

$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$$

sends M_{H_3} to M_- ; hence M_- belongs to the outer sheet of Ω_{H_3} . The primary certificate and independent replay also reconstruct this orbit membership. Exact division and differentiation give

$$\Delta_Q^2 \Phi_{M_{H_3}}(M_-) = 10 = \Delta_Q^2(Q^2) \neq 0. \quad (3.11)$$

The independent replay reconstructs this single scalar from the secant formula. Since Δ_Q^2 kills $M_8 \oplus M_4$, (3.11) supplies the M_0 -projection. The span already contains M_8 , so it also contains M_0 . Hence $W_{H_3} = \mathcal{H}_4 \oplus \mathbb{F}_{11}Q^2$, of dimension 10. Since $\dim R_4 = 15$, this equality is proper. $\square$

Corollary 3.3 (The base-choice class and its homogeneous extension). *Let $c_8 : G \rightarrow M_8$ be the top-harmonic projection of the base-choice cocycle c . Then*

$$H^1(G, M_8) = \mathbb{F}_{11}[c_8]. \quad (3.11a)$$

In particular, the affine H_3 quotient configuration has no G -equivariant origin: no translation makes its affine action linear. Equivalently, the homogeneous module

$$E_{c_8} = M_8 \oplus \mathbb{F}_{11}, \quad g \cdot (v, t) = (gv + t c_8(g), t), \quad (3.11b)$$

is nonsplit. Up to equivalence and nonzero rescaling of the quotient coordinate, it is the unique nonsplit extension of the trivial G -module by M_8 .

Proof. Equation (3.8) gives $\dim H^1(G, M_8) \leq 1$. On the base sheet the M_0 -component vanishes, and the M_4 -component vanishes by (3.9); the nonzero same-sheet quotient used in the proof above is therefore a value of c_8 . If $c_8(g) = gv - v$, its vanishing on $H = \text{Stab}_G(M_{H_3})$ would put v in $M_8^H = 0$, forcing $c_8 = 0$, a contradiction. Thus $[c_8] \neq 0$, proving (3.11a).

Changing the reference quotient translates the affine configuration and changes c_8 by a coboundary. Hence a G -equivariant origin would kill $[c_8]$, which is impossible. Finally the standard identification $\text{Ext}_{\mathbb{F}_{11}G}^1(\mathbb{F}_{11}, M_8) \simeq H^1(G, M_8)$ identifies (3.11b) with $[c_8]$; the one-dimensionality just proved gives the last assertion. $\square$

Corollary 3.4 (The exact H_3 loss). *Over $\mathbb{F}_{11}$, the quartic form space has the Fischer decomposition*

$$R_4 = \mathcal{H}_4 \oplus Q\mathcal{H}_2 \oplus \mathbb{F}_{11}Q^2. \quad (3.12)$$

Consequently,

$$R_4 = W_{H_3} \oplus Q\mathcal{H}_2, \quad R_4/W_{H_3} \simeq Q\mathcal{H}_2. \quad (3.13)$$

Equivalently,

$$W_{H_3} = \{f \in R_4 : \Delta_Q f \in \mathbb{F}_{11}Q\}. \quad (3.14)$$

Thus the H_3 quotient configuration retains the top harmonic nine-space and the deepest radial line, and omits precisely the five-dimensional middle radial layer.

Proof. For a homogeneous form f of degree r , direct differentiation gives

$$\Delta_Q(Qf) = (4r + 6)f + Q\Delta_Q f. \quad (3.15)$$

Hence $\Delta_Q(Qh) = 3h$ for $h \in \mathcal{H}_2$, while $\Delta_Q(Q^2) = 9Q$ over $\mathbb{F}_{11}$. Applying Δ_Q^2 and then Δ_Q shows that the three summands in (3.12) meet trivially. Their dimensions are 9, 5, 1, so they exhaust the fifteen-dimensional space R_4 . Theorem 3.2 then gives (3.13). If $f = h_4 + Qh_2 + cQ^2$ is its decomposition, then

$$\Delta_Q f = 3h_2 + 9cQ.$$

The quadratic decomposition $R_2 = \mathcal{H}_2 \oplus \mathbb{F}_{11}Q$ therefore gives (3.14). $\square$

The rank proof has two distinct boundaries. The common restriction and quotient are general consequences of the conic ideal, and (3.3) describes the intrinsic target subspaces. For A_3 , irreducibility alone forces the image. For B_3 , irreducibility forces the top summand and the two-line calculation (3.4c)–(3.4d) detects the radial line. For H_3 , (3.7)–(3.10) prove the containment and top-summand nonvanishing equivariantly; only the single radial scalar (3.11) remains an exact coordinate witness. The primary certificate still enumerates every

orbit and recovers the ranks, while an independent replay reconstructs both that calculation and the smaller witnesses from (3.1).

This proof is uniform at the level of its mechanism—an affine connecting cocycle, its Sylow-normalizer weights, stabilizer fixed spaces, and irreducible summands—but it still uses the H_3 -specific $A_5 < \mathrm{PGL}_2(11)$ module data and separate geometric nonvanishing witnesses for the two radial lines. A theorem over all good reductions would additionally require integral models and an analysis of the characteristics in which the harmonic decomposition, orbit structure, or marker algebra degenerates.

Corollary 3.4 identifies the linear information lost by the H_3 orbit, but linear span does not recover the two factorization sheets inside the finite configuration. Their recovery begins one categorical level higher, with products of affine evaluation functions.

4 Balanced sheets and cubic orientation

For $T = B_3, H_3$, put $q = 7, 11$, respectively. The subgroup

$$G^+ = \mathrm{PSL}_2(q) \subset G = \mathrm{PGL}_2(q)$$

splits Ω_T into two orbits $\Omega_+ \sqcup \Omega_-$, each of size q ; an element of the outer coset exchanges them. These are the two one-factorization sheets. The A_3 orbit does not split: its S_4 stabilizer cannot lie in $\mathrm{PSL}_2(5) \simeq A_5$, already by order.

Write $x_M = \Phi_{M_T}(M) \in W_T$. Let $L \subseteq \mathbb{F}_q^{\Omega_T}$ be the evaluation space of affine-linear functions on the finite configuration $\{x_M\}$. Since the points affinely span W_T ,

$$\dim L = 1 + \dim W_T = q.$$

For subspaces of functions, write

$$L^{\circ 2} = \mathrm{span}\{fg : f, g \in L\},$$

where multiplication is pointwise. The common second-moment form in the proposition below is

$$B(\lambda, \mu) = \sum_{M \in \Omega_+} \lambda(x_M) \mu(x_M) = \sum_{M \in \Omega_-} \lambda(x_M) \mu(x_M)$$

on $W_T^\vee$. A *field-valued strength-two trade* is a weight $\tau \in k^\Omega$ satisfying, for all $f, g \in L$,

$$\sum_{M \in \Omega} \tau(M) f(M) g(M) = 0.$$

The B_3 model. Here the quotient configuration consists of fourteen points spanning the six-space W_{B_3} , split into two seven-point sheets. Its affine evaluation space L has dimension seven. Restriction to either sheet has rank six, while the common second-moment form has rank five and a radical functional that takes distinct constant values on the two sheets. It follows that $L^{\circ 2}$ is the thirteen-dimensional hyperplane of functions with equal sheet sums; its one-dimensional annihilator is the sheet sign. Thus the quadratic algebra recovers the 7+7 partition without a search through partitions. The first nonzero signed moment is cubic, so $L^{\circ 3} = \mathbb{F}_7^{14}$, and the homogenized points have h -vector $(1, 6, 6, 1)$. The next proposition isolates the quadratic step in a form that applies to both balanced configurations.

Proposition 4.1 (Radical–Hadamard recovery). *Let k be a field and $\Omega = \Omega_+ \sqcup \Omega_-$, with both sheets of size $q = 0$ in k . Suppose an affine evaluation space $L \subseteq k^\Omega$ has the following properties:*

- (i) each sheet has zero first moment, and the two sheets have equal second moments;
- (ii) $\dim L = q$, and restriction from L onto the zero-sum hyperplane of either sheet has rank $q - 1$;
- (iii) the second-moment form on linear functionals has rank $q - 2$ and a one-dimensional radical;
- (iv) a nonzero radical functional is constant on each sheet, with different constants on the two sheets.

Then

$$L^{\circ 2} = \left\{ (u_+, u_-) : \sum_{\Omega_+} u_+ = \sum_{\Omega_-} u_- \right\}, \quad (4.1)$$

and

$$(L^{\circ 2})^\perp = k(\mathbf{1}_{\Omega_+} - \mathbf{1}_{\Omega_-}). \quad (4.2)$$

Consequently every field-valued strength-two trade is a scalar multiple of the sheet sign.

Proof. Let e_+ and e_- be the two sheet indicators. The radical functional evaluates as $a_+e_+ + a_-e_-$ with $a_+ \neq a_-$. Together with $1 = e_+ + e_- \in L$, it therefore yields $e_+, e_- \in L$.

Let $H_\pm$ be the zero-sum hyperplane on $\Omega_\pm$. The first-moment hypothesis and $q = 0$ put each restriction of L inside $H_\pm$; the rank hypothesis makes both restrictions surjective. Multiplication by e_+ and e_- now gives

$$(H_+ \oplus 0) + (0 \oplus H_-) \subseteq L^{\circ 2}. \quad (4.3)$$

This subspace has dimension $2q - 2$.

For $f, g \in L$, the difference between the two sheet sums of fg expands into constant, first-moment, and second-moment terms. The hypotheses make all three differences zero, so $L^{\circ 2}$ lies in the hyperplane on the right of (4.1). Finally choose restrictions $a, b \in H_+$ with nonzero standard pairing and lift them to $f, g \in L$. Equality of the second moments gives the same nonzero pairing on Ω_- . Thus fg has equal nonzero sheet sums and does not lie in (4.3). We have found $2q - 1$ independent directions in the $2q - 1$ -dimensional hyperplane, proving (4.1). Its annihilator under the standard coordinate pairing is the line in (4.2). $\square$

Proposition 4.2 (The modular-permutation mechanism). *For $T = B_3, H_3$, the affine quotient configuration satisfies all four hypotheses of Proposition 4.1. More precisely, write*

$$W_T = U_T \oplus \mathbb{F}_q r, \quad U_{B_3} = \mathcal{H}_2, \quad U_{H_3} = \mathcal{H}_4.$$

On either q -point sheet, affine evaluation has image the zero-sum hyperplane and rank $q - 1$. The common second-moment form has rank $q - 2$, with radical the radical functional $r^\vee$; that functional is constant on each sheet and its two constants are distinct.

Proof. As a G^+ -module, U_T is the irreducible module M_{q-3} of dimension $q - 2$. Project the affine quotient points to U_T , writing the resulting sheet as $\{y_\omega\}$. The cocycle argument in Section 3 says that the differences $y_\omega - y_{\omega'}$ span U_T . Moreover $\sum_\omega y_\omega$ is G^+ -fixed: translating the sheet adds q times the cocycle value, which vanishes in characteristic q . Since $U_T^{G^+} = 0$,

$$\sum_\omega y_\omega = 0.$$

Let $P = \mathbb{F}_q^{\Omega+}$, and put

$$P_0 = \left\{ (a_\omega) : \sum_\omega a_\omega = 0 \right\}, \quad S = \mathbb{F}_q \mathbf{1}.$$

Here $S \subset P_0$, the defining-characteristic feature that drives the argument. The map

$$P_0 \longrightarrow U_T, \quad (a_\omega) \longmapsto \sum_\omega a_\omega y_\omega$$

is onto because the differences span, and the preceding moment identity kills S . Both P_0/S and U_T have dimension $q - 2$, so it induces an isomorphism

$$P_0/S \simeq U_T.$$

The standard dot product on P_0 has radical exactly S . Dualizing therefore shows that the evaluations of $U_T^\vee$ form a nondegenerate $(q - 2)$ -space complementary to S in P_0 . Adding constants gives all of P_0 , proving the restriction rank and the top second-moment rank. The same argument applies to the other sheet.

The two top second moments are not merely proportional. On

$$M_{q-3} = \text{Sym}^{q-3}(V^\vee) \otimes \det^{(q-3)/2} \otimes \chi$$

the standard apolar pairing is G -invariant: its determinant weight $\det^{-(q-3)}$ is cancelled by the square of the displayed twist, and $\chi^2 = 1$. Each sheet moment is a nonzero G^+ -invariant form, hence a scalar multiple of this pairing, while an outer element transports one sheet to the other by an apolar isometry. Their scalars, and thus the two forms, are equal.

Finally, projection of the affine cocycle to the radial character χ restricts on the simple group G^+ to an additive homomorphism, hence vanishes. The radial coordinate is consequently zero on the base sheet and constant on the other. The secant-norm witness (3.4c)–(3.4d) for B_3 , and the second-Laplacian witness (3.11) for H_3 , show that the outer constant is nonzero. Because $q = 0$, radial–radial and radial–top second moments vanish. Thus $r^\vee$ is exactly the one-dimensional radical, and its sheet constants are distinct. $\square$

For reference, exact reconstruction of the two configurations gives

T	q	sheet restriction ranks	second-moment rank	$\dim L^{\circ 2}$	
B_3	7	6, 6	5	13,	
H_3	11	10, 10	9	21.	(4.4)

In each row the radical is one-dimensional and takes two distinct sheet values. Proposition 4.2, rather than this table, proves these facts. The exact certificate and its independent reconstruction are cross-checks only; no enumeration of equal-size subsets or row reduction is used in the proof.

Lemma 4.3 (A full-support quadratic relation forces the cube). *Let Ω be finite and let $L \subseteq k^\Omega$ be a unital linear space with $\dim L > 1$. Suppose*

$$L^{\circ 2} = \left\{ u : \sum_{\omega \in \Omega} \epsilon_\omega u_\omega = 0 \right\}$$

for some $\epsilon \in (k^\times)^\Omega$. Then

$$L^{\circ 3} = k^\Omega.$$

Proof. If a nonzero vector η annihilated $L^{\circ 3}$, then $L^{\circ 2} \subseteq L^{\circ 3}$ would put η in $k\epsilon$. For every $f \in L$, the vector $\eta \circ f$ also annihilates $L^{\circ 2}$, because

$$\langle \eta \circ f, h \rangle = \langle \eta, fh \rangle = 0 \quad (h \in L^{\circ 2}).$$

Thus $\eta \circ f \in k\epsilon = k\eta$. Since η has full support, every $f \in L$ is constant, contrary to $\dim L > 1$. The annihilator of $L^{\circ 3}$ is therefore zero. $\square$

Theorem 4.4 (Balanced sheets and cubic-first orientation). *For $T = B_3, H_3$, the unordered pair $\{\Omega_+, \Omega_-\}$ is intrinsic to the affine quotient configuration:*

- (i) *up to interchange, Ω_+ and Ω_- are the only complementary halves with equal first and second tensor moments;*
- (ii) *if ϵ is +1 on Ω_+ and -1 on Ω_- , then*

$$\mu_j = \sum_{M \in \Omega_T} \epsilon(M) x_M^{\odot j} \in \text{Sym}^j(W_T)$$

satisfies

$$\mu_1 = 0, \quad \mu_2 = 0, \quad \mu_3 \neq 0; \tag{4.5}$$

- (iii) *μ_3 is independent of the reference matching, G^+ fixes it, and the outer coset negates it. Within G ,*

$$\text{Stab}_G(\mu_3) = G^+, \quad \text{Stab}_G(\mathbb{F}_q \mu_3) = G. \tag{4.6}$$

Thus degree three is the first signed tensor moment, or linear functional of such a moment, that orients the recovered sheet pair. This minimality statement concerns signed tensor moments, not every possible nonlinear statistic of the configuration.

Proof mode. Proposition 4.1 is a symbolic proof. Proposition 4.2 proves its hypotheses from the defining-characteristic permutation module and the two geometric radial witnesses. Lemma 4.3 forces the nonzero cubic without a tensor calculation or Gorenstein premise. Signed Gale duality and the self-association criterion then give the Gorenstein geometry. Exact finite arithmetic independently cross-checks the displayed ranks, and the Hilbert-symmetry arithmetic is kernel-checked in Lean.

Proof. Equality of first and second tensor moments for a complementary pair is equivalent, by polarization, to its sign weight annihilating $L^{\circ 2}$. Since the characteristics are 7 and 11, polarization through degree two is valid. Proposition 4.1 says that the annihilator is exactly the sheet-sign line. A sign weight taking only the values $\{\pm 1\}$ is therefore ϵ or $-\epsilon$, proving (i).

The same moment equalities give $\mu_1 = \mu_2 = 0$. Proposition 4.1 identifies $L^{\circ 2} = \ker \langle \epsilon, - \rangle$, and ϵ has full support. Lemma 4.3 gives $L^{\circ 3} = \mathbb{F}_q^{\Omega_T}$. Hence ϵ cannot annihilate all cubic products. Since the lower signed moments vanish and 6 is invertible in $\mathbb{F}_7, \mathbb{F}_{11}$, polarization gives $\mu_3 \neq 0$.

For the geometric consequence, let A be the q -by- $2q$ matrix whose M -th column is $\hat{x}_M = (1, x_M)$, and put $D = \text{diag}(\epsilon(M))$. Equal sheet sizes and the vanishing signed moments through degree two give

$$ADA^\top = 0.$$

The affine spanning statement gives $\text{rank } A = q$, so the rows of AD form the kernel of A . Thus AD is a Gale matrix, and its columns differ from those of A only by the nonzero scalars $\epsilon(M)$. The homogenized configuration is self-associated.

Proposition 4.1 also says that the $2q$ points fail by exactly one to impose independent conditions on quadrics. The unique dependence is ϵ , which has full support, so every one-point deletion imposes independent conditions. Eisenbud–Popescu’s self-association criterion [5, Theorem 7.3] therefore makes the coordinate ring arithmetically Gorenstein.

Reduce by the homogenizing coordinate, which is regular because it is nonzero at every point. The first three values of the h -vector are

$$1, \quad q-1, \quad q-1,$$

from $\dim L = q$ and $\dim L^{\circ 2} = 2q-1$. A Gorenstein h -vector is symmetric, and its entries sum to the length $2q$. These two facts force

$$h = (1, q-1, q-1, 1).$$

This recovers from the Gorenstein geometry the cubic fullness already forced by Lemma 4.3.

If the reference matching changes, every point translates by one vector c . Expanding the signed third moment after $x_M \mapsto x_M + c$ leaves μ_3 unchanged: the other terms contain μ_2, μ_1 , or $\mu_0 = \sum_M \epsilon(M) = 0$.

Every element of G^+ preserves both sheets and hence fixes the signed sum. Every element of the outer coset exchanges the sheets and negates it. Since $\mu_3 \neq 0$, these two transformation laws give (4.6). Finally (4.5) proves the stated minimality inside the signed tensor-moment family. $\square$

Corollary 4.5 (Graded evaluation algebra). *Put $L^{\circ 0} = \langle 1 \rangle$, and let $L^{\circ d}$ be the span of the pointwise products of d elements of L . For $T = B_3, H_3$, the pointwise powers of the affine evaluation space satisfy*

$$\dim L^{\circ d} = \begin{cases} 1, & d = 0, \\ q, & d = 1, \\ 2q-1, & d = 2, \\ 2q, & d \geq 3. \end{cases}$$

In particular, $L^{\circ 3} = \mathbb{F}_q^{\Omega_T}$. The homogenized configuration

$$\widehat{X}_T = \{[1 : x_M] : M \in \Omega_T\} \subseteq \mathbb{P}^{q-1}$$

therefore has Hilbert function

$$1, \quad q, \quad 2q-1, \quad 2q, \quad 2q, \dots$$

and h -vector $(1, q-1, q-1, 1)$.

Proof. Pair functions on Ω_T with the sheet sign by $\langle \epsilon, f \rangle = \sum_M \epsilon(M) f(M)$. Proposition 4.1 gives $L^{\circ 2} = \ker \langle \epsilon, - \rangle$, with ϵ of full support. Lemma 4.3 gives $L^{\circ 3} = \mathbb{F}_q^{\Omega_T}$. Multiplication by 1 gives the same equality in every higher degree. Homogeneous degree- d forms on $\widehat{X}_T$ restrict to $L^{\circ d}$, which gives the Hilbert function; its first difference is the stated h -vector. Changing the reference matching translates every x_M . Such a translation preserves L and all its pointwise powers and acts projectively on $\widehat{X}_T$, so the graded algebra and Hilbert function are independent of the reference. $\square$

Corollary 4.6 (Self-association and cubic duality). *For $T = B_3, H_3$, the configuration $\widehat{X}_T \subseteq \mathbb{P}^{q-1}$ is a reduced self-associated arithmetically Gorenstein set of $2q$ points. Every one-point deletion imposes independent conditions on quadrics, so $\widehat{X}_T$ has the Cayley–Bacharach property in degree two. Its dualizing residue vector is the sheet sign ϵ . The Artinian reduction by the homogenizing coordinate has Hilbert function $(1, q-1, q-1, 1)$, and its Macaulay inverse system is the cubic line $\mathbb{F}_q \mu_3$.*

Proof. The proof of Theorem 4.4 already establishes signed Gale self-duality, the Cayley–Bacharach property in degree two, the Gorenstein coordinate ring, and the displayed h -vector. These arguments commute with base change to an algebraic closure, so the Gorenstein conclusion descends to $\mathbb{F}_q$.

Let z be the homogenizing coordinate. It is nonzero at every point and is therefore regular on the coordinate ring. The degree d part of the reduction modulo z is $L^{\circ d}/L^{\circ(d-1)}$, so its Hilbert function is the h -vector from Corollary 4.5. The functional

$$f \mapsto \sum_M \epsilon(M) f(x_M)$$

annihilates degrees below three and is nonzero in degree three. Thus ϵ is the socle residue vector. Degree-three polarization identifies its Macaulay dual generator with μ_3 . $\square$

The implication above is the zero-defect case of the modern self-dual-code criterion: the Gorenstein defect is measured by the defect of the Schur square, or equivalently by decomposability [6]. Here $\dim L^{\circ 2} = 2q - 1$ already places both configurations in the zero-defect case. The ideal, deletion, socle, inverse-system, and minimal-resolution calculations are also checked directly in the verification bundle. Self-association does not select an affine origin. Translating all x_M acts projectively on the columns of A and preserves the signed Gale identity, in agreement with the nontrivial base-choice obstruction behind the quotient construction.

The cubic remembers an orientation that the linear quotient does not. Indeed, the signed linear relation

$$\sum_{M \in \Omega_+} x_M = \sum_{M \in \Omega_-} x_M$$

already shows that the sheet sign is killed by the linear quotient map. Quadratic products recover the unordered partition through (4.2), and the cubic supplies the sign exchanged when the two recovered levels are swapped.

Corollary 4.7 (Secant-product tensor syzygies). *For $T = B_3, H_3$, let $m_Q : R_{m-2} \rightarrow R_m$ denote multiplication by Q . Regarding the displayed powers as symmetric tensors, not polynomial powers, the canonically scaled secant products satisfy*

$$\begin{aligned} \sum_M \epsilon(M) P_M &= 0, \\ \sum_M \epsilon(M) P_M^{\odot 2} &= 0, \\ \sum_M \epsilon(M) P_M^{\odot 3} &= \text{Sym}^3(m_Q)(\mu_3) \neq 0. \end{aligned} \tag{4.7}$$

Proof. Substitute $P_M = P_{M_T} + Qx_M$ and expand. Each sheet has $q = 0$ elements in $\mathbb{F}_q$, and the signed quotient moments vanish through degree two. All terms below cubic degree cancel, leaving $\text{Sym}^3(m_Q)(\mu_3)$ in degree three. Since $\mathbb{F}_q[X, Y, Z]$ is a domain, multiplication by Q is injective; over a field, the symmetric powers of an injective linear map are injective. $\square$

Conclusion

Restriction to a conic erases the pairing of marked points, but division by its equation retains a structured record of the factorization. The four-endpoint switch produces an affine quotient configuration, and the three rank-three systems span spaces of dimensions

3, 6, 10. In type H_3 , the quotient retains the top harmonic part and deepest radial line while losing exactly QH_2 ; its base-choice cocycle also obstructs an equivariant affine origin.

In the balanced B_3 and H_3 cases, quadratic products recover the unordered factorization sheets, and their first signed tensor moment is cubic. The same cubic is the Macaulay inverse system of the self-associated arithmetically Gorenstein homogenization. Thus the information forgotten by conic restriction returns in two stages: quadratic data recover the unoriented factorization, and cubic data recover its orientation.

The completeness theorem shows that, among full projective matching orbits, a unique two-level quadratic trade whose levels are one-factorizations can occur only for $B_3/\mathbb{F}_7$ and $H_3/\mathbb{F}_{11}$. The remaining structural question is whether the one-factorization hypothesis follows from weaker intrinsic properties of the quotient configuration. The linear ranks have one proof from representations—the affine connecting cocycle, its irreducible Fischer summands, and a radial trace—but the two radial nonvanishing witnesses remain type-specific.

The appendix refinements require a chosen A_4 -structure and do not canonically identify the depth plane with the relative-cubic plane.

The appendices record further consequences and refinements of the H_3 configuration that are not needed for the main theorem. Appendices A–C treat decorated incidence profiles, their modular depth, and arithmetic marker splitting; Appendices D and E isolate the resulting cubic structures. The final appendix gives the verification architecture for all statements.

A Six profiles and matching-row reconstruction

We now specialize to H_3 . Put

$$G = \mathrm{PGL}_2(11), \quad G^+ = \mathrm{PSL}_2(11), \quad X = \Omega_{H_3} \simeq G/H,$$

where $H \simeq A_5$ is the stabilizer of the base matching in (3.1). Thus $X = X_+ \sqcup X_-$, with $|X_+| = |X_-| = 11$. The construction has four stages:

$$(A_+, A_-) \mapsto K \mapsto \text{four oriented cell pairs} \mapsto D(M) \mapsto K \backslash G/H.$$

The ordered parent pair supplies the A_4 -subgroup K ; its cell partition supplies four signed incidence counts; those counts recover the double-coset label. Only the first two stages require coordinates.

A.1 The decorated parent correspondence

On $V = \mathbb{F}_{11}^3$, set

$$q_0(x, y, z) = x^2 + y^2 + z^2, \quad \mathcal{C}_0 = \mathbb{P}\{q_0 = 0\}.$$

The projectivity

$$\psi = \begin{pmatrix} 10 & 2 & 1 \\ 1 & 2 & 10 \\ 3 & 0 & 3 \end{pmatrix} \quad \text{satisfies} \quad q_0(\psi(X, Y, Z)^t) = 3(XZ - Y^2), \quad (\text{A.1})$$

so $\psi \circ \nu$ identifies the endpoint conic of Section 2 with $\mathcal{C}_0$. In these coordinates take the two six-point sets

$$\begin{aligned} A_+ &= \{(0 : 1 : 4), (0 : 1 : 7), (1 : 0 : 3), (1 : 0 : 8), (1 : 4 : 0), (1 : 7 : 0)\}, \\ A_- &= \{(0 : 1 : 3), (0 : 1 : 8), (1 : 0 : 4), (1 : 0 : 7), (1 : 3 : 0), (1 : 8 : 0)\}. \end{aligned} \quad (\text{A.2})$$

Let $\mathcal{P} = G \cdot A_+$, where G is viewed as the full projective stabilizer of $\mathcal{C}_0$. For $A \in \mathcal{P}$ and $a \in A$, let $C_{A,a}$ be the conic through $A \setminus \{a\}$. The six pairs

$$C_{A,a} \cap \mathcal{C}_0, \quad a \in A, \quad (\text{A.3})$$

form a perfect matching M_A of the twelve points of $\mathcal{C}_0$. On this 22-point orbit, $A \mapsto M_A$ is a G -equivariant bijection onto X . This is the *matching-decorated parent correspondence*; its bijectivity and the assertion that (A.3) consists of six disjoint pairs are exact finite inputs. It is the secant-factorization counterpart of the self-polar matching and Brianchon-support reconstruction obtained from the Clebsch decoding geometry [7].

Put $H_{\pm} = \text{Stab}_G(A_{\pm})$ and

$$K = H_+ \cap H_-, \quad J = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}. \quad (\text{A.4})$$

Then $K \simeq A_4$, $J(A_+) = A_-$, and J normalizes K and exchanges X_+ with X_- . Thus the *ordered golden pair* means precisely the ordered pair (A_+, A_-) .

Choose linear representatives of K preserving q_0 , and let $\widetilde{K} = \mathbb{F}_{11}^{\times} K$. For the following eight vectors, define $R_i = \mathbb{P}(\widetilde{K}r_i)$:

i	r_i	i	r_i
1	(0, 1, 4)	10	(0, 1, 3)
3	(0, 1, 2)	13	(0, 1, 5)
6	(1, 3, 2)	14	(1, 1, 5)
9	(1, 4, 2)	11	(1, 2, 4)

(A.5)

These are cells in the scalar- K orbit partition of $\mathbb{P}(V)$. The involution J exchanges the four pairs

$$(R_1, R_{10}), \quad (R_3, R_{13}), \quad (R_6, R_{14}), \quad (R_9, R_{11}). \quad (\text{A.6})$$

Only these four oriented pairs enter the argument. Reversing the golden pair reverses all four orientations.

This profile layer uses more decoration than the balanced-sheet theorem. The quotient configuration intrinsically recovers the unordered sheets, but it does not select an ordered golden pair or the resulting scalar- K refinement. The reconstruction below is relative to that selected refinement.

A.2 Incidence profiles

For $M \in X$, let S_M be the union of its six secants in $\mathbb{P}^2(\mathbb{F}_{11})$, and put

$$Z_M(R) = |S_M \cap R|.$$

Equivalently, $Z_M(R)$ counts the points of R at which P_M vanishes. Define the *depth profile*

$$D(M) = (Z_M(R_1) - Z_M(R_{10}), Z_M(R_3) - Z_M(R_{13}), \\ Z_M(R_6) - Z_M(R_{14}), Z_M(R_9) - Z_M(R_{11})) \in \mathbb{F}_{11}^4. \quad (\text{A.7})$$

The entries here and in the profile table below are first formed as integer differences of cardinalities, using the canonical cell orientation in (A.6), and only then reduced modulo 11. This definition uses only zero sets of secant products, so it is unchanged if their equations are rescaled.

Theorem A.1 (Six-profile reconstruction). *The K -orbits on each sheet $X_{\pm}$ have sizes 1, 4, 6. With the orientation in (A.6), the corresponding profiles and fibre sizes are*

$sheet$	$ D^{-1}(v) $	v	
+	1	$v_1 = (-6, 0, 12, -12),$	
+	4	$v_2 = (-3, 3, 0, 3),$	
+	6	$v_3 = (3, -2, -2, 0),$	(A.8)
-	1	$-v_1,$	
-	4	$-v_2,$	
-	6	$-v_3.$	

The six vectors are pairwise distinct. Hence D recovers the double-coset row in

$$K \backslash G / H$$

of every matching. Its linear span nevertheless has dimension only two: over $\mathbb{F}_{11}$ it is the plane

$$2a + 2b + c = 0, \quad 9a + 8b + d = 0. \quad (\text{A.9})$$

The two singleton profiles recover their individual matchings. Under the matching-decorated parent correspondence, they therefore recover the corresponding individual H_3 parent. A profile in a fibre of size four or six recovers only that K -orbit, not an individual matching.

Proof mode. The subgroup-mark reduction and the implications from equivariance are conceptual. The six representative incidence rows in (A.10) below are certificate-checked exact counts and are reconstructed by an independent implementation. A companion certificate verifies that the matching decoration uniquely recovers its parent on the 22-point H_3 locus.

Proof. The permutation character of K on either eleven-point sheet has values

$$(11, 3, 2)$$

on elements of orders 1, 2, 3. The marks of the transitive A_4 -sets are

K -set	size	marks on orders 1, 2, 3
K/K	1	(1, 1, 1)
K/V_4	3	(3, 3, 0)
K/C_3	4	(4, 0, 1)
K/C_2	6	(6, 2, 0)
$K/1$	12	(12, 0, 0).

Comparing marks gives the unique nonnegative decomposition

$$X_+|_K \simeq K/K \sqcup K/C_3 \sqcup K/C_2.$$

The orbit sizes are therefore 1, 4, 6; conjugation by J gives the same decomposition on X_- .

For $k \in K$, the transformation k preserves every cell in (A.6) and carries S_M to S_{kM} . Thus $D(kM) = D(M)$. The involution J exchanges the two cells in every oriented pair and carries S_M to S_{JM} , so

$$D(JM) = -D(M).$$

It remains to count one representative on each positive-sheet orbit. The exact incidence rows are

orbit size	$\text{Stab}_K(M)$	zero counts in the four pairs of (A.6)	$D(M)$	
1	A_4	(0, 6; 0, 0; 12, 0; 0, 12)	$v_1,$	
4	C_3	(3, 6; 3, 0; 3, 3; 6, 3)	$v_2,$	
6	C_2	(3, 0; 1, 3; 2, 4; 6, 6)	$v_3.$	(A.10)

The J -images supply the three negative rows. These six vectors are pairwise distinct, so their fibres are precisely the six K -orbits.

Row reduction over $\mathbb{F}_{11}$ gives rank two and the equations (A.9). This rank drop does not merge orbit labels: linear rank and set-theoretic separation are different assertions. Finally, a singleton fibre contains its matching without residual choice. The stabilizer of the matching M_A from (A.3) is the same A_5 as the stabilizer of its parent A . The equivariant decorated transform between the two G/H -orbits is therefore a bijection, so the matching also determines its parent. The stated limitation for the other four rows follows from their fibre sizes. $\square$

Corollary A.2 (Decorated sheet classifier). *Relative to the orientation in (A.6), the profile $D(M)$ determines the sheet containing M . The antipodal singleton pair $\{v_1, -v_1\}$ recovers the unordered golden matching pair and hence the unordered golden parent pair; choosing the orientation selects one member.*

Proof. The positive and negative profile sets in (A.8) are disjoint, and the only singleton fibres are v_1 and $-v_1$. Apply the matching–parent bijection from Theorem A.1. $\square$

This pointwise classifier does not contradict the failure of a linear sheet sign in Section 4. The map D uses zero incidences with the selected scalar- K refinement; it is not a linear functional on the affine quotient configuration. Without that extra decoration, the quadratic and cubic moment statements remain the intrinsic recovery mechanism.

The profile compression also retains the cubic-first orientation from Section 4. The three positive vectors satisfy the integral weighted relation

$$v_1 + 4v_2 + 6v_3 = 0. \quad (\text{A.11})$$

Corollary A.3 (Orbit weights from normalized profiles). *The three incidence-normalized vectors v_1, v_2, v_3 recover the orbit-size multiset $\{1, 4, 6\}$. They also recover the stabilizer orders $\{12, 3, 2\}$ inside $K \simeq A_4$. The underlying rational rays alone do not recover these weights.*

Proof. The coordinates of each v_i are fixed by the cardinality differences in (A.7), so independent rescaling is not allowed. These three integral vectors span a plane over $\mathbb{Q}$, and (A.11) is the primitive positive generator of their relation lattice. Its coefficients recover the three orbit sizes without labels. Orbit–stabilizer then gives $12/1, 12/4, 12/6$. If the vectors are replaced by primitive generators of their rays, the relation becomes $1 : 2 : 1$, which proves the final warning. $\square$

B Modular depth and the projective-cover bridge

Because an H_3 sheet has eleven points in characteristic eleven, the sum of its basis vectors is simultaneously constant and augmentation-zero. Linearizing the three incidence rows therefore kills exactly this constant direction and leaves the two-dimensional profile plane. The following module statement identifies that loss.

Retain the notation of Section A, put

$$k = \mathbb{F}_{11}, \quad L = G^+ = \text{PSL}_2(11), \quad V = k[X_+] \simeq \text{Ind}_H^L \mathbf{1},$$

and regard V as the permutation module on one eleven-point sheet. Let e_1, e_4, e_6 be the indicator functions of the three K -orbits, in orbit-size order. Thus

$$V^K = ke_1 \oplus ke_4 \oplus ke_6, \quad \mathbf{1} = e_1 + e_4 + e_6. \quad (\text{B.1})$$

Linearizing the incidence construction in (A.7) gives

$$\mathcal{D} : V^K \longrightarrow k^4, \quad e_i \longmapsto i u_i, \quad (\text{B.2})$$

where $u_1 = v_1$, $u_4 = v_2$, and $u_6 = v_3$ are the profiles on the orbits of the indicated sizes. Equation (A.11) says exactly that $\mathcal{D}(\mathbf{1}) = 0$.

Proposition B.1 (Modular depth quotient). *The L -module V is the projective cover $P(\mathbf{1})$ and has Loewy dimensions*

$$1 \mid 9 \mid 1.$$

The map $\mathcal{D}$ has kernel $k\mathbf{1}$ and induces an isomorphism

$$P(\mathbf{1})^K / \text{soc } P(\mathbf{1}) \xrightarrow{\sim} \langle u_1, u_4, u_6 \rangle_k. \quad (\text{B.3})$$

The sheet exchanged by J carries the negative copy of this quotient.

Proof mode. Projectivity, indecomposability, fixed-point exactness, and the socle quotient are conceptual. Absolute irreducibility of the nine-dimensional middle layer and the rank and kernel of $\mathcal{D}$ are exact generator-matrix calculations in the `relative-cubic-depth` bundle, reconstructed by its independent replay.

Proof. The order of $H \simeq A_5$ is prime to 11, so its trivial module is projective and induction makes V projective. The action of L on X_+ is doubly transitive. Hence $\text{End}_L(V)$ is generated by the identity and the all-ones operator E . In characteristic eleven,

$$E^2 = 11E = 0,$$

so this endomorphism algebra is local. Thus V is indecomposable. Its trivial head occurs once, and consequently $V = P(\mathbf{1})$. The constant vector is its one-dimensional socle. The remaining nine-dimensional layer is absolutely irreducible; equivalently, the exact generator matrices on it span the full 9-by-9 matrix algebra. This gives the displayed Loewy dimensions.

Taking K -fixed points is exact because $11 \nmid |K| = 12$. The orbit sizes 1, 4, 6 give (B.1). The three vectors in (B.2) span the plane (A.9), and their only relation is (A.11), so $\ker \mathcal{D} = k\mathbf{1}$. This proves (B.3). Finally $D(JM) = -D(M)$, which attaches the outer character on the opposite sheet. $\square$

Corollary B.2 (The canonical nine-space bridge). *For the fixed marked conic and sheet X_+ , there is a canonical G^+ -module isomorphism*

$$\text{rad } P(\mathbf{1}) / \text{soc } P(\mathbf{1}) \xrightarrow{\sim} M_8 \downarrow_{G^+}. \quad (\text{B.4})$$

It sends the augmentation class of $\sum_{M \in X_+} a_M [M]$ to

$$\sum_{M \in X_+} a_M \pi_8(x_M), \quad \sum_M a_M = 0, \quad (\text{B.5})$$

where π_8 is the top-harmonic projection. Formula (B.5) is independent of the reference matching.

Proof. The unique trivial head map in Proposition B.1 is the augmentation map, so

$$\text{rad } P(\mathbf{1}) = \ker(V \xrightarrow{\sum} k).$$

Changing the reference matching translates every x_M by the same vector, so the augmentation condition makes (B.5) reference-independent. The affine cocycle term in x_{gM} cancels for the same reason, hence the map is G^+ -equivariant. The same-sheet first-moment identity of Proposition 4.1 sends the constant vector $\mathbf{1}$, which lies in the augmentation module because $|X_+| = 11 = 0$ in k , to zero. Thus (B.5) descends to $\text{rad } P(\mathbf{1})/\text{soc } P(\mathbf{1})$.

Its image contains a nonzero same-sheet quotient and is G^+ -stable. The irreducibility of $M_8 \downarrow_{G^+}$ therefore makes it surjective. Both sides have dimension nine, so it is an isomorphism. $\square$

Corollary B.3 (The homogeneous projective-cover bridge). *For the cocycle representative c_8 determined by the chosen reference matching, let E_{c_8} be the G -module defined in (3.11b). There is a G^+ -module isomorphism*

$$P(\mathbf{1})/\text{soc } P(\mathbf{1}) \xrightarrow{\sim} E_{c_8} \downarrow_{G^+}, \quad (\text{B.6})$$

given by

$$\Theta \left(\sum_{M \in X_+} a_M [M] \right) = \left(\sum_{M \in X_+} a_M \pi_8(x_M), \sum_{M \in X_+} a_M \right). \quad (\text{B.7})$$

Thus the extension class, unlike a cocycle representative or an affine origin, is canonical: the top-harmonic homogeneous extension restricts to the nonsplit exact sequence

$$0 \longrightarrow M_8 \downarrow_{G^+} \longrightarrow P(\mathbf{1})/\text{soc } P(\mathbf{1}) \longrightarrow \mathbf{1} \longrightarrow 0.$$

If $\tilde{E}_c$ denotes the homogenization of the full affine quotient with translation space $M_8 \oplus M_0$, then

$$\tilde{E}_c \downarrow_{G^+} \simeq P(\mathbf{1})/\text{soc } P(\mathbf{1}) \oplus \mathbf{1}, \quad (\text{B.8})$$

where the last summand is the radial module $M_0 \downarrow_{G^+}$.

Proof. The affine transformation law $\pi_8(x_{gM}) = g\pi_8(x_M) + c_8(g)$ makes Θ G^+ -equivariant. The constant vector maps to zero: its second coordinate is $|X_+| = 11 = 0$ in k , and its first coordinate vanishes by the same-sheet first-moment identity. Hence Θ descends through the socle.

On the kernel of augmentation, the descended map is the isomorphism of Corollary B.2; on the one-dimensional head it induces the identity through the second coordinate. It is therefore an isomorphism. The exact sequence cannot split: Proposition B.1 gives $P(\mathbf{1})/\text{soc } P(\mathbf{1})$ the simple socle M_8 , whereas a splitting would put a trivial summand in its socle. Finally, $M_0 = \chi$ is trivial on G^+ , and its cocycle vanishes there, proving (B.8). $\square$

Remark B.4 (Ambient and intrinsic sheet recovery). With the harmonic–radial decomposition retained, the M_0 -projection already separates the two H_3 sheets: it is zero on X_+ , constant on X_- , and nonzero by the witness (3.11). The quadratic theorem of Section 4 is stronger in a different sense: it recovers the sheets intrinsically from the abstract affine point configuration, without retaining the ambient Fischer operator.

This quotient identifies the precise modular loss in the six-profile construction. Before linearization, the six profiles remain distinct and recover all six double-coset labels. After linearization, the constant orbit trade dies and only the two-dimensional depth plane remains. Conversely, the three rational rays in that plane remain distinct but do not retain their orbit weights. The weights 1, 4, 6 are recovered only when the quotient is supplied with the incidence-normalized vectors of (A.8), as in Corollary A.3.

Remark B.5 (Reduction of the profile table). The four-coordinate profiles in (A.8) are integral before reduction. The greatest common divisor of their nonzero 2-by-2 minors is 3. Their span therefore drops below rank two only in characteristic 3. The six signed rows remain distinct away from characteristics 2 and 3: characteristic 2 identifies antipodes, while characteristic 3 sends the rows v_1 and v_2 to zero. This is a statement about the displayed integral table, not a construction of the marked conic geometry in those characteristics.

C Arithmetic splitting of the marker algebra

The arithmetic representatives of the three configurations come in conjugate pairs. Besides the base matchings in (3.1), we use

$$\begin{aligned} M_{B,-} &= \{\{0, \infty\}, \{1, 3\}, \{2, 6\}, \{4, 5\}\}, \\ M_{B,+} &= \{\{0, \infty\}, \{1, 5\}, \{2, 3\}, \{4, 6\}\}, \\ M_{H,-} &= \{\{0, 10\}, \{1, \infty\}, \{2, 7\}, \{3, 5\}, \{4, 8\}, \{6, 9\}\}. \end{aligned} \tag{C.1}$$

The pair for A_3 consists of two copies of M_{A_3} , the pair for B_3 is $(M_{B,-}, M_{B,+})$, and the pair for H_3 is $(M_{H_3}, M_{H,-})$. To make the arithmetic bridge part of the definition, let $\bar{k}$ be an algebraic closure of the working field and define the root-specialization maps

$$\begin{array}{ccc} T & \text{root} & \sigma_T(\text{root}) \\ \hline A_3 & \sqrt{2}, -\sqrt{2} & M_{A_3}, M_{A_3} \\ B_3 & 3, 4 & M_{B,-}, M_{B,+} \\ H_3 & 4, 8 & M_{H_3}, M_{H,-}. \end{array} \tag{C.2}$$

Thus the *marker datum* is the quadratic algebra together with σ_T , not the polynomial alone. Exact projective transport places the displayed B_3 and H_3 values in Ω_{B_3} and Ω_{H_3} . The certificate also verifies that σ_T is the reduction of the corresponding integral Coxeter frames; the theorem below needs only the explicit map (C.2). We call the datum *fused* when its two geometric roots have one rational matching value, and *split* when they give two distinct rational matching sheets.

Lemma C.1 (Split-inert frame calculation). *The three marker polynomials factor as follows:*

$$\begin{array}{lll} A_3, q = 5 & t^2 - 2 & \text{irreducible,} \\ B_3, q = 7 & t^2 - 2 = (t - 3)(t - 4) & \text{split,} \\ H_3, q = 11 & t^2 - t - 1 = (t - 4)(t - 8) & \text{split.} \end{array} \tag{C.3}$$

In the first row the two geometric roots form one Frobenius orbit over $\mathbb{F}_{25}$. In each split row the nontrivial involution of the quadratic marker algebra exchanges the two rational roots.

Proof. The nonzero squares modulo 5 are 1 and 4, so 2 is not a square. If $u^2 = 2$ in $\mathbb{F}_{25}$, the nontrivial $\mathbb{F}_5$ -automorphism sends u to $-u$, and hence exchanges the two roots. The remaining factorizations follow by direct substitution. $\square$

Theorem C.2 (Marker specialization and H_3 arithmetic gluing). *For these arithmetic representatives of the rank-three matching configurations, the following hold.*

- (a) *The A_3 marker datum is fused over $\mathbb{F}_5$. The B_3 and H_3 marker data split into two rational matching sheets over $\mathbb{F}_7$ and $\mathbb{F}_{11}$, respectively.*

- (b) In the H_3 case, the stabilizers H_+ and H_- of the singleton matching in each rational sheet are isomorphic to A_5 , satisfy

$$H_+ \cap H_- = K \simeq A_4, \quad \langle H_+, H_- \rangle = \mathrm{PSL}_2(11), \quad (\mathrm{C.4})$$

and are exchanged by a projectivity J of nonsquare determinant. The normalizer $N_G(K)$ is a rational S_4 whose determinant-square kernel is K .

- (c) The determinant character that distinguishes the two rational sheets also negates the depth profiles. Thus choosing one root in the split marker algebra chooses one sheet, while its singleton profile chooses the corresponding decorated matching row. Forgetting that choice leaves the unordered pair.

Proof mode. Lemma C.1 and the determinant-character deductions are conceptual. The agreement of the specialization map (C.2) with the integral Coxeter frames, the matching stabilizers and intersections, and the generated subgroup in (C.4) are exhaustive exact finite checks. The classical subgroup names use the standard subgroup structure of $\mathrm{PGL}_2(11)$ [1, 4].

Proof. Part (a) records the specialization data in (C.2). Its two A_3 roots have the same matching value. The two values in the B_3 row are exchanged by $x \mapsto -x$, and the two values in the H_3 row are exchanged by

$$x \mapsto \frac{x-1}{x+1}.$$

The exact transport check verifies these identities on the displayed matchings. Together with Lemma C.1, they prove (a).

For H_3 , exact projective closure gives

$$|H_+| = |H_-| = 60, \quad |H_+ \cap H_-| = 12, \quad |\langle H_+, H_- \rangle| = 660.$$

Both stabilizers lie in the determinant-square subgroup, while the displayed transporter has nonsquare determinant. The normalizer $N_G(K)$ has order 24, and its determinant-square kernel has order 12. The cited subgroup identifications give (C.4) and the S_4/A_4 hinge, proving (b).

The determinant-square subgroup has the two matching sheets as its orbits, and J exchanges them. Section A proves $D(JM) = -D(M)$ and identifies the unique singleton in each sheet. Theorem A.1 then returns the decorated matching row from that singleton. This proves (c). $\square$

Arithmetic splitting and modular depth retain different amounts of information. Splitting produces a two-point choice of sheet. The depth quotient is defined only after the A_4 -refinement has been selected and remembers all three orbit rays modulo the constant socle. The first choice orients the second, but does not canonically identify it with any cubic quotient of the ten-dimensional factorization space. Appendix E makes that boundary precise.

D Cubic structure of the profile plane

Lemma D.1 (Three-ray cubic forcing). *Let k have characteristic different from 2 and 3. Suppose u_1, u_2 form a basis of a two-dimensional k -space and $n_1, n_2, n_3 \in k^\times$ satisfy*

$$n_1 u_1 + n_2 u_2 + n_3 u_3 = 0.$$

For the signed antipodal measure

$$\xi = \sum_{i=1}^3 n_i (\delta_{u_i} - \delta_{-u_i}),$$

the moments of degrees one and two vanish, while the cubic moment is nonzero.

Proof. The relation kills the first moment, and antipodality kills every even moment. Write

$$u_3 = -\frac{n_1}{n_3}u_1 - \frac{n_2}{n_3}u_2.$$

In $\sum_i n_i u_i^{\odot 3}$, the coefficient of $u_1^{\odot 2} \odot u_2$ is

$$-\frac{3n_1^2 n_2}{n_3^2},$$

which is nonzero. The signed cubic is twice this tensor and is therefore nonzero as well. $\square$

Corollary D.2 (The mass-zero cubic flag). *In Lemma D.1, suppose also that $n_1 + n_2 + n_3 = 0$. Then its half-cubic factors as*

$$\frac{1}{2}M_3(\xi) = \sum_{i=1}^3 n_i u_i^{\odot 3} = \frac{n_1 n_2}{(n_1 + n_2)^2} (u_1 - u_2)^{\odot 2} \odot ((2n_1 + n_2)u_1 + (n_1 + 2n_2)u_2). \quad (\text{D.1})$$

The two displayed lines are distinct. Hence the cubic intrinsically defines a doubled line and a residual simple line; its Hessian recovers the doubled line.

Proof. Since $n_3 = -(n_1 + n_2)$, the weighted vector relation gives

$$u_3 = \frac{n_1 u_1 + n_2 u_2}{n_1 + n_2}.$$

Substitution and expansion give (D.1). The residual factor would equal the doubled factor only if $3(n_1 + n_2) = 0$, which is impossible under the hypotheses. The Hessian of a binary cubic $L^2 R$, with L and R distinct, is a nonzero scalar multiple of L^2 . $\square$

For the incidence-normalized profiles of Section A, put

$$\eta = \delta_{v_1} + 4\delta_{v_2} + 6\delta_{v_3} - \delta_{-v_1} - 4\delta_{-v_2} - 6\delta_{-v_3}.$$

Then

$$M_j(\eta) = (1 - (-1)^j) \sum_{i=1}^3 n_i v_i^{\odot j}, \quad (n_1, n_2, n_3) = (1, 4, 6). \quad (\text{D.2})$$

Equation (A.11) kills the first moment, and antipodality kills every even moment. The first two profiles are independent over $\mathbb{F}_{11}$, since the determinant of their first two coordinates is $-18 \neq 0$. Lemma D.1 therefore forces the cubic to be nonzero. In the frozen normalization, its first coordinate is

$$2((-6)^3 + 4(-3)^3 + 6(3)^3) = 6 \neq 0 \quad \text{in } \mathbb{F}_{11}. \quad (\text{D.3})$$

Thus the six-row quotient preserves the cubic-first signed moment even though its linear image is only a plane.

Remark D.3 (The H_3 cubic flag). Here $1 + 4 + 6 = 0$ in $\mathbb{F}_{11}$, so Corollary D.2 forces the double-line factorization. In the basis $e_1 = v_2, e_2 = v_3$ of the profile plane, the projective cubic is

$$f(x, y) = x^3 + 7x^2y + 5xy^2 + 9y^3 = (x - y)^2(x - 2y),$$

with Hessian $7x^2 + 8xy + 7y^2 = 7(x - y)^2$. Hence the cubic intrinsically distinguishes a doubled line and a residual simple line. This flag is internal to the decorated profile plane: it supplies no canonical map from the full relative-cubic space and no descended G^+ -action.

E The relative-cubic Tate plane

Let $W = W_{H_3}$ be the ten-dimensional quotient space from Theorem 3.2, let χ be the determinant character of $G = \mathrm{PGL}_2(11)$, and put

$$M = \mathrm{Sym}^3(W) \otimes \chi, \quad R = M^G.$$

The signed cubic of Section 4 lies in R , which has dimension three. There is a canonical two-dimensional quotient of this source space, but it is not canonically the depth plane of Proposition B.1.

We fix the monomial basis of $\mathrm{Sym}^3(W)$ induced by the harmonic–radial coordinates of Section 3. The *frozen relative basis* is the reduced-echelon basis of the simultaneous equations $gf = \chi(g)f$ for the standard generators defining the marked H_3 orbit, ordered by its pivots. Thus the coordinates below specify a reproducible basis, not an additional intrinsic structure. The *three labelled cubic orbit classes* are indexed, in order, by the three K -orbits of sizes 1, 4, 6 on X_+ . Finally, the *invariant symmetric-form pencil* is $\mathbb{P}((\mathrm{Sym}^2 W)^G)$. Its unique rank-one point has the form $[\ell^2]$; contraction by its covector ℓ is the induced map

$$\iota_\ell : R \longrightarrow (\mathrm{Sym}^2 W)^G.$$

Proposition E.1 (Canonical relative-cubic Tate plane). *The invariant space R and the coinvariant space M_G both have dimension three. The canonical projection and group norm fit into*

$$R \xrightarrow{\pi} M_G \xrightarrow{N} R, \quad \mathrm{rank} \pi = 2, \quad \mathrm{rank} N = 1, \quad (\text{E.1})$$

with

$$\mathrm{im} \pi = \ker N, \quad \mathrm{im} N = \ker \pi = \langle [1 : 3 : 9] \rangle. \quad (\text{E.2})$$

Contraction by the covector whose square is the unique rank-one member of the invariant symmetric-form pencil gives a second construction of the same quotient. In the frozen relative basis its matrix is

$$\begin{pmatrix} 8 & 1 & 0 \\ 0 & 8 & 1 \end{pmatrix}, \quad (\text{E.3})$$

and its kernel is again $\langle [1 : 3 : 9] \rangle$.

Proof mode. The invariant/coinvariant construction and the implications of the displayed ranks are conceptual. The dimensions, norm and projection ranks, kernel line, rank-one pencil member, contraction matrix, and flattening census are exact finite linear algebra in the **relative-cubic-depth** bundle and its independent replay.

Proof. Exact invariant and commutator calculations give $\dim R = \dim M_G = 3$, the two ranks in (E.1), and $N\pi = 0$. Thus $\mathrm{im} \pi \subseteq \ker N$; both spaces have dimension two, so they are equal. The computed image of N is the displayed kernel of π , proving (E.2).

The invariant symmetric-form pencil on W has a unique rank-one member ℓ^2 , where ℓ transforms by χ . Contracting a χ -twisted cubic by ℓ cancels the two characters and lands in the invariant quadratic pencil. Exact contraction gives (E.3), whose kernel proves the second assertion. $\square$

The kernel line is intrinsic rather than an artifact of the displayed basis. Among the 133 projective lines of R , it is the unique line whose first flattening has rank 9; the remaining census is one line of rank 1 and 131 lines of rank 10.

Remark E.2 (Why the two canonical planes do not glue). The three labelled cubic orbit classes project to M_G with relation $[2, 9, 1]$, whereas the three labelled depth profiles have relation $[2, 8, 1]$. There is therefore no label-preserving linear map carrying these source classes to the depth rays. The integral transfer makes the obstruction structural. In orbit-size order set

$$s = (1, 1, 1)^t, \quad B = s(1, 4, 6).$$

Then $B^2 = 11B$. Divided transfer kills every integral balanced relation a with $(1, 4, 6)a = 0$, but $Bs = 11s$, so it fixes the depth socle after division by 11. It separates the source relation from the depth socle instead of identifying them. The doubled-line and residual-line flag of Remark D.3 is internal to the depth plane and supplies no missing map.

F Verification architecture

Table 1 separates the mathematical arguments from their exact finite inputs. Here “certificate” means a deterministic calculation over a prime field, checked against a canonical JSON output and a SHA-256 manifest; an independent replay uses a separate implementation. “Lean” means that the stated finite implication is checked by the Lean kernel. Neither term replaces the cited identification of the classical groups or configurations.

Result	Proof in the text or classical input	Exact evidence
Matching-secant quotient	unique conic divisibility and the four-endpoint switch	none
Balanced-orbit completeness	Dickson–Giudici maximal subgroups, exceptional normalizers, and unique two-point block systems	none
Ranks and the middle Fischer layer	affine cocycle, irreducible Fischer modules; Edge–Dye configurations	quotient matrices as cross-checks; one radial scalar for H_3 ; independent replay
Balanced sheets and cubic orientation	Propositions 4.1 and 4.2; Lemma 4.3; signed Gale duality	finite moments and ranks as cross-checks; Lean Hilbert arithmetic; Gorenstein replay
Six-profile reconstruction	subgroup marks, orbit–stabilizer, and the three-ray lemmas	profile rows and replay; decorated-parent bijection
Modular depth and the Tate plane	projectivity, fixed-point exactness, and standard modular-module input	invariant, coinvariant, and depth calculations with replay
Arithmetic splitting and gluing	displayed root calculations; cited subgroup identifications	Lean gate and normalized certificate/replay

Table 1: Proof modes for the principal results.

The claim-by-claim map is `verification/trust_manifest.json`; the exact twenty-one-statement identity is `verification/statement_identity.json`. From the root of the archival source, the aggregate entry point is

```
python3 papers/clebsch-factorization/verification/verify_release.py
```

It checks the statement identity and all admitted checksum manifests, runs the admitted primary checks and independent replays, elaborates the arithmetic-gluing and Hilbert-symmetry Lean gates, and rebuilds this manuscript through the repository Makefile.

The enumerated review closure accompanying this candidate is pinned by

```
verification/evidence_fingerprint.json.
```

Its SHA-256 digest is

71e27a2468c12177c36b8df2f8b7061bcc0a7b
e78bda9230e3288bcb609a2972.

That fingerprint fixes a normalized manuscript, statement contents, extractor, paper Makefile, verification documentation, eight evidence manifests, command vectors, Lean gates, runner, trust manifest, and environment: Python 3.13.12, Lean 4.32.0-rc1, and Mathlib revision

571b8a8e54219b4d393f75f4b8653fac08197fcc.

A successful replay ends with

`clebsch factorization release: CHECK OK.`

The normalization replaces only the displayed fingerprint digest and the statement identity's derived full-source hash, avoiding a circular self-hash. This is an explicit review-source allowlist, not a claim to hash every file in the repository or every transitive tool dependency. An immutable public archive locator must be added before release.

The largest formal component is the gate

`RelativeConicArcs.Gates.
ClebschArithmeticGluIng.`

Its twenty-three terminals check the displayed roots, matching involutions, projective orders, determinant halves, the A_3/B_3 orbit calculations, and the split-fused conclusion. For H_3 , Lean checks the normalized certificate rows, intersections, determinant classes, the $11 + 11$ signature split, and the shape of the bounded generation words. Evaluation and completeness of the 660-element coset and word certificates remain exact generator/replay obligations. The classical names S_4 , A_4 , A_5 , and $\mathrm{PSL}_2(11)$ are cited inputs: no group name is inferred from an order.

The separate structural gate

`RelativeConicArcs.Gates.
ClebschHilbertSymmetry`

checks the arithmetic implication used after the classical Gorenstein criterion: a symmetric finite Hilbert function of total length $2q$ whose first values are $1, q - 1, q - 1$ has socle degree three and final value one. It does not formalize the Eisenbud–Popescu criterion, the identification of the first Hilbert values, or Gorenstein symmetry.

The other computations are load-bearing only at the points shown in Table 1. The matching-module bundle supplies the three finite quotient ranks and cross-checks the moment data. The H_3 -equivariant-rank bundle checks the Sylow-normalizer weights, A_5 -fixed dimensions, sheet membership, and minimal top and radial witnesses. The balanced-sheet bundle independently reconstructs the permutation-module conclusions, while the Gorenstein bundle checks the ideals, deletion ranks, socles, and resolutions. The profile-incidence bundle computes one row per double coset; the decorated-parent bundle identifies a singleton matching with its parent; the relative-cubic-depth bundle computes the modular and Tate rank patterns; and the arithmetic-gluIng gate checks the normalized splitting data. All use exact prime-field arithmetic. The balanced-orbit classification, general divisibility theorem, radical–Hadamard and hyperplane-square lemmas, modular-permutation mechanism, three-ray cubic lemma, and mass-zero factorization are proved in the text and do not depend on those enumerations.

This verification surface proves only the three marked configurations defined here. It does not supply an all-characteristic construction, a canonical map from the relative-cubic Tate plane to the depth plane, or any passage, holonomy, theta, Fourier, quantum, or Mathieu identification.

References

- [1] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson, *Atlas of Finite Groups*, Clarendon Press, Oxford, 1985.
- [2] W. L. Edge, Conics and orthogonal projectivities in a finite plane, *Canadian Journal of Mathematics* **8** (1956), 362–382.
- [3] R. H. Dye, Hexagons, conics, A_5 and $\mathrm{PSL}_2(K)$, *Journal of the London Mathematical Society* (2) **44** (1991), 270–286. doi:10.1112/jlms/s2-44.2.270.
- [4] M. Giudici, Maximal subgroups of almost simple groups with socle $\mathrm{PSL}(2, q)$, arXiv:math/0703685, 2007.
- [5] D. Eisenbud and S. Popescu, The projective geometry of the Gale transform, *Journal of Algebra* **230** (2000), 127–173. doi:10.1006/jabr.1999.7940.
- [6] G. Rodríguez-Pajares, D. Ruano, and F. Salizzoni, A combinatorial description of when a self-associated set of points fails to be arithmetically Gorenstein, arXiv:2512.16766v1, 2025.
- [7] T. Rudd, *A conic deep-hole syndrome locus characterizes the Clebsch code*, working paper, 2026.
- [8] T. Braun and G. Nebe, Orthogonal representations of $\mathrm{SL}_2(q)$ in defining characteristic, *Beiträge zur Algebra und Geometrie* (2025). doi:10.1007/s13366-024-00763-w.