

Arcs complete outside a prescribed conic

A prescribed-hole defect identity and a certified classification over $\mathbb{F}_{16}$

Tavis Rudd

July 2026

Abstract

Let $\mathcal{C}$ be a nonsingular conic in $\text{PG}(2, q)$. We study arcs $A \subseteq \text{PG}(2, q) \setminus \mathcal{C}$ whose secants cover every point outside $A \cup \mathcal{C}$, and write $\rho_{\mathcal{C}}(q)$ for their minimum size. Starting from the classical first two equations for the secant-index distribution of an arc, we prove an exact defect identity for an arbitrary prescribed set of points that may remain uncovered, together with equality and stability criteria. Its conic specialization gives

$$q^2 - k \leq \binom{k}{2}(q-1) - \frac{6}{\lfloor k/2 \rfloor} \binom{k}{4} - \frac{I_{\mathcal{C}}(A)}{\lfloor k/2 \rfloor}$$

for every $\mathcal{C}$ -complete k -arc, together with an exact equality criterion and a quantitative stability estimate. In particular, for every prime power q ,

$$\rho_{\mathcal{C}}(q) \geq \sqrt{2q} + \frac{3}{2} - \frac{8}{\sqrt{2q}}.$$

Conversely, if H is any prescribed set and an ordinary complete b -arc satisfies $b|H| < q^2 + q + 1$, projective averaging moves it off H ; the conic case gives $\rho_{\mathcal{C}}(q) \leq t_2(2, q)$ whenever $t_2(2, q) \leq q$. We also derive even-characteristic nucleus constraints and formulate the quadratic obstruction as a finite-field evaluation criterion. Together with the lower bounds below, explicit witnesses give

$$\rho_{\mathcal{C}}(8) = \rho_{\mathcal{C}}(9) = \rho_{\mathcal{C}}(11) = 6, \quad \rho_{\mathcal{C}}(16) = 9.$$

The last equality follows from a certified classification of all 2633 projective eight-arcs over $\mathbb{F}_{16}$: no nonzero quadratic contains an eight-arc's ordinary-uncovered locus while avoiding the arc. Supplementary checkers and a companion Lean development verify the witnesses, theorem chain, and finite classification certificates. The arc-MDS dictionary also turns the defect into a syndrome-leader collision identity; for the $q = 11$ witness it yields a non-GRS MDS code whose deep-hole locus is the conic and whose simultaneous extensions form the icosahedral independence complex.

1 Introduction

An arc in a projective plane is a set of points no three of which are collinear. An arc is complete when it cannot be enlarged while remaining an arc; equivalently, every point outside it lies on a secant determined by two of its points. Complete arcs and the closely related problem of finding small saturating sets have a substantial literature; see, for example, Ball [3], Kim–Vu [12], and Nagy [15].

Fix a nonsingular conic $\mathcal{C} \subset \text{PG}(2, q)$. We impose the arc condition but allow the points of $\mathcal{C}$ to remain uncovered.

Definition 1.1. An arc $A \subseteq \text{PG}(2, q) \setminus \mathcal{C}$ is $\mathcal{C}$ -complete if every point of

$$\text{PG}(2, q) \setminus (A \cup \mathcal{C})$$

lies on a secant of A . Define

$$\rho_{\mathcal{C}}(q) = \min\{|A| : A \text{ is a } \mathcal{C}\text{-complete arc}\}.$$

For an arbitrary arc disjoint from $\mathcal{C}$, its uncovered locus is

$$\mathcal{U}_{\mathcal{C}}(A) = \text{PG}(2, q) \setminus \left(A \cup \mathcal{C} \cup \bigcup_{\{a,b\} \in \binom{A}{2}} ab \right).$$

Thus A is $\mathcal{C}$ -complete exactly when $\mathcal{U}_{\mathcal{C}}(A) = \emptyset$.

The minimum in Definition 1.1 is attained. Indeed, in the finite family of arcs disjoint from $\mathcal{C}$, any member of maximum cardinality is maximal under insertion and hence is $\mathcal{C}$ -complete. The same argument works for every prescribed hole set.

Equivalently, A is maximal among arcs in the point set $\text{PG}(2, q) \setminus \mathcal{C}$. All nonsingular conics in $\text{PG}(2, q)$ are projectively equivalent, so the numerical value of $\rho_{\mathcal{C}}(q)$ does not depend on the chosen conic.

The problem differs from two nearby standard notions. A saturating set need not be an arc and has no prescribed exceptional locus. An almost-complete subset of a conic, in the sense of Bartoli–Davydov–Marcugini–Pambianco [5], consists of selected points *on* the conic whose chords cover the complementary points. Ng–Wild’s arcs covering a line [16] prescribe a one-dimensional set that must be covered, rather than an exceptional set that may remain uncovered. Here the selected arc is required to avoid the conic.

There is also classical work on *complete exterior sets*: sets of $(q+1)/2$ exterior points whose pairwise joins miss the conic [6, p. 143]. This condition constrains the joins of the selected points, whereas Definition 1.1 requires those joins to cover the complementary off-conic locus. The two conditions therefore point in opposite directions. If

$$U(A) = \text{PG}(2, q) \setminus \left(A \cup \bigcup_{\{a,b\} \in \binom{A}{2}} ab \right)$$

is the ordinary-uncovered locus, then

$$\begin{aligned} \text{complete exteriority} &\implies \mathcal{C}(\mathbb{F}_q) \subseteq U(A), \\ \mathcal{C}\text{-completeness} &\iff U(A) \subseteq \mathcal{C}(\mathbb{F}_q), \\ \text{both conditions} &\implies U(A) = \mathcal{C}(\mathbb{F}_q). \end{aligned}$$

The strict $q = 7$ example and the exceptional equality at $q = 11$ return in Section 8.

The point index with respect to an arc, and the first two double-counting equations for its distribution, are classical; compare Hirschfeld [11, Chapter 9] and Ball [3]. The use of secant counts to bound the smallest complete (k, n) -arcs is also classical; for a recent explicit lower bound see Alabdullah–Hirschfeld [1]. The main observation of this note is that, after splitting those equations over a prescribed exceptional set and its complement, the inequalities normally used to control overlap have an exact remainder. This produces:

- (i) an exact prescribed-hole defect identity and equality criterion;

- (ii) a quantitative stability bound for near equality;
- (iii) a strengthened lower bound with additive asymptotic term $3/2$;
- (iv) a projective-averaging upper-bound transfer from ordinary complete arcs.

We then develop three applications of this spine: characteristic-two nucleus constraints, an evaluation and coding interpretation, and verified small-field results culminating in $\rho_C(16) = 9$.

The classical inputs are the secant-index equations, the Lunelli–Sce $\sqrt{2q}$ scale, the arc–MDS correspondence, and the known $q = 11$ Clebsch geometry. The contributions here are the prescribed-hole remainder and its consequences, the uncovered-locus evaluation obstruction, and the certified relative-conic conclusions stated above.

2 The classical secant equations

Throughout the remainder of the paper, $q \geq 3$ and $k \geq 3$. Let Π be a projective plane of order q , and let A be a k -arc in Π . Write $r_A(x)$ for the number of secants of A through $x \in \Pi \setminus A$. We usually write $r(x)$. Put

$$N = \binom{k}{2}, \quad m = \left\lfloor \frac{k}{2} \right\rfloor.$$

Lemma 2.1. *For every $x \notin A$, one has $r(x) \leq m$.*

Proof. Distinct secants through x cannot share an endpoint in A . Indeed, two lines through the same pair x, a coincide. Hence the secants through x use pairwise disjoint pairs of points of A , of which there are at most $\lfloor k/2 \rfloor$. $\square$

Proposition 2.2 (Classical first and second index equations). *For every k -arc A in a projective plane of order q ,*

$$\sum_{x \notin A} r(x) = \binom{k}{2} (q - 1), \tag{1}$$

$$\sum_{x \notin A} \binom{r(x)}{2} = 3 \binom{k}{4}. \tag{2}$$

Proof. There are $\binom{k}{2}$ secants. Each contains $q + 1$ points, exactly two of them in A , and therefore contributes $q - 1$ incidences with points outside A . This proves (1).

For (2), count unordered pairs of distinct secants through a common external point. By Lemma 2.1, their four endpoints in A are distinct. Conversely, a four-subset of A has three partitions into two unordered pairs. Each partition gives two secants, which meet in a unique point because the plane is projective. Their intersection is outside A : otherwise one of the two secants would contain three points of the arc. Thus every four-subset contributes exactly three. $\square$

Only the projective-plane axioms and the common line size $q + 1$ enter these proofs. In particular, the second equation need not hold in a general finite linear space where two lines may be disjoint.

3 An exact defect identity for prescribed holes

The next theorem is stated for an arbitrary exceptional point set. The conic enters only in later specializations.

Let $\mathcal{H} \subseteq \Pi \setminus A$ be any set of points allowed to remain uncovered, and define

$$\begin{aligned} V_{\mathcal{H}}(A) &= \Pi \setminus (A \cup \mathcal{H}), \\ \mathcal{X}_{\mathcal{H}}(A) &= \{x \in V_{\mathcal{H}}(A) : r(x) > 0\}, \\ \mathcal{U}_{\mathcal{H}}(A) &= V_{\mathcal{H}}(A) \setminus \mathcal{X}_{\mathcal{H}}(A), \\ I_{\mathcal{H}}(A) &= \sum_{y \in \mathcal{H}} r(y). \end{aligned}$$

Thus $\mathcal{X}_{\mathcal{H}}(A)$ is the covered required locus, and $\mathcal{U}_{\mathcal{H}}(A)$ is the uncovered required locus.

Splitting Proposition 2.2 over $V_{\mathcal{H}}(A)$ and $\mathcal{H}$ gives

$$\sum_{x \in V_{\mathcal{H}}(A)} r(x) = N(q-1) - I_{\mathcal{H}}(A), \quad (3)$$

$$\sum_{x \in V_{\mathcal{H}}(A)} \binom{r(x)}{2} + \sum_{y \in \mathcal{H}} \binom{r(y)}{2} = 3 \binom{k}{4}. \quad (4)$$

Theorem 3.1 (Prescribed-hole defect identity). *Let A be a k -arc in a projective plane of order q , let $\mathcal{H} \subseteq \Pi \setminus A$, and put $m = \lfloor k/2 \rfloor$. Define*

$$\Delta_{\mathcal{H}}(A) = N(q-1) - \frac{6}{m} \binom{k}{4} - \frac{I_{\mathcal{H}}(A)}{m} - |\mathcal{X}_{\mathcal{H}}(A)|.$$

Then

$$m\Delta_{\mathcal{H}}(A) = \sum_{x \in \mathcal{X}_{\mathcal{H}}(A)} (r(x) - 1)(m - r(x)) + \sum_{y \in \mathcal{H}} r(y)(m - r(y)). \quad (5)$$

In particular, $\Delta_{\mathcal{H}}(A) \geq 0$.

Proof. Only points of $\mathcal{X}_{\mathcal{H}}(A)$ contribute to the sum in (3). Multiplying the definition of $\Delta_{\mathcal{H}}(A)$ by m and using (3) gives

$$m\Delta_{\mathcal{H}}(A) = m \sum_{x \in \mathcal{X}_{\mathcal{H}}(A)} r(x) + (m-1)I_{\mathcal{H}}(A) - m|\mathcal{X}_{\mathcal{H}}(A)| - 6 \binom{k}{4}.$$

By (4),

$$6 \binom{k}{4} = 2 \sum_{x \in \mathcal{X}_{\mathcal{H}}(A)} \binom{r(x)}{2} + 2 \sum_{y \in \mathcal{H}} \binom{r(y)}{2}.$$

Substitution and termwise simplification yield

$$\begin{aligned} m(r-1) - 2 \binom{r}{2} &= (r-1)(m-r), \\ (m-1)r - 2 \binom{r}{2} &= r(m-r), \end{aligned}$$

which proves (5). Every term is nonnegative by Lemma 2.1. □

Corollary 3.2 (Coverage and uncovered-locus bounds). *Under the hypotheses of Theorem 3.1,*

$$|\mathcal{X}_{\mathcal{H}}(A)| \leq N(q-1) - \frac{6}{m} \binom{k}{4} - \frac{I_{\mathcal{H}}(A)}{m}, \quad (6)$$

and

$$|\mathcal{U}_{\mathcal{H}}(A)| \geq |V_{\mathcal{H}}(A)| - N(q-1) + \frac{6}{m} \binom{k}{4} + \frac{I_{\mathcal{H}}(A)}{m}. \quad (7)$$

Equality holds in either inequality if and only if

$$r(x) \in \{1, m\} \quad (x \in \mathcal{X}_{\mathcal{H}}(A)), \quad r(y) \in \{0, m\} \quad (y \in \mathcal{H}).$$

Proof. The two inequalities are respectively $\Delta_{\mathcal{H}}(A) \geq 0$ and its rearrangement using $|\mathcal{U}_{\mathcal{H}}(A)| = |V_{\mathcal{H}}(A)| - |\mathcal{X}_{\mathcal{H}}(A)|$. The equality statement follows from the vanishing of every nonnegative term in (5). $\square$

For later use, we record the complete-outside consequence before imposing any geometry on the holes.

Corollary 3.3 (Arbitrary prescribed holes). *Let $|\mathcal{H}| = h$. If A is a k -arc disjoint from $\mathcal{H}$ and complete outside $\mathcal{H}$, then*

$$q^2 + q + 1 - k - h \leq \binom{k}{2}(q-1) - \frac{6}{m} \binom{k}{4} - \frac{I_{\mathcal{H}}(A)}{m}. \quad (8)$$

Proof. Completeness gives $|\mathcal{X}_{\mathcal{H}}(A)| = |V_{\mathcal{H}}(A)| = q^2 + q + 1 - k - h$. Substitute this equality into (6). $\square$

The exact remainder also quantifies near equality.

Corollary 3.4 (Quantitative stability). *Assume $m \geq 3$ and put*

$$\begin{aligned} M &= \{x \in \mathcal{X}_{\mathcal{H}}(A) : 2 \leq r(x) \leq m-1\}, \\ J &= \{y \in \mathcal{H} : 1 \leq r(y) \leq m-1\}. \end{aligned}$$

Then

$$(m-2)|M| + (m-1)|J| \leq m\Delta_{\mathcal{H}}(A). \quad (9)$$

Proof. For $2 \leq r \leq m-1$, $(r-1)(m-r) \geq m-2$. For $1 \leq r \leq m-1$, $r(m-r) \geq m-1$. Apply these estimates to (5). $\square$

Thus small defect forces almost every multiply covered required point to have the maximum possible multiplicity m , while almost every hole point met by a secant also has multiplicity m . This is the precise stability content behind the equality heuristic.

4 Coding interpretation

We record the standard arc-code dictionary because it makes the defect identity more informative. Choose nonzero column representatives of a k -arc $A \subset \text{PG}(2, q)$ and form a $3 \times k$ matrix H_A . For $k \geq 4$, the code

$$C_A = \{c \in \mathbb{F}_q^k : H_A c^T = 0\}$$

is a $[k, k - 3, 4]_q$ MDS code: every three columns are independent, the columns span $\mathbb{F}_q^3$, and any four columns support a dependence. Conversely, a projective parity-check system for a code with these parameters is a k -arc. This correspondence and its coset-weight consequences are classical; see Davydov–Marcugini–Pambianco [8]. We call such a code *projectively GRS* when, after permuting and scaling columns and applying a projective coordinate change, its projective parity-check columns lie on a normal rational curve; in $\text{PG}(2, q)$ this is a nonsingular conic. This is the standard normal-rational-curve/GRS correspondence [18].

Proposition 4.1 (Syndrome form of relative completeness). *Let $s \neq 0$ have projective direction $x = [s] \in \text{PG}(2, q)$. Then:*

- (i) *the minimum weight of a word c with $H_{AC}^\top = s$ is one, two, or three according as $x \in A$, $x \notin A$ lies on a secant of A , or x is ordinarily uncovered;*
- (ii) *if the minimum weight is two, the number of weight-two leaders is exactly $r_A(x)$;*
- (iii) *if the minimum weight is three, the number of weight-three leaders is exactly $\binom{k}{3}$; and*
- (iv) *adjoining x as one more projective parity-check column preserves the MDS property exactly when x is ordinarily uncovered.*

Consequently, A is complete outside a prescribed set $\mathcal{H}$ exactly when all its projective distance-three syndrome directions lie in $\mathcal{H}$ (and $A \cap \mathcal{H} = \emptyset$). These directions are deep-hole directions only when covering radius three has separately been established.

Proof. A weight-one representation is a scalar multiple of a selected column. A weight-two representation uses a unique unordered column pair whose projective span is a secant through x ; independence of the pair makes its two coefficients unique. If neither case occurs, any three columns form a basis and represent s , so the distance is three. For each three-subset the representation is unique, and none of its coefficients can vanish; hence there are exactly $\binom{k}{3}$ leaders. The extension statement is the same three-column independence test. The leader count also follows as the $d = 4$ case of the general farthest-coset formula in Davydov–Marcugini–Pambianco [8, Theorem 7.7]; the deep-hole/MDS-extension formulation is standard in the Reed–Solomon setting as well [13]. $\square$

Thus $r_A(x)$ is not merely an incidence index: it is the exact number of minimum weight-two leaders of every affine syndrome on direction x . All $q - 1$ nonzero scalar multiples of a projective syndrome direction have the same distance and leader count: multiplication by the scalar bijects their coefficient words without changing Hamming support. This is the normalization behind every projective-to-affine factor of $q - 1$ below. The first moment counts leaders, the second counts collisions between distinct leader supports, and Theorem 3.1 is an exact leader-collision remainder identity after prescribed directions are removed. In particular, Corollary 3.3 is also a length obstruction for projective codimension-three MDS codes whose distance-three syndrome directions are confined to the prescribed projective set.

5 Specialization to a conic

Let $\mathcal{C}$ be a nonsingular conic in $\text{PG}(2, q)$, and let $A \subseteq \text{PG}(2, q) \setminus \mathcal{C}$ be a k -arc. Since $|\mathcal{C}| = q + 1$, there are $q^2 - k$ points outside $A \cup \mathcal{C}$. Write

$$I_{\mathcal{C}}(A) = \sum_{y \in \mathcal{C}} r(y) = \sum_{\ell \text{ an } A\text{-secant}} |\ell \cap \mathcal{C}|.$$

Corollary 5.1 (Conic-loss and uncovered-locus inequalities). *For every such arc,*

$$|\mathcal{U}_C(A)| \geq q^2 - k - \binom{k}{2}(q-1) + \frac{6}{m} \binom{k}{4} + \frac{I_C(A)}{m}. \quad (10)$$

If A is $\mathcal{C}$ -complete, then

$$\boxed{q^2 - k \leq \binom{k}{2}(q-1) - \frac{6}{m} \binom{k}{4} - \frac{I_C(A)}{m}.} \quad (11)$$

The first correction in (11) accounts for unavoidable concurrence among secants; the last accounts for secant incidence spent on the exempt conic.

Define

$$L_1(q) = \min \left\{ k \geq 3 : q^2 - k \leq \binom{k}{2}(q-1) \right\},$$

$$L_2(q) = \min \left\{ k \geq 3 : q^2 - k \leq \binom{k}{2}(q-1) - \frac{6}{\lfloor k/2 \rfloor} \binom{k}{4} \right\}.$$

Then

$$\rho_C(q) \geq L_2(q) \geq L_1(q). \quad (12)$$

The definition of $L_2(q)$ involves only integer arithmetic after multiplication by $\lfloor k/2 \rfloor$.

For calculation, the corrected capacity has the parity-dependent forms

$$\binom{k}{2}(q-1) - \frac{6}{\lfloor k/2 \rfloor} \binom{k}{4} = \begin{cases} \frac{k-1}{2}(k(q-1) - (k-2)(k-3)), & k \text{ even,} \\ \frac{k}{2}((k-1)(q-1) - (k-2)(k-3)), & k \text{ odd.} \end{cases} \quad (13)$$

For comparison, some values are

q	5	8	9	11	13	16	17	19
$L_1(q)$	4	5	5	6	6	7	7	7
$L_2(q)$	4	6	6	6	7	8	8	8

Thus the second-moment correction alone proves, for example, that a $\mathcal{C}$ -complete arc in $\text{PG}(2, 16)$ has at least eight points.

Theorem 5.2 (Explicit additive lower bound). *For every prime power q ,*

$$\boxed{\rho_C(q) \geq \sqrt{2q} + \frac{3}{2} - \frac{8}{\sqrt{2q}}.} \quad (14)$$

Consequently,

$$\liminf_{q \rightarrow \infty} (\rho_C(q) - \sqrt{2q}) \geq \frac{3}{2}. \quad (15)$$

Proof. Let k be the size of a $\mathcal{C}$ -complete arc and put $s = \sqrt{2q}$. The elementary first-moment inequality $q^2 - k \leq \binom{k}{2}(q-1)$ implies $k \geq s$ for $q \geq 2$: otherwise $k < s \leq q$, while $k^2 < 2q$ gives $\binom{k}{2} < q$, so its right side is smaller than $q(q-1)$ whereas $q^2 - k \geq q(q-1)$. If $k \geq s+2$, then (14) is immediate. Otherwise write $k = s + a$, so $0 \leq a < 2$ and $s \geq 2$.

Since $m = \lfloor k/2 \rfloor \leq k/2$,

$$\frac{6}{m} \binom{k}{4} \geq \frac{12}{k} \binom{k}{4} = \frac{(k-1)(k-2)(k-3)}{2}.$$

Dropping the nonnegative conic term in (11) therefore gives the necessary inequality

$$q^2 - k \leq \frac{k-1}{2} (k(q-1) - (k-2)(k-3)).$$

After substituting $q = s^2/2$ and $k = s + a$, the right side minus the left side is exactly

$$\frac{2a-3}{4} s^3 + \frac{a^2-7a+10}{4} s^2 + \frac{-3a^2+10a-8}{2} s + \frac{-a^3+5a^2-8a+6}{2}.$$

For $0 \leq a \leq 2$, the last three coefficients are at most $5/2$, 1 , and 3 , respectively. Since $s \geq 2$,

$$\frac{5}{2} s^2 + s + 3 \leq 4s^2,$$

so their total contribution is at most $4s^2$. The displayed difference is nonnegative, and hence

$$0 \leq \frac{2a-3}{4} s^3 + 4s^2.$$

Dividing by s^2 gives $a \geq 3/2 - 8/s$, which is (14). Letting q tend to infinity gives (15). $\square$

Remark 5.3 (The scale of the conic term). Every secant meets $\mathcal{C}$ in at most two points, so

$$0 \leq I_{\mathcal{C}}(A) \leq 2 \binom{k}{2}.$$

At the relevant scale $k \asymp \sqrt{q}$, the term $I_{\mathcal{C}}(A)/m$ is therefore $O(\sqrt{q})$, while the universal overlap correction $6 \binom{k}{4}/m$ is $\Theta(q^{3/2})$. Consequently, improving the lower bound on $I_{\mathcal{C}}(A)$ within (11) can be useful for finely balanced finite cases, but cannot improve the additive constant $3/2$ in Theorem 5.2. Any stronger asymptotic result needs an additional mechanism, not merely a spectral estimate for $I_{\mathcal{C}}(A)$.

6 An upper-bound transfer from complete arcs

Let $t_2(2, q)$ denote the smallest size of an ordinary complete arc in $\text{PG}(2, q)$.

Theorem 6.1 (Projective averaging). *Let $H \subseteq \text{PG}(2, q)$ and let B be an ordinary complete b -arc. If*

$$b|H| < q^2 + q + 1, \tag{16}$$

then some projective image of B is disjoint from H and complete outside H .

Proof. Choose a uniformly random projectivity $g \in \text{PGL}(3, q)$. Point transitivity gives

$$\mathbb{E}|gB \cap H| = \frac{b|H|}{q^2 + q + 1} < 1.$$

Since the intersection size is a nonnegative integer, it is zero for some g . Projectivities preserve ordinary completeness, so gB is complete outside H . $\square$

Corollary 6.2 (Conic transfer). *If $t_2(2, q) \leq q$, then*

$$\rho_{\mathcal{C}}(q) \leq t_2(2, q). \quad (17)$$

More generally, every complete b -arc with $b \leq q$ has a projective image disjoint from the prescribed conic.

Proof. Take $H = \mathcal{C}$, so $|H| = q + 1$. For integral $b \leq q$ one has $b(q + 1) < q^2 + q + 1$, and Theorem 6.1 applies. $\square$

Kim and Vu proved that, for all sufficiently large q , every projective plane of order q has a complete arc of size at most $\sqrt{q}(\log q)^c$ for an absolute constant c [12]. This is less than q for large q , so Corollary 6.2 gives the following immediate consequence.

Corollary 6.3. *For some absolute constant c and all sufficiently large prime powers q ,*

$$\rho_{\mathcal{C}}(q) \leq \sqrt{q}(\log q)^c.$$

The transfer does not exploit the conic and is unlikely to be sharp. It nevertheless supplies a uniform upper bound without having to preserve the arc condition during a separate random covering construction.

7 Even characteristic and the nucleus

Assume q is even and let ν be the nucleus of $\mathcal{C}$. The nucleus is outside $\mathcal{C}$, all tangents to $\mathcal{C}$ pass through ν , and every line through ν is one of these $q + 1$ tangents.

Proposition 7.1 (The nucleus belongs to the arc). *Let A be a k -arc disjoint from $\mathcal{C}$ with $\nu \in A$. Exactly $k - 1$ A -secants are tangent to $\mathcal{C}$. In particular,*

$$I_{\mathcal{C}}(A) \geq k - 1, \quad I_{\mathcal{C}}(A) \equiv k - 1 \pmod{2}.$$

If A is $\mathcal{C}$ -complete, then

$$q^2 - k \leq \binom{k}{2}(q - 1) - \frac{6}{m} \binom{k}{4} - \frac{k - 1}{m}. \quad (18)$$

Proof. For every $a \in A \setminus \{\nu\}$, the line νa is tangent to $\mathcal{C}$, giving $k - 1$ tangent secants. No line joining two points of $A \setminus \{\nu\}$ can be tangent: every tangent contains ν , which would place three points of A on one line. Hence these are exactly the tangent A -secants.

Every tangent contributes one to $I_{\mathcal{C}}(A)$, while every other line contributes zero or two. This proves the lower bound and parity statement. Inequality (18) follows from Corollary 5.1. $\square$

Proposition 7.2 (The nucleus lies outside the arc). *Let A be a $\mathcal{C}$ -complete arc with $\nu \notin A$. Then*

$$1 \leq r(\nu) \leq m,$$

the tangent A -secants are exactly the secants through ν , and

$$I_{\mathcal{C}}(A) \equiv r(\nu) \pmod{2}, \quad I_{\mathcal{C}}(A) \geq r(\nu) \geq 1.$$

Consequently,

$$q^2 - k \leq \binom{k}{2}(q - 1) - \frac{6}{m} \binom{k}{4} - \frac{1}{m}. \quad (19)$$

Proof. The nucleus is a required point outside $A \cup \mathcal{C}$, so relative completeness gives $r(\nu) \geq 1$; Lemma 2.1 gives the upper bound. A line is tangent to $\mathcal{C}$ exactly when it passes through ν . Thus the number of tangent A -secants is $r(\nu)$. Reducing the sum of the intersection sizes $|\ell \cap \mathcal{C}|$ modulo two gives $I_{\mathcal{C}}(A) \equiv r(\nu) \pmod{2}$. The remaining claims follow from Corollary 5.1. $\square$

Corollary 7.3 (Universal even-characteristic loss). *Every $\mathcal{C}$ -complete arc in even characteristic satisfies*

$$I_{\mathcal{C}}(A) \geq 1.$$

Consequently, inequality (11) can always be strengthened by replacing its final term by at least $-1/m$.

Proof. If the nucleus belongs to A , Proposition 7.1 gives $I_{\mathcal{C}}(A) \geq k - 1 \geq 1$. Otherwise Proposition 7.2 gives $I_{\mathcal{C}}(A) \geq r(\nu) \geq 1$. $\square$

These two cases are useful structural reductions, but Remark 5.3 limits what the incidence term alone can accomplish. For example, at $(q, k) = (16, 8)$ inequality (11) would be contradicted only by $I_{\mathcal{C}}(A) > 268$, whereas the universal upper bound is $I_{\mathcal{C}}(A) \leq 56$. Thus the exact determination of $\rho_{\mathcal{C}}(16)$ below requires information beyond a lower bound on $I_{\mathcal{C}}(A)$ inserted into the present inequality.

8 Verified finite-field examples

The verified values at $q = 8, 9, 11$ use the lower bound $L_2(q)$ and explicit coordinates. At $q = 16$, the same ingredients give $8 \leq \rho_{\mathcal{C}}(16) \leq 9$; a checked projective classification excludes eight.

Complete exteriority supplies a useful foil. For every four-arc, the six secants have three repeated off-arc intersection points, and hence

$$|U(A)| = (q^2 + q + 1 - 4) - (6(q - 1) - 3) = (q - 2)(q - 3).$$

Blokhuis–Seress–Wilbrink’s $q = 7$ exterior four-arc therefore has $|U(A)| = 20 > 8 = |\mathcal{C}(\mathbb{F}_7)|$: its conic lies strictly inside the uncovered locus [6, pp. 143, 146]. The $q = 11$ witness below is exceptional because the reverse inclusion also holds.

8.1 Evaluation obstruction

The algebraic rejection used in that classification is an instance of the following elementary linear-algebra observation. Linear systems of quadrics containing arcs are an established tool; see Glynn [10] and Ball [4]. Fix nonzero vector representatives of the projective points under consideration. If V is a finite-dimensional linear system of homogeneous forms, write $\text{ev}_x \in V^*$ for evaluation at the representative of x . The zero or nonzero status of this evaluation is independent of the chosen representative.

Indeed, if an arc is complete outside a conic, then its ordinary-uncovered locus lies in the zero set of the conic equation while every selected point avoids that zero set. The next lemma isolates exactly the linear-algebraic obstruction used to rule this out at $q = 16$.

Lemma 8.1 (Uncovered evaluation obstruction). *Let A and U be finite sets of projective points. There is no nonzero $f \in V$ such that $U \subseteq Z(f)$ and $A \cap Z(f) = \emptyset$ if either of the following holds:*

1. *the evaluation map $f \mapsto (\text{ev}_u(f))_{u \in U}$ is injective;*

2. for some $a \in A$, the functional ev_a belongs to the span of $\{\text{ev}_u : u \in U\}$.

Proof. If f vanishes on U , injectivity in the first case gives $f = 0$. In the second case, applying a spanning relation for ev_a to f gives $f(a) = 0$, contrary to $A \cap Z(f) = \emptyset$. $\square$

For the full system of degree- d forms, the evaluation functionals are the degree- d Veronese images of the points, up to nonzero scalar. Thus the first alternative says that the uncovered Veronese images span the dual coefficient space; the second supplies a selected Veronese image already in their span. In particular, for the full degree- d system it is enough to exhibit $\binom{d+2}{2}$ uncovered points whose Veronese evaluation matrix is invertible. This formulation is not specific to conics.

The following sharp form is useful when a zero locus must avoid several selected points simultaneously. The vector-space covering threshold is classical; see, for example, Khare [14].

Proposition 8.2 (Finite-field evaluation dichotomy). *Let W be a finite-dimensional vector space over $\mathbb{F}_q$, let $\nu : X \rightarrow W$ be any feature map, and let $U, A \subseteq X$ be finite with $|A| \leq q$. There is a linear form $f \in W^*$ such that*

$$f(\nu(u)) = 0 \quad (u \in U), \quad f \neq 0, \quad f(\nu(a)) \neq 0 \quad (a \in A)$$

if and only if

$$\langle \nu(U) \rangle \neq W \quad \text{and} \quad \nu(a) \notin \langle \nu(U) \rangle \quad (a \in A).$$

The uniform threshold is sharp: $q + 1$ distinct hyperplanes through a common codimension-two subspace can cover W .

Proof. Put $S = \langle \nu(U) \rangle$. Necessity follows by applying f to the stated span relations. Conversely, $S \neq W$ makes the annihilator $S^\perp \subseteq W^*$ nonzero. For each $a \in A$, the condition $\nu(a) \notin S$ says that evaluation at $\nu(a)$ cuts out a proper hyperplane of $S^\perp$. At most q proper hyperplanes cannot cover a nonzero vector space over $\mathbb{F}_q$: their union has size at most $1 + q(|S^\perp|/q - 1) < |S^\perp|$. Any form outside their union has the required properties. For sharpness, quotient by a common codimension-two subspace and use all $q + 1$ lines in $\mathbb{F}_q^2$. $\square$

Taking ν to be the degree- d Veronese map gives the same exact dichotomy for homogeneous degree- d forms. Thus Lemma 8.1 is the one-selected-point exclusion mechanism, while Proposition 8.2 determines exactly when a single form can avoid as many as q selected points.

8.2 Small values and the $q = 16$ classification

Proposition 8.3. *For the standard conic*

$$\mathcal{C} : XZ = Y^2$$

in the indicated projective planes,

$$\rho_{\mathcal{C}}(8) = 6, \quad \rho_{\mathcal{C}}(9) = 6, \quad \rho_{\mathcal{C}}(11) = 6.$$

Proof. Equation (12) gives

$$L_2(8) = L_2(9) = L_2(11) = 6.$$

Appendix C lists $\mathcal{C}$ -complete arcs of size six in each plane. Their verification is described below. $\square$

Theorem 8.4 (Quadratic avoidance over $\mathbb{F}_{16}$). *Let A be any eight-arc in $\text{PG}(2, 16)$, and let $U(A)$ be the points outside A that lie on no secant of A . There is no nonzero homogeneous quadratic form Q such that*

$$U(A) \subseteq Z(Q) \quad \text{and} \quad A \cap Z(Q) = \emptyset.$$

The assertion includes singular quadratic forms.

Proof. The proof has three steps. First, canonical augmentation reduces every eight-arc to one of 2633 projective classes. Starting from a normalized four-frame, the certified lists contain respectively 4, 61, 454, and 2633 classes at sizes five through eight. Coverage certificates check that every legal extension occurs in the next list, while explicit projective equivalences and frame reduction transport an arbitrary eight-arc to a listed leaf. Thus the exhaustiveness claim does not depend on trusting the generator's canonical labels. The final count agrees with the published classification of Al-Seraji–Al-Ogali [2, Theorem 3.8.1].

Second, for 2630 of the classified leaves, six points of $U(A)$ give an invertible 6×6 evaluation matrix, giving the first alternative of Lemma 8.1. In each of the remaining three leaves the evaluation rows have rank five, but an explicit linear combination places the evaluation functional of a point of A in their span, giving the second alternative.

Third, the obstruction is projectively invariant. If gA is a normalized representative, replace Q by $Q \circ g^{-1}$. This is nonzero because g is invertible, its zero set is $gZ(Q)$, and projectivities carry secants to secants, so $U(gA) = gU(A)$. Thus a quadratic as in the statement would contradict the obstruction for the normalized leaf. Appendix A records the independent regeneration and formal trust boundary. $\square$

Corollary 8.5 (Exact relative value over $\mathbb{F}_{16}$). *For every nonsingular conic $\mathcal{C} \subseteq \text{PG}(2, 16)$,*

$$\rho_{\mathcal{C}}(16) = 9.$$

Proof. Equation (12) gives $L_2(16) = 8$. If a $\mathcal{C}$ -complete eight-arc existed, its ordinary-uncovered locus would lie in the zero set of the nonzero quadratic defining $\mathcal{C}$, while the arc is disjoint from that zero set. This contradicts Theorem 8.4. The nine-point witness in Appendix C gives the matching upper bound. $\square$

The published ordinary classification records arc counts, stabilizers, secant-index distributions, and conic-contained classes. The additional datum used here is the uncovered-locus evaluation split: 2630 full-rank leaves and three rank-five leaves forced to meet the arc.

Remark 8.6 (Anatomy of the three exceptional leaves). Using the binary $\mathbb{F}_{16}$ encoding from Appendix C and coefficient order $(X^2, Y^2, Z^2, XY, XZ, YZ)$, the one-dimensional kernels of the three exceptional evaluation matrices are generated by

$$(1, 1, 1, 1, 1, 0), \quad (0, 0, 0, 5, 4, 1), \quad (2, 1, 1, 5, 5, 1).$$

The first and third forms split over $\mathbb{F}_{16}$ as

$$(X + 6Y + 6Z)(X + 7Y + 7Z) \quad \text{and} \quad 2(X + 4Y + 15Z)(X + 15Y + 4Z),$$

respectively, and each union of lines contains two selected points. The middle form is nonsingular and contains seven of the eight selected points. Thus the exceptional leaves are not merely unexplained rank defects: two are reducible quadratic configurations, while the third is an arc differing by one point from a seven-point subset of a conic. These descriptions follow by direct field arithmetic and are not used to establish classification completeness.

8.3 The $q = 11$ coding application

For reference, the verified incidence statistics are

q	$ A $	$\binom{ A }{2}$	$q^2 - A $	$I_C(A)$	$\max_{x \notin A} r(x)$
8	6	15	58	16	3
9	6	15	75	13	3
11	6	15	115	0	3
16	9	36	247	32	4

The row at $q = 11$ has a stronger geometric interpretation. Since $I_C(A) = 0$, none of the 15 secants meets $\mathcal{C}$: every secant of the witness is exterior to the conic. If N_i denotes the number of required points of index i , relative completeness and the maximum-index bound give only $i = 1, 2, 3$, while the two moment equations give

$$N_1 + N_2 + N_3 = 115, \quad N_1 + 2N_2 + 3N_3 = 150, \quad N_2 + 3N_3 = 45.$$

Consequently $(N_1, N_2, N_3) = (90, 15, 10)$.

Proposition 8.7 (The $q = 11$ code and coset spectrum). *Let A be the six-point witness in Appendix C, and let C_A be its parity-check code. Then:*

- (i) C_A is a projectively non-GRS $[6, 3, 4]_{11}$ MDS code of covering radius three, and its projective deep-hole syndrome locus is exactly the twelve-point conic $\mathcal{C}$;
- (ii) the affine cosets of distances 0, 1, 2, 3 number $(1, 60, 1150, 120)$, and the distance-two cosets split as $(900, 150, 100)$ according as they have one, two, or three minimum-weight leaders.

Proof. Every witness triple has nonzero determinant and the six columns span $\mathbb{F}_{11}^3$, giving the MDS parameters. The 6×6 evaluation matrix of the quadratic monomials

$$X^2, XY, XZ, Y^2, YZ, Z^2$$

is nonsingular, so no conic contains all six projective columns; by the preceding definition and the classical normal-rational-curve dictionary, the code is not projectively GRS. Direct finite arithmetic gives the secant-index spectrum

$$0 : 12, \quad 1 : 90, \quad 2 : 15, \quad 3 : 10,$$

where index zero is precisely the conic. Multiplying each nonzero projective direction by its ten affine scalars gives all nonzero affine syndromes bijectively. Proposition 4.1 identifies the determinant-zero witness pairs with weight-two leader supports; uniqueness on an independent support and scalar multiplication then give the claimed coset and leader counts. The index-zero directions give the covering radius. $\square$

Proposition 8.8 (The $q = 11$ extension complex). *For the same witness:*

- (i) the simultaneous one-column MDS extensions form the independence complex of the icosahedral graph, with independence polynomial

$$1 + 12t + 36t^2 + 20t^3;$$

exactly six two-column and twenty three-column extensions are maximal, giving six complete eight-arcs and twenty complete nine-arcs containing the fixed seed, and there is no ten-arc extension;

(ii) *each witness column colours five pairwise disjoint edges of the icosahedron, missing an antipodal pair, and the six colour classes partition its thirty edges. Adding the six antipodal edges turns them into six perfect matchings of the augmented graph.*

Proof. For simultaneous extensions, two conic columns conflict exactly when their chord contains a witness column. No three conic columns are collinear, so there is no additional triple obstruction and valid extension sets are exactly graph-independent sets. Exhausting the 2^{12} subsets gives the displayed polynomial and maximal-set counts. The same determinant table partitions the thirty conflict edges into the six stated matchings. Relative completeness confines every further extension point to the conic, so maximality in the independence complex upgrades to ordinary completeness. The six missing antipodal edges are distinct; adjoining the appropriate one to each colour class gives six disjoint perfect matchings of the augmented graph. Appendix A records the two independent implementations and the formal checks. $\square$

The six-set is a classical Clebsch hexagon with icosahedral A_5 symmetry; see Edge, Dye, and Storme–Van Maldeghem for the classical configuration and orbit geometry [7, 9, 17]. More precisely, Dye’s edge criterion at $q = 11$ and the complete-exterior-set formulation of Blokhuis–Seress–Wilbrink give $\mathcal{C}(\mathbb{F}_{11}) \subseteq U(A)$ [9, discussion preceding Theorem 6, p. 281] [6, pp. 143, 146]. Dye’s ten Brianchon concurrences and the chord-defect identity then give $|U(A)| = 12$, hence equality. Proposition 8.7 combines these classical inputs with the syndrome counts, while Proposition 8.8 records the simultaneous-extension calculation.

Appendix A records the computational and formal verification boundary. The auxiliary game-theoretic consequence of the $q = 11$ extension graph is separated in Appendix B.

9 Further questions

The elementary theory leaves several concrete problems.

Problem 9.1. Determine whether the uncovered-quadratic-rank obstruction used at $q = 16$ extends to other even orders or gives useful lower bounds on the size of a $\mathcal{C}$ -complete arc beyond the scalar defect inequality.

Problem 9.2. Develop the higher-dimensional analogue for caps complete outside a prescribed quadric in $\text{PG}(n, q)$. The evaluation-avoidance lemma already applies to arbitrary feature maps and every Veronese degree; the missing ingredients are the appropriate cap secant moments and useful examples in dimensions at least three.

Problem 9.3. Determine whether the exact defect remainder yields new lower bounds for arc-constrained saturating sets or the corresponding covering codes.

Problem 9.4. Determine whether small-order examples beyond $q = 11$ produce further recognizable simultaneous-extension complexes.

Problem 9.5. Construct $\mathcal{C}$ -complete arcs of size $O(\sqrt{q})$, or prove that this is impossible for an infinite family. Theorem 6.1 currently supplies only the general complete-arc bound $O(\sqrt{q}(\log q)^c)$.

Problem 9.6. Classify equality and bounded-defect configurations in Theorem 3.1. The equality pattern requires all nontrivial secant concurrence to occur at points of maximum possible index, a severe combinatorial restriction.

Problem 9.7. Determine whether the conic stabilizer or its coherent configuration yields useful finite-order restrictions not reducible to a bound on $I_C(A)$. Remark 5.3 shows that an estimate for $I_C(A)$ alone cannot improve the asymptotic constant obtained here.

The central distinction is between universal overlap, already measured by the classical second index equation, and genuinely conic-specific geometry. The former gives the lower bound in this paper. Progress beyond it will probably require an explicit construction, an algebraic obstruction, or a classification of near-extremal secant arrangements rather than another application of the same incidence budget.

A Computational and formal verification

The supplementary Python program `verify_relative_conic_arcs.py` performs exact finite-field arithmetic and, for each witness:

- (a) enumerates and normalizes all $q^2 + q + 1$ projective points;
- (b) checks that $XZ = Y^2$ contains exactly $q + 1$ points;
- (c) verifies disjointness from the conic and that every pair-line contains exactly two selected points;
- (d) verifies that every point outside $A \cup C$ lies on a selected pair-line; and
- (e) independently checks both equations in Proposition 2.2.

Its SHA-256 digest is

`e9508958d604e68c6c3d09fd3afadfaa8a3126508a51f1dfa993e7a7aed5d36a.`

The verifier is deliberately simpler than a search program: it certifies the listed upper-bound witnesses but makes no claim about search completeness.

Two further implementations independently regenerate Propositions 8.7 and 8.8. Their separately written source files are

`check_q11_structure.py`, `check_q11_structure.cpp`.

Their SHA-256 digests are

`0abe909c9aadce0db4c75f296c8de25e929dd1065c906996da8dec017e534d69,`
`1753674172d48f1d056d350e30baa9eb67de0810c84a96da0440947768ae041c.`

They agree on the projective, syndrome, chord, and group-action data. Both include transformed-coordinate invariance checks; the adversarial controls replace one witness point and mutate a group generator, respectively, and reject the exceptional conclusions. The small-field hyperplane checker `check_evaluation_dichotomy.py`, with digest

`6ed309bd2461ce9998cbd3bcaa5396379e6973b7503ce2dcdfb32c9806386566,`

exhausts every family of at most q projective hyperplanes for $q = 2, 3, 5$ and dimensions one through three, and checks the sharp $q + 1$ plane cover.

The exact-classification generator `search_rhoc16.cpp` has SHA-256 digest

`589af8430e94b4c9c23ce895e6d32d2b3b5b9b387b1fb23ed6d3875cdee39031.`

Its frozen report `search_rhoc16_output.txt` has digest

6989079b5cb64b0e57d5c42b872093fff99f861300b8fbb909daef450c15cc63.

The generator emits local Lean certificates; its own assertions and canonical labels are not part of the trusted proof boundary.

The companion Lean library `lean/RelativeConicArcs/` formalizes the incidence, defect, conic, asymptotic, averaging, nucleus, evaluation-dichotomy, syndrome, and coding arguments. Its finite certificates check the code parameters, deep-hole locus, leader distributions, quadratic obstruction, chord partition, extension complex, and the exhaustiveness and projective transport used at $q = 16$. The generic Boolean checker inspects conic disjointness, the arc condition, and coverage of the canonical representatives $[1 : y : z]$, $[0 : 1 : z]$, and $[0 : 0 : 1]$; a proved soundness theorem transports acceptance to relative completeness. The concrete checks use kernel reduction rather than `native_decide`. The accompanying trust manifest records the theorem map and axiom profiles. In particular, the finite results do not depend on the external Kim–Vu hypothesis used for the conditional asymptotic upper bound.

B Auxiliary residual-game structure

After the six $q = 11$ witness points are occupied, all twelve points of $\mathcal{C}$ remain legal. Join two conic points when their chord contains a witness point. The resulting 12-vertex, 30-edge, 5-regular conflict graph is the icosahedral graph. Its antipodal involution is fixed-point-free, preserves adjacency, and pairs only nonadjacent vertices, so the copycat strategy makes the corresponding independent-set building position a P-position. The formal development checks the localization from the projective cap game to this graph and hence the actual six-point projective position. It also checks that the displayed $q = 9$ six-arc is terminal.

C Witness coordinates and field encodings

All coordinates are homogeneous and use the standard conic $XZ = Y^2$.

For $q = 8$, write $\mathbb{F}_8 = \mathbb{F}_2[\alpha]/(\alpha^3 + \alpha + 1)$ and encode $a_0 + a_1\alpha + a_2\alpha^2$ by the binary integer $a_0 + 2a_1 + 4a_2$. A witness is

$$(0, 1, 1), (0, 1, 2), (1, 0, 1), \\ (1, 0, 2), (1, 1, 2), (1, 1, 6).$$

For $q = 9$, write $\mathbb{F}_9 = \mathbb{F}_3[\alpha]/(\alpha^2 + 1)$ and encode $a_0 + a_1\alpha$ by $a_0 + 3a_1$. A witness is

$$(1, 0, 4), (1, 0, 5), (1, 1, 0), \\ (1, 1, 2), (1, 2, 3), (1, 2, 4).$$

For $q = 11$, coordinates lie in the prime field. A witness is

$$(1, 10, 0), (1, 9, 1), (1, 4, 7), \\ (1, 8, 5), (0, 1, 4), (1, 1, 7).$$

For $q = 16$, write $\mathbb{F}_{16} = \mathbb{F}_2[\alpha]/(\alpha^4 + \alpha + 1)$ and use binary coefficient encoding. A witness of size nine is

$$(1, 5, 14), (1, 5, 3), (1, 9, 2), (1, 0, 9), (1, 13, 13), \\ (1, 6, 10), (1, 3, 9), (0, 1, 11), (1, 10, 2).$$

References

- [1] S. Alabdullah and J. W. P. Hirschfeld, A new lower bound for the smallest complete (k, n) -arc in $\text{PG}(2, q)$, *Designs, Codes and Cryptography* **87** (2019), 679–683. doi:10.1007/s10623-018-00592-8.
- [2] N. A. M. Al-Seraji and R. A. B. Al-Ogali, Classification of arcs in finite projective plane of order sixteen, *Al-Mustansiriyah Journal of Science* **29**(1) (2018), 138–149. doi:10.23851/mjs.v29i1.184.
- [3] S. Ball, On small complete arcs in a finite plane, *Discrete Mathematics* **174** (1997), 29–34. doi:10.1016/S0012-365X(96)00315-9.
- [4] S. Ball, On arcs and quadrics, in *Arithmetic of Finite Fields—WAIFI 2016*, Lecture Notes in Computer Science **10064**, Springer, 2017, 95–102. doi:10.1007/978-3-319-55227-9_7.
- [5] D. Bartoli, A. A. Davydov, S. Marcugini, and F. Pambianco, On the smallest size of an almost complete subset of a conic in $\text{PG}(2, q)$ and extendability of Reed–Solomon codes, *Problems of Information Transmission* **54** (2018), 101–115. doi:10.1134/S0032946018020011.
- [6] A. Blokhuis, A. Seress, and H. A. Wilbrink, Characterization of complete exterior sets of conics, *Combinatorica* **12**(2) (1992), 143–147. doi:10.1007/BF01204717.
- [7] W. L. Edge, Conics and orthogonal projectivities in a finite plane, *Canadian Journal of Mathematics* **8** (1956), 362–382.
- [8] A. A. Davydov, S. Marcugini, and F. Pambianco, On the weight distribution of the cosets of MDS codes, *Advances in Mathematics of Communications* **17**(5) (2023), 1115–1138. doi:10.3934/amc.2021042. arXiv:2101.12722v2. (Numbered results are cited by their arXiv v2 numbering.)
- [9] R. H. Dye, Hexagons, conics, A_5 and $\text{PSL}_2(K)$, *Journal of the London Mathematical Society* (2) **44** (1991), 270–286. doi:10.1112/jlms/s2-44.2.270.
- [10] D. G. Glynn, On the construction of arcs using quadrics, *Australasian Journal of Combinatorics* **9** (1994), 3–19.
- [11] J. W. P. Hirschfeld, *Projective Geometries over Finite Fields*, second edition, Oxford University Press, 1998.
- [12] J. H. Kim and V. H. Vu, Small complete arcs in projective planes, *Combinatorica* **23** (2003), 311–363. doi:10.1007/s00493-003-0024-1.
- [13] K. Kaipa, Deep holes and MDS extensions of Reed–Solomon codes, *IEEE Transactions on Information Theory* **63** (2017), 4940–4948.
- [14] A. Khare, Vector spaces as unions of proper subspaces, *Linear Algebra and its Applications* **431** (2009), 1681–1686. doi:10.1016/j.laa.2009.06.001.
- [15] Z. L. Nagy, Saturating sets in projective planes and hypergraph covers, *Discrete Mathematics* **341** (2018), 1078–1083. doi:10.1016/j.disc.2018.01.011.
- [16] S.-L. Ng and P. R. Wild, On k -arcs covering a line in finite projective planes, *Ars Combinatoria* **58** (2001), 289–300.

- [17] L. Storme and H. Van Maldeghem, Primitive arcs in $\text{PG}(2, q)$, *Journal of Combinatorial Theory, Series A* **69** (1995), 200–216. doi:10.1016/0097-3165(95)90051-9.
- [18] L. Storme and J. A. Thas, Generalized Reed–Solomon codes and normal rational curves: an improvement of results by Seroussi and Roth, in *Advances in Finite Geometries and Designs*, Oxford University Press, 1991, 369–389. doi:10.1093/oso/9780198535928.003.0031.