

Clebsch series figure set

This document previews the reusable figures of `papers/clebsch-series-figures`. It is a proof sheet, not a manuscript: each figure is defined in its own `\input`-able file and is meant to be placed by a host paper with its own caption. See `README.md` for the input recipe and for the list of edges that encode unproved claims.

D1 — the series map

The five-paper map, with Paper III on top descending the golden sign into the central torsor, Papers I and II joined to it by horizontal recognition passages, and Papers IV and V below. Two conventions carry meaning. The Paper IV edge is *dashed*: no theorem carries its reconstructed plane to the oriented cubic torsor, so the bridge is an open obligation rather than an established arrow. The Paper V node states an *intent*: that paper is unwritten and the Rosetta theorem is not proved.




Figure 1: The series map with no node emphasised.

D1 — the five “you are here” variants

Each numbered paper’s forward version carries the same map with its own node emphasised by a heavier rule and a light tint.




Figure 2: \ClebschSeriesMap{I} — for Paper I.




Figure 3: \ClebschSeriesMap{II} — for Paper II.




Figure 4: $\backslash\text{ClebschSeriesMap}\{\text{III}\}$ — for Paper III.




Figure 5: $\backslash\text{ClebschSeriesMap}\{\text{IV}\}$ — for Paper IV.





Figure 6: $\backslash\text{ClebschSeriesMap}\{V\}$ — for Paper V.

D2 — the recognition template

The programme is a template with instances, not a study of one object. The shape is drawn once; the table instantiates its two slots.




	<i>rich object</i>	<i>coarse observation</i>
Paper I	the Clebsch code, a six-arc in $\text{PG}(2, 11)$	its projective deep-hole syndrome locus
Paper II	the icosahedral conic matching and its two sheets	a two-valued strength-two quadratic trade
Paper III	the order-six conference operator C	four-local principal-minor data (the aligned four-sets)
Paper IV	the marked conic plane $\text{PG}(2, 13)$	the minimum-word layer of its passant code
AME paper	a stabilizer $\text{AME}(2m, q)$ state	its product-unitary symmetries

Figure 7: One shape, five instances. The rich object is forgotten down to a coarse observation, and a theorem of the named paper recovers it.

D3 — the quantifier ladder

Each bar is a theorem’s quantifier range. Where the theorem’s answer is a single exceptional object, it is marked as a distinguished point at the end of the general range. The visual argument is general hypothesis, exceptional conclusion.



- a filled disc marks a theorem whose answer is a single exceptional object, sitting at the end of the general range; a bar without one has a conclusion uniform across the whole range.

Figure 8: Quantifier ranges of the portfolio’s general theorems.

D4 — the four shadows, billed by type

A redesign of Paper III's four-shadow figure. The four descriptions are not four coincidences of equal standing, and the diagram now separates three registers.




Figure 9: The four descriptions of the marked cubic, separated by what kind of statement each agreement is.

D5 — the determinantal ladder for a Seidel matrix

The reconstruction theorem deliberately consults the weakest rung.




Figure 10: Principal minors of a Seidel matrix by order, and which rung the reconstruction theorem uses.

D6 — Paper III's causal spine




Figure 11: From the incidence cover to the operator's shadows, with the proved recognition return underneath.