

Frame transport for framed formal monodromy:
the lemma, its hypotheses, and the receiver problem
Third version, reporting the source checks and correcting the second

Memo for a reader of *Irrationality of Cubic Threefolds after One Stabilization*

15 August 2026

Abstract

Proposition 4.7 of the draft transports framed formal monodromy across the comparison isomorphisms of Iritani and Iritani–Koto, and the transport step itself is asserted in one sentence. This memo states and proves the missing lemma over a differential field, with its hypotheses made explicit; shows that those hypotheses are jointly unsatisfiable over the coefficient receiver the draft builds, so that the lemma as proved has no verified instance in the application; corrects the receiver; and reduces the remaining question to one inequality that depends on a normalization the draft is free to choose. The inequality fails under the draft’s current instruction to take the separating weight parameter large, and holds at the cubic endpoint under the smallest admissible choice. Section 4 examines an iterated receiver that would satisfy the inequality trivially, checks it against the sources, and finds that it needs a semipositivity hypothesis on the ambient variety which weak factorization does not supply.

The second version of this memo proposed the framing theory of Hinault, Yu, Zhang and Zhang as the probable repair, on the ground that its decomposition gauge is regular in the loop coordinate. Three compatibility checks against the pinned sources have since been run, and the route does not survive them; Section 5 now reports what those checks found. Two consequences are worth more than the route was. First, no receiver is needed for the comparison isomorphisms at all: Iritani’s Ψ and Iritani–Koto’s Φ are polynomial in the loop coordinate with inverses in the same ring, so the transport theorem below applies to them with $\Gamma = 0$. Second, the diagnosis of the second version was wrong — the pro-Laurent bulk gauge is not the draft’s own choice but Iritani–Koto’s own fundamental solution, which the sources meet and handle by Birkhoff factorization. What is left is a single statement about one bulk displacement, recorded in Section 5. Section 6 then attacks that statement directly through the flatness identities, and gets far enough to prove that multiplicity-one blocks carry no framed monodromy, to write down how a block residue evolves under bulk motion, and to reduce what remains to one solvability condition on the coalesced blocks. Section 9 lists what the earlier versions got wrong; readers of those versions should start there.

1 The assertion at issue

References are to Section 4 of the draft, *Framed formal monodromy*. I name statements by their semantic labels as well as by number, because the edits proposed in Section 8 renumber the section: Definition 4.1 (`def:framed-sixth-multiplicity`), Definition 4.2 (`def:pro-laurent-gauge-group`), Lemma 4.3 (`lem:pv-base-change`), Lemma 4.5 (`lem:formal-base-shift`), Proposition 4.7 (`prop:framed-operations`), Lemma 4.9 (`lem:divisor-tagging`).

Lemma 4.3 ends with the sentence

If the gauge uses only integral powers of z at every finite level, this conjugacy preserves the original- z -disc frame,

whose proof is the single sentence “the deck action fixes every integral power of z at each finite level, so the compatible gauge is single-valued for the framed turn.” Lemma 4.5 then *defines*

$$M^{\text{bulk}} = G M_{f,V}^{\text{small,shifted}} G^{-1} \quad (4.1c)$$

and Proposition 4.7 concludes that “each center target summand has the framed monodromy of its specialized small connection.” The conclusion does not follow from the display: conjugation preserves a characteristic polynomial for trivial reasons, but says nothing about whether the conjugated matrix is the intrinsic turn operator on horizontal solutions of the gauged module. What is missing is a transport lemma together with an algebra in which the turn acts on solutions.

2 The lemma over a differential field

Notation for Exp_V , $K = \mathbf{C} \oplus V$ and Ω_V is the draft’s, from (4.0) and (4.0a). Recall that Exp_V is defined on $(K, +)$, not on Ω_V .

The ground field

Let Γ be a totally ordered abelian group and let

$$\mathcal{H} = \Omega_V((\Gamma \oplus \mathbf{Z}e_z))$$

be the Hahn field of series with well-ordered support and coefficients in Ω_V , the value group ordered lexicographically with the e_z -coordinate *last*, as in the draft. Write $z = x^{e_z}$ and let θ multiply the coefficient at (γ, k) by k ; this is a derivation because k is additive on the value group, and $\theta z = z$. Its constants are

$$\mathcal{C} := \mathcal{H}^{\theta=0} = \Omega_V((\Gamma)).$$

Taking $\Gamma = 0$ gives $\mathcal{H} = \Omega_V((z))$. For $e \geq 1$ put $\mathcal{H}_e = \Omega_V((\Gamma \oplus \frac{1}{e}\mathbf{Z}e_z))$, a degree- e extension containing $w = x^{e_z/e}$, and $\theta_w = e\theta$, so $\theta_w w = w$. All residues below are taken with respect to θ_w on the ramified side; the draft’s $M_{\text{RS},V}$ uses the same convention, and the two differ by the factor e if one is careless.

The solution algebra and the turn

Definition 1. Assume the exponential factors under consideration have *constant* coefficients, that is $\varphi \in w^{-1}\Omega_V[w^{-1}]$. The *universal formal solution algebra* $\mathcal{U}_e$ is the $\mathcal{H}_e$ -algebra generated by symbols

$$w^\rho \ (\rho \in K), \quad \ell, \quad E(\varphi),$$

subject to $w^\rho w^{\rho'} = w^{\rho+\rho'}$, w^n = the element w^n of $\mathcal{H}_e$ for $n \in \mathbf{Z}$, $E(\varphi)E(\varphi') = E(\varphi + \varphi')$, $E(0) = 1$, with

$$\theta_w(w^\rho) = \rho w^\rho, \quad \theta_w(\ell) = 1, \quad \theta_w(E(\varphi)) = \theta_w(\varphi)E(\varphi).$$

The two hypotheses buried in that definition — constant coefficients in φ , and exponents ρ in K rather than in Ω_V — are not bookkeeping. The first is false for the modules the draft transports and is discussed at Remark 5; the second is forced because Exp_V is defined only on K , so a module whose residues leave K needs Exp rebuilt over a larger field.

Lemma 2 (Structure). *Fix representatives $\Lambda \subset K$ for $K/\mathbf{Z}$ with $0 \in \Lambda$. Then $\mathcal{U}_e$ is free as an $\mathcal{H}_e$ -module with basis $\{w^\rho \ell^m E(\varphi)\}$, $\rho \in \Lambda$, $m \geq 0$. In particular $\mathcal{U}_e \neq 0$ and $\mathcal{H} \subset \mathcal{U}_e$.*

Proof. One cannot argue that $K/\mathbf{Z}$ is torsion-free; it is not, since it contains $\mathbf{Q}/\mathbf{Z}$, and the rational classes are exactly the ones ν_6 counts. Instead use base change. The group algebra $\mathcal{H}_e[K]$ is free over $\mathcal{H}_e[\mathbf{Z}] = \mathcal{H}_e[t, t^{-1}]$ on any set Λ of coset representatives, and $\mathcal{U}_e$ is obtained from it by the base change $\mathcal{H}_e[t, t^{-1}] \rightarrow \mathcal{H}_e$, $t \mapsto w$, a surjection with kernel $(t - w)$ generated by a nonzerodivisor. Base change of a free module is free on the image of the same basis. The crossed multiplication $w^\rho w^{\rho'} = w^n w^{\rho''}$ is well defined, and the derivation is consistent with the integer identification because w^n sits at e_z -exponent n/e and θ_w multiplies it by n . The factors $E(\varphi)$ form the group algebra of a torsion-free group and ℓ is a polynomial variable. $\square$

Lemma 3 (The turn). *There is a unique automorphism σ of $\mathcal{U}_e$ with*

$$\sigma(w) = e^{2\pi i/e} w, \quad \sigma(w^\rho) = \text{Exp}_V(\rho/e) w^\rho, \quad \sigma(\ell) = \ell + \frac{2\pi i}{e}, \quad \sigma(E(\varphi)) = E(\sigma\varphi),$$

acting on $\mathcal{H}_e$ by multiplying the coefficient at $(\gamma, j/e)$ by $e^{2\pi i j/e}$. It fixes $\mathcal{H}$, hence $\mathcal{C}$, pointwise, and commutes with θ_w .

Proof. The action on $\mathcal{H}_e$ multiplies coefficients by a character of the value group, hence is a support-preserving ring automorphism, and is the identity on $\mathcal{H}$ because there $j \in e\mathbf{Z}$. Multiplicativity in ρ is the homomorphism property of Exp_V on $(K, +)$, and the two prescriptions agree on integral exponents because $\text{Exp}_V(n/e) = e^{2\pi i n/e}$ by (4.0a). Commutation with θ_w is not a diagonality argument, since neither $\sigma(\ell) = \ell + 2\pi i/e$ nor $\theta_w(\ell^m) = m\ell^{m-1}$ is diagonal; it is the direct check $\sigma\theta_w(\ell^m) = m(\ell + \frac{2\pi i}{e})^{m-1} = \theta_w\sigma(\ell^m)$, together with the diagonal cases. $\square$

Lemma 4 (Constants). *Under the constant-coefficient hypothesis of Definition 1, $\mathcal{U}_e^{\theta_w=0} = \mathcal{C}$.*

Proof. Let v be the Hahn valuation of $\mathcal{H}_e$, additive because the coefficient field is a field; θ_w multiplies coefficients by integers and never enlarges a support, so $v(\theta_w c) \geq v(c)$.

Write $x = \sum c_{\rho, m, \varphi} w^\rho \ell^m E(\varphi)$ in the basis of Lemma 2 with $\theta_w x = 0$; the equation splits over φ . For $\varphi \neq 0$, let M be the top ℓ -degree in that block; for each ρ , $\theta_w(c_{\rho, M}) + \rho c_{\rho, M} + \theta_w(\varphi) c_{\rho, M} = 0$. Every monomial of $\theta_w(\varphi)$ has Γ -part 0 and strictly negative e_z -part, so $v(\theta_w(\varphi)c) < v(c) \leq v(\theta_w c + \rho c)$ for $c \neq 0$ and the lowest term cannot cancel; hence $c_{\rho, M} = 0$, and descending in M kills the block.

For $\varphi = 0$ the coefficient equations are $\theta_w(c_{\rho, m}) + \rho c_{\rho, m} + (m+1)c_{\rho, m+1} = 0$. Descending in m : the top-degree coefficient satisfies the homogeneous equation $\theta_w(c) + \rho c = 0$, and on each monomial this reads $(j + \rho)c = 0$ with $j \in \mathbf{Z}$, so $c = 0$ unless $\rho \in \mathbf{Z}$, that is unless $\rho = 0$. For $\rho = 0$ the top coefficient is θ_w -constant and the next equation gives $M c_{0, M} = -\theta_w(c_{0, M-1})$, whose right side has zero component in e_z -degree 0; comparing that component gives $c_{0, M} = 0$ for $M \geq 1$. What survives is $\rho = 0$, $m = 0$, $\varphi = 0$ with θ_w -constant coefficient, and the θ_w -constants of $\mathcal{H}_e$ are those supported in e_z -degree zero, namely $\mathcal{C}$. $\square$

Remark 5 (The constant-coefficient hypothesis is necessary, and fails in the application). Allow φ a Novikov coefficient of positive weight, $\varphi = x^\gamma w^{-1}$ with $\gamma \in \Gamma$, $\gamma > 0$ — the shape of every exponential factor the draft actually meets, since the eigenvalues λ of $c_1 \star$ have positive Novikov weight. Then $v(\theta_w \varphi) > 0$ in the coefficient-dominant order and the domination argument reverses. Worse, the equation $\theta_w(c) = \theta_w(\varphi)c$ is solved by

$$c = \sum_{n \geq 0} \frac{(-1)^n}{n!} x^{n\gamma} w^{-n},$$

whose support $\{(n\gamma, -n/e)\}_{n \geq 0}$ is increasing, hence well ordered, so $c \in \mathcal{H}_e$ and $cE(\varphi)$ is a θ_w -constant outside $\mathcal{C}$. Lemma 4 is therefore false in that regime, and with it Proposition 6. The structural reason is worth stating on its own: *in the draft's coefficient-dominant order, $\exp(-\lambda/z)$ is already a unit of the coefficient field*, so adjoining $E(\varphi)$ as a free symbol duplicates an existing element and inflates the constants. In that order the exponential factors are not symbols at all, and the entire irregularity of the formal solution sits in its divergent unit part, which is exactly where the rate comparison of Section 3 lives.

Framed operators and transport

Proposition 6 (The framed operator is the matrix of the turn). *Let D be a differential module over $\mathcal{H}$, free of rank n , with connection matrix Ω , and let $Y \in \mathrm{GL}_n(\mathcal{U}_e)$ satisfy $\theta Y = -\Omega Y$. Then $\mathrm{Sol}(D) = \{v : \theta v + \Omega v = 0\} = Y\mathcal{C}^n$, free of rank n and σ -stable; the matrix M with $\sigma(Y) = YM$ lies in $\mathrm{GL}_n(\mathcal{C})$ and is well defined up to $\mathrm{GL}_n(\mathcal{C})$ -conjugacy. For D over $\Omega_V((z))$ in Levelt–Turrittin form, M is the draft's framed operator: in a solution basis adapted to one deck orbit of length d the matrix of σ is block-cyclic with the draft's intertwiners A_i as blocks, and the cyclic-block determinant gives $\chi(X) = \det(X^d I - U)$ with $U = A_{d-1} \cdots A_0$, which is (4.0b).*

Proof. From $\theta(Y^{-1}) = Y^{-1}\Omega$ one gets $\theta(Y^{-1}v) = Y^{-1}(\theta v + \Omega v)$, so horizontality of v means $Y^{-1}v$ has θ_w -constant entries, and Lemma 4 identifies those with $\mathcal{C}$. Since σ commutes with θ and fixes Ω , it preserves $\mathrm{Sol}(D)$; so $M = Y^{-1}\sigma(Y)$ has entries in $\mathcal{C}$ and is invertible, being a product of invertibles, with inverse $\sigma(Y^{-1})Y$. Two solution matrices differ on the right by an element of $\mathrm{GL}_n(\mathcal{C})$, which conjugates M . $\square$

Theorem 7 (Frame transport). *Let D, D' be as in Proposition 6, with solution matrices Y, Y' , and let $G \in \mathrm{GL}_n(\mathcal{H})$ satisfy $\Omega' = G\Omega G^{-1} - \theta(G)G^{-1}$. Then GY is a solution matrix for D' and $\sigma(GY) = (GY)M$. Consequently*

$$M_{f,V}(D') = C_0^{-1} M_{f,V}(D) C_0 \quad \text{for some } C_0 \in \mathrm{GL}_n(\mathcal{C}),$$

so the two framed operators have the same characteristic polynomial in $\mathcal{C}[X]$ and the same multiplicity at every root of unity.

Proof. $\theta(GY) = \theta(G)Y - G\Omega Y = -\Omega'(GY)$. Since σ fixes $\mathcal{H}$ pointwise, $\sigma(G) = G$, so $\sigma(GY) = G\sigma(Y) = GYM$. Writing $Y' = (GY)C_0$ with $C_0 \in \mathrm{GL}_n(\mathcal{C})$ gives $M' = C_0^{-1}MC_0$. $\square$

Remark 8 (Which object is $M_{f,V}$). The framed operator is an automorphism of the horizontal module; its matrix depends on a choice of *solution* frame and is well defined up to conjugacy over the constants. The conjugate GMG^{-1} appearing in the draft's (4.1c) is not the matrix of the turn in any solution frame — Y' differs from GY by a constant matrix, not by G — but

the matrix of the transported automorphism read in the *module* frame of D' . Both readings are legitimate, and they give the same characteristic polynomial, but the draft should say which it means, since conflating them is exactly the step at issue.

Remark 9 (The hypothesis on G is the whole content). Since σ fixes $\mathcal{H}$ pointwise, $\sigma(G) = G$ is automatic once G has entries in $\mathcal{H}$; what matters is that G lies in $\mathcal{H}$ and not in $\mathcal{U}_e$. No hypothesis-free statement exists: Y itself is a gauge over $\mathcal{U}_e$ carrying D to the trivial module, whose framed operator is the identity. The rank-one case shows what survives: if $g \in \mathcal{H}^\times$ and $\theta(g)/g = \delta$, then each monomial gives $(k - \delta)g_{(\gamma,k)} = 0$, so $\delta \in \mathbf{Z}$; a gauge over $\mathcal{H}$ shifts residues only by integers, which Exp_V kills.

The scissors: these hypotheses have no common instance in the application

Theorem 7 is true, and in the application it is empty. Its two hypotheses pull in opposite directions.

- If $\Gamma = 0$, so $\mathcal{H} = \Omega_V((z))$, then Levelt–Turrittin supplies solution matrices, but the pro-Laurent comparison gauge has unbounded negative z -order and is not in $\text{GL}_n(\mathcal{H})$.
- If $\Gamma \neq 0$ with the draft's coefficient-dominant order, the gauge lies in $\mathcal{H}$, but $\mathcal{H}$ is not a formal Laurent series field over its constants and Levelt–Turrittin is unavailable. Concretely, the e_z -component of the Hahn valuation is not a valuation on $\mathcal{H}$: for $f = x^{(0,0)} + x^{(1,-100)}$ and $g = -x^{(0,0)}$ its values on f and g are 0 while its value on $f + g$ is -100 . So there is no Newton polygon, no slope filtration and no Turrittin algorithm. Nor can one descend to the bounded- z -order elements: $f = 1 - zx^{-\gamma}$ has inverse $-\sum_{k \geq 1} z^{-k} x^{k\gamma}$, so they do not form a field.

An abstract Picard–Vessiot ring for a module over $\mathcal{H}$ does exist, with constants $\mathcal{C}$ after passing to the divisible hull. So the missing ingredient is *not* a solution algebra. It is a canonical Levelt–Turrittin normal form over $\mathcal{H}$, singling out which element of the differential Galois group deserves to be called the turn. That is the precise shape of the remaining problem, and it also shows why a purely valuation-theoretic argument cannot supply it.

3 The receiver

The draft's receiver, in the proof of Proposition 4.7, is

$$\mathcal{R}_j = \Omega_{V,j} \otimes_{H_{0,j}} H_{\text{tot},j}, \quad H_{\text{tot},j} = \mathbf{C}((\Gamma_j^{\text{coeff}} \oplus \mathbf{Z}e_z)). \quad (4.4c)$$

Why the tensor product cannot simply be dissolved

The map $a \otimes h \mapsto a \cdot h$ into the Hahn field with $\Omega_{V,j}$ coefficients is not well defined: the copy of $H_{0,j}$ inside $\Omega_{V,j}$ consists of degree-zero constants, the copy inside $H_{\text{tot},j}$ of monomial series, and the tensor product exists precisely to identify them. Nor can the two copies be assigned different jobs, with module coefficients read as constants and gauge bookkeeping as monomials: the gauge relation $\Omega' = G\Omega G^{-1} - \theta(G)G^{-1}$ ties G and Ω through the same Novikov variables, so one copy is forced. With the constants reading the gauge leaves the field; with the monomial reading the exponential factors become Γ -supported and Remark 5 applies.

Two things should also be said about $\mathcal{R}_j$ itself. It is a tensor product of two field extensions and need not be a domain, and the draft reads characteristic polynomials and algebraic multiplicities in $\mathcal{R}_j[X]$; that deserves an explicit remark, which the draft partly gives by proving each factor injects.

A better receiver

Extend the Γ -valuation of $H_{0,j}$ to its algebraic closure and then to $\Omega_{V,j}$, and take the field of series $\sum_k c_k z^k$ with $c_k \in \Omega_{V,j}$ whose support $\{(v_\Omega(c_k), k)\}$ is well ordered in the chosen order. This has a single copy of the coefficients, is a field rather than a tensor product of two fields, hosts the pro-Laurent gauge, and hosts the exponential factors as honest units. It is strictly better than (4.4c). It does not by itself supply Levelt–Turrittin, and it does not remove the rate criterion below.

The two rates

Weight a monomial of coefficient weight d and z -power k by $a d + b k$, $a, b > 0$, and let ε be the minimal weight of a generator of J_j .

- *Pro-Laurent gauges gain one filtration unit per unit of z -pole.* By the recursion (4.1a) and the draft’s own conclusion, G_α has no power of z below $-|\alpha|$ and a term of bulk degree m lies in $F^m B$; every monomial of J_j^N has weight at least N . Their weights grow like $m(a-b)$, which together with the draft’s finite-below argument gives well ordered support when $a > b$.
- *Splitting gauges lose $w(\Delta\lambda)$ per unit of z -power.* This is the draft’s own construction (4.9d): at order n the block off-diagonal coefficient is removed by a Sylvester equation $[K_0, A_n] = B_n$ with K_0 the leading matrix, so one division by an eigenvalue difference per order, and $w(A_n) \geq -n w(\Delta\lambda)$. The draft’s displayed cubic entry $-14/(81r^2)$ at z -order two exhibits the rate: one factor of r^{-1} per unit of z -power. Three caveats. After a ramification $z = w^e$ the divisions occur once per w -order, so the loss is $e w(\Delta\lambda)$ per unit of z -power; at the cubic the draft proves the exponential blocks are unramified in z , so $e = 1$ there. The claim that the regular-singular normalization costs nothing needs the homogeneity argument that residue eigenvalues are rational, verified in the cubic block where $\det L_s = (s + 1/6)(s + 5/6)$. And $w(\Delta\lambda)$ is not controlled by $w(\lambda)$ when the Picard rank exceeds one, since w and the quantum grading are different functionals and leading terms can cancel.

Both conditions hold together iff

$$e w(\Delta\lambda) < \varepsilon$$

The criterion is not an artifact of choosing an order

For $C := e w(\Delta\lambda)/\varepsilon \geq 1$ the obstruction is not about well ordering at all. The z^n -coefficient of the single product GP of a pro-Laurent gauge with a splitting gauge is $\sum_m g_m p_{n+m}$, whose terms have weight at least $m(1-C) - nC$; for $C \geq 1$ infinitely many terms have bounded weight, so that coefficient is an infinite sum in no completion, ordered or not. Any repair that forms this product must therefore establish $C < 1$. A repair that never forms it — see route 2 below — is not bound by this.

The criterion depends on a normalization the draft may choose

This is the practical point, and it reverses the recommendation of the first version of this memo. The draft fixes $w(u) = 1$ and $w(s_{j,\ell}) = 1$ independently of L , and sets $w(Q^{i_*d}u^{\rho_C \cdot d}) = L(H \cdot i_*d) + \rho_C \cdot d$. Hence $\varepsilon = 1$ always, while the weights of the Novikov generators grow linearly in L . The criterion is therefore

$$e(L(H \cdot i_*d_0) + \rho_C \cdot d_0) < c_1 \cdot d_0,$$

using that the eigenvalues of $c_1\star$ are homogeneous of quantum degree one when Q^d has degree $c_1 \cdot d$ — a convention I have checked against the draft’s own low-dimensional vanishing proof, where the Q^β -component of $c_1\star$ maps H^i to $H^{i+2-2c_1\cdot\beta}$, and against its explicit cubic block, where K_0 has eigenvalues $\pm 6r, 0, 0$ with $r = (3q)^{1/2}$ and $c_1 \cdot \ell = 2$. Consequences:

- Under the draft’s instruction to choose L “so large that” separation holds, the criterion fails at essentially every comparison with two distinct exponential factors — including the cubic endpoint.
- Under the smallest admissible L it can hold. For a projective bundle with $c_1(V) \cdot \ell = 0$ one may take $L = 1$, giving $w(Q^\ell) = 1 = \varepsilon$ and $w(\Delta\lambda) = w(q^{1/2}) = 1/2$.
- Independently of L : separation forces $w \geq \varepsilon$ on every generator of J_j , and the image of Q^{d_0} is a generator, so $c_1 \cdot d_0 \geq 2$ is necessary under any admissible weight. Index one fails outright.

So the cheapest available repair is to choose the separating weight minimally rather than large, and to check the resulting inequality case by case. The first version of this memo asserted the opposite, on the false ground that the ratio was scale-invariant; rescaling w does move both rates equally, but L is not a rescaling, because the bulk generators’ weights are pinned at one.

4 The iterated receiver

The criterion above is a consequence of one architectural choice, and a reader of the second version of this memo proposed the alternative: the draft puts the intrinsic Novikov variables of the endpoint and the new completion variables into a single ordered group, which is what gives an eigenvalue difference positive weight. Nothing forces that. Treat the intrinsic Novikov fraction field as *coefficients*, and complete only in the new exceptional and bulk directions afterwards.

Two facts make the proposal concrete, and both are visible in the draft.

- *The unbounded negative z -order is bulk-driven, not Novikov-driven.* In the recursion (4.1a) the coefficient G_α has no power of z below $-|\alpha|$, where α indexes the *bulk* parameter η , and the filtration depth is acquired by substituting $\eta \in F^1B$. A completion in the bulk directions alone therefore already hosts the pro-Laurent gauge; the Novikov directions are not needed for it.
- *With the Novikov directions as coefficients, the pathology disappears and the criterion is satisfied rather than evaded.* Eigenvalues of $c_1\star$ become constants of weight zero, so $\exp(-\lambda/z)$ is not a unit of the receiver but a genuine exponential symbol, the constant-coefficient hypothesis of Definition 1 holds, and Remark 5 evaporates. Since $w(\Delta\lambda) = 0$, the inequality $ew(\Delta\lambda) < \varepsilon$ holds trivially, and the product GP of Section 3 converges coefficientwise.

The shifts that do involve the exceptional Novikov root do not force those directions back into the filtration. The rank-one unit twist attached to $-(c-1)\lambda_j$, the fixed divisor shift, and the ambient target's $\tau^\circ = O(q^{-1})$ all enter Lemma 4.5 as its a_0 and a_2° terms, which are required only to be *constant in the positive filtration*, not to lie in it. With the Novikov directions as coefficients, $\exp(-a_0/z)$ is an honest exponential symbol rather than the object the draft must argue about separately.

What the sources actually permit

I have checked this against the pinned sources rather than left it as a question, and the answer is positive for the comparison and negative for the gauge.

Iritani's Theorem 5.18 states Ψ as an isomorphism of $\mathbf{C}[z]((q^{-1/s}))[[Q, \tilde{\tau}]]$ -modules, with the change of variables defined over $\mathbf{C}((q^{-1/(r-1)}))[[Q]]$. So the ring is *Laurent* in the exceptional root and *formal* in the ambient Novikov variables and the bulk. Two consequences.

The comparison survives. Inverting the Novikov monomials and completing in the exceptional and bulk directions is a ring homomorphism out of that ring. Ψ , its inverse, and the two inverse identities are module maps over it, so they base-change along it unchanged, and the framed operators' entries lie in the coefficient field, which injects. Nothing in the transport step needs Iritani's proof, only his statement, so the formality of his ring in Q is not by itself an obstacle.

The gauge does not. The source's own reconstruction writes $\tau(\tilde{\tau}) = \tau^\circ + t(\tilde{\tau})$ and $\varsigma_j(\tilde{\tau}) = \varsigma_j^\circ + s_j(\tilde{\tau})$, with $t(\tilde{\tau}) \in H^*(X)((q^{-1}))[[Q, \tilde{\tau}]]$ and s_j of the same shape over $H^*(Z)$. The initial terms are harmless: τ° and ς_j° are by definition the values at $Q = \tilde{\tau} = 0$, hence carry no Novikov dependence at all, and the exceptional direction remains a completion direction as the draft already has it. But t and s_j vanish only at $Q = \tilde{\tau} = 0$, so at $\tilde{\tau} = 0$ they retain pure-Novikov terms. Those have to be removed by the normalized gauge of Lemma 4.5, and that construction converges by topological nilpotence of the bulk shift. Inverting the Novikov directions destroys exactly that.

So the iterated receiver is not free, and the decisive question is now small and graded rather than general:

Is the pure-Novikov part of the coordinate shift — t and s_j restricted to $\tilde{\tau} = 0$ — confined to $H^0 \oplus H^2$?

If it is, the string and divisor equations absorb it into the a_0 and a_2° terms of Lemma 4.5, which are required only to be constant in the positive filtration, and the iterated receiver goes through for the ambient summand with $w(\Delta\lambda) = 0$. If it is not, the Novikov directions must stay inside the filtration, eigenvalue differences reacquire positive weight, and the criterion of Section 3 returns.

The grading answers this, and the answer is negative in general. Iritani's Theorem 5.18(5) states that $\tau(\tilde{\tau})$ and $\varsigma_j(\tilde{\tau})$ are homogeneous of degree 2, and his conventions are $\deg Q^d = 2c_1(X) \cdot d$ and $\deg \tau_i = 2 - \deg \phi_i$. A pure-Novikov term $Q^d \phi$ therefore satisfies

$$\deg \phi = 2 - 2c_1(X) \cdot d,$$

so $c_1 \cdot d = 1$ contributes in H^0 , $c_1 \cdot d = 0$ in H^2 , and $c_1 \cdot d = -1, -2, \dots$ in $H^4, H^6, \dots$; classes with $c_1 \cdot d \geq 2$ cannot contribute at all. Hence the required confinement to $H^0 \oplus H^2$ follows exactly when every contributing effective class satisfies $c_1 \cdot d \geq 0$, and it is not automatic: Iritani's

theorem is stated for an arbitrary smooth projective X , and after several blowups in a weak factorization there is no reason for the intermediate ambient variety to have nef anticanonical class.

So the iterated receiver does not by itself remove arbitrary intermediate fourfolds from the problem. It removes them under a semipositivity hypothesis on the ambient, which the weak factorization does not supply. This is worth stating plainly, because it is the second proposed repair that fails on inspection, and it fails for a reason the draft would have inherited silently.

For the center the same analysis applies, with the additional and unchanged difficulty that its Novikov map is the noninjective one, which is why divisor tagging exists.

The scope consequence is the reason to try this first. If the ambient endpoint is handled this way, no inequality is needed for arbitrary intermediate fourfolds — the case that looked least tractable. What remains is the center summands, and there the draft's own low-dimensional classification is restrictive: for nef canonical class it produces an integral- z gauge to a regular-singular connection, so there is no exponential separation at all and $w(\Delta\lambda) = 0$ again; the remaining minimal cases are $\mathbf{P}^1$, $\mathbf{P}^2$, ruled surfaces, and point blowups. Proving primitive-sixth vanishing directly for those, in the coordinates Iritani's specialization actually produces, is a smaller problem than a general transport theorem.

5 The framing route, checked against the sources and closed

The second version of this memo proposed the following route and called it the probable repair. I state it as it was proposed, then report the three checks, then state what survives. The checks were run against arXiv:2411.02266 as cached, arXiv:2307.03696v4, and arXiv:2307.13555v3; the full verdicts, locators and file hashes are in the companion report *The three framing-compatibility checks, run against Iritani's decomposition*.

The route as proposed

Everything above treats the pro-Laurent bulk gauge of Lemma 4.5 as unavoidable. It may not be. Hinault, Yu, Zhang and Zhang work with F -bundles on the formal u -disc, where a framing is a regular flat connection and, in a framing-flat trivialization, the meromorphic connection takes the normal form

$$\nabla_{u\partial_u} = u\partial_u + u^{-1}K + (\mu D + H).$$

Their Theorem 4.34 characterizes when two connections of that shape are gauge-equivalent under

$$\Phi(u) \in \mathrm{GL}(H[[u]]), \quad \Phi_{ij}(u) \in k[c_1, \dots, c_r][[u]],$$

with Φ then unique given its value at $u = 0$; the conditions are that the leading operators K be conjugate by a parameter-polynomial ϕ , that the two μ agree, and that the diagonal entries of H agree after conjugation modulo the parameter ideal. Their applications are uniqueness of the decomposition for projective bundles and for blowups, complementing the existence theorems of Iritani–Koto and Iritani.

The shape of that gauge is exactly what this memo has been missing. A gauge in $\mathrm{GL}_n(K[[u]])$ has only nonnegative integral powers of the loop coordinate, adjoins no root of u , and is therefore fixed by the turn $u \mapsto e^{2\pi i}u$. Taking $\Gamma = 0$, so that $\mathcal{H} = \Omega_V((u))$ and Levelt–Turrittin is available, Theorem 7 applies with no further work:

$$\Phi \in \mathrm{GL}_n(K[[u]]), \quad \nabla' = \Phi \nabla \Phi^{-1} - \theta(\Phi) \Phi^{-1} \implies M_{f,V}(\nabla') \text{ and } M_{f,V}(\nabla) \text{ conjugate.}$$

No Hahn receiver, no product GP , and no inequality $ew(\Delta\lambda) < \varepsilon$, provided the decomposition of Proposition 4.7 can be identified with an F -bundle isomorphism of the class Theorem 4.34 governs. That identification, the matching of the $u = 0$ datum, and the separation of the center specialization were the three checks.

Check one: the identification does not exist as posed

It fails three times over, and the first two are independent of the draft.

Domain. Section 4.4 of Hinault–Yu–Zhang–Zhang is titled *Equivalence of F -bundles over a point*, and both of its applications evaluate at the closed point $q = t = 0$ of the base, the large-radius limit; the proof of their Theorem 5.16 opens by observing that there the quantum product reduces to the classical cup product, so that the leading operator K_{split} of their (5.18) is a cup-product operator and hence nilpotent. The draft’s ν_6 is not evaluated there. It is read where the eigenvalues of $c_1\star$ separate — at the cubic, where the block carries $r = (3q)^{1/2}$ and the indicial roots $\pm 1/6, \pm 5/6$. At $q = 0$ that separation collapses and the exponential blocks the framed operator is built from do not exist. Their Theorems 5.20 and 5.24 do reconstruct the isomorphism over the whole base from its restriction at the limit point, but that is uniqueness of a map whose existence is Iritani’s and Iritani–Koto’s theorem; it is not a transport of framed monodromy between base points.

Hypotheses, for the blowup. Section 4.4 assumes the generalized eigenspaces of K all have the same dimension, and the matrix calculus of Definition 4.33 and Theorem 4.34 rests on the resulting splitting $H_0^{\oplus m} \cong H$. The blowup decomposition is $H^*(X) \oplus \bigoplus_{i=1}^{m-1} H^*(Z)[-2i]$, whose summands differ in dimension whenever $\dim H^*(Z) \neq \dim H^*(X)$; the authors write a general endomorphism there as a matrix with $\Phi_{1,1} \in \text{End}(H^*(X))$ and $\Phi_{i,i} \in \text{End}(H^*(Z))$, so the mismatch is explicit in their own notation. Their blowup statement, Theorem 5.22, does assert existence of the isomorphism, and Theorem 5.24 gives the corresponding uniqueness; the point is not that it is unproved but that it is not *derived from* Theorem 4.34, whose hypotheses do not cover the unequal block sizes, and that its existence is referred to the earlier decomposition work. So Theorem 5.22 supplies no leverage of its own on the blowup half of Proposition 4.7 — equation (4.3), the half the headline theorem cannot do without — beyond what Iritani’s theorem already gives. The projective-bundle half is genuinely covered: there the eigenspaces are m copies of $H^*(X)$ by their Lemma 5.8, matching Section 4.4, and Theorem 5.16 is proved.

Purpose. Even where Theorem 4.34 applies, its conclusion compares the source at its base point with the target based at a shifted class $\Delta(a)$, pinned by their (5.17) for projective bundles and (5.23) for blowups. That shift is exactly what the pro-Laurent gauge of Lemma 4.5 exists to undo. The route relocates the bulk displacement rather than removing it.

The two normalizations do agree on where the displacement sits, which is a useful check on the draft’s reading of the sources. In their limit-point coordinates the eigenvalue $\lambda_i \in \mathbf{C}[c_1, \dots, c_m]$ is cohomology-valued, so $\Delta(a)$ acquires components in $H^{\geq 4}$ from the higher Chern classes, while the H^2 components cancel identically — their Lemma 5.8(2) gives $m\lambda_i \equiv m\xi^{i-1} - c_1V$, cancelling the c_1V on the right of (5.17) — and the H^2 part of $\Delta(a)$ is left free. In Iritani’s coordinates the same data reads $\varsigma_j^\circ = -(c-1)\lambda_j + h_{Z,j} + O(q^{-1/(c-1)})$ with λ_j a *scalar* multiple of $q^{1/(c-1)}$, so the H^0 part is that scalar, the H^2 part is $h_{Z,j}$, and everything of degree at least four has moved into the tail. Same content, different bookkeeping; the draft’s split of ς_j° into unit twist, fixed divisor, and positive-filtration tail is right. The tail is irreducible: for a trivial rank-two bundle all $c_i(V)$ vanish and their $\Delta(a)$ is purely $H^0 \oplus H^2$, yet Iritani–Koto’s ς_j° still carries its $O(q^{-1/r})$ tail, so even $X \times \mathbf{P}^1$ does not escape the bulk gauge.

Check two: the $u = 0$ datum, which passes

Their Theorems 5.20(2) and 5.24(2) determine the base map from its restriction to the base point up to a multiplicative constant in each logarithmic direction; the proof of their Proposition 4.31 shows why, since df is recovered through $\Psi(d \log p_i) = d \log f_i$ and integrating a logarithmic form leaves one constant. That ambiguity is a character of the Novikov lattice: $q \mapsto cq$ is $Q^d \mapsto c^{E \cdot d} Q^d$, which is the divisor substitution (4.1) of the draft with a_2° a logarithmic multiple of the dual class. The draft already proves such a substitution is a z -constant coefficient automorphism and leaves ν_6 unchanged, and already checks it is well defined on the image monoid. The bundle map carries no ambiguity at all once its restriction is fixed.

The ambiguity also never has to be tolerated, because the sources pin it. Iritani–Koto’s (5.11) gives $\varsigma_j^\circ = r\lambda_j + [z^{-1}] \log(q^{c_1(V)/(rz)} F_j(1))$ in closed form and their (5.12) gives Φ_j° ; Iritani’s Section 5.8.1 fixes $\Psi|_{Q=\tilde{\tau}=0}$ from his Lemma 5.13 and the explicit λ_j . One manuscript sentence citing those, and noting that any residual constant is a divisor substitution already covered by (4.1), settles this point for good, whichever repair eventually lands.

Check three: the center specialization, which stays separate

It stays separate more firmly than the check anticipated. Their A -model F -bundle is built throughout on the one-variable Novikov projection $q^\beta \mapsto q^{\beta \cdot \omega}$ of their (2.21), and in the blowup application the center copies are the maximal F -bundles of Z associated to $\sigma^* \omega_X$ — the Novikov direction pulled back from the ambient variety, which is the draft’s specialization in a coarser form. No statement of theirs can therefore certify that a specialized center module carries an intrinsic framed monodromy. Divisor tagging remains the only route to intrinsic center invariants.

One genuine gain came out of it. Their Lemma 2.24 proves that the collapsed potential determines the full genus-zero potential, by expanding in the H^2 bulk directions and separating curve classes by their intersection numbers with an H^2 basis, under the finiteness of their Assumption 2.22. That is the mechanism of `lem:divisor-tagging`, in print, in a closely related setting. It does not close that lemma — it works at the level of the potential over a base retaining all H^2 directions, whereas the draft needs the separation at the level of a framed operator at a fixed bulk point, and the draft’s collapse is by the pair $(i_*, \rho_C \cdot)$ rather than by a single nef class — but it makes the shape of the argument a cited one.

What survives, and the corrected diagnosis

Two things survive, and both are worth more than the route was.

The comparison isomorphisms need no receiver. Iritani’s Ψ is an isomorphism of $\mathbf{C}[z]((q^{-1/s}))[[Q, \tilde{\tau}]]$ -modules (Theorem 5.18), and Iritani–Koto’s Φ is induced by a $\mathbf{C}[z, q][[Q, \tilde{\tau}]]$ -module map and is an isomorphism over $\mathbf{C}[z]((q^{-1/r}))[[Q, \tilde{\tau}]]$ (proof of their Theorem 5.1(6)). Both are polynomial in z with inverses in the same ring, hence lie in $\mathrm{GL}_n(F((z)))$ for F the Laurent coefficient field, hence are fixed by the turn. Theorem 7 applies to them with $\Gamma = 0$, where Levelt–Turrittin is available: no Hahn field, no product GP , no inequality. The receiver problem was never a property of the comparison.

The pro-Laurent gauge is not the draft’s choice. This corrects the sentence the second version of this memo built its recommendation on. Iritani–Koto transport along the same object: their reconstruction in Section 5.8 uses $M = \bigoplus_j e^{-\varsigma_j^\circ/z} M_B(\varsigma_j^\circ + s_j; Qq^{-c_1(V)/r})$, described in their footnote as a fundamental solution for the summands with respect to $(Q, s_0, \dots, s_{r-1})$ but not

for q and z , satisfying $M = \text{id} + O(z^{-1})$ and lying in $\text{End}[z^{-1}]((q^{-1/r}))[[Q, s]]$. Per bulk degree it is a *polynomial* in z^{-1} — which is the draft's own bound (4.1a), that G_α has no power of z below $-|\alpha|$ — and it is unbounded only after summing over bulk degrees. The sources meet this object and handle it not by an ordered receiver but by Birkhoff factorization, splitting the z -regular factor Φ from the $\text{id} + O(z^{-1})$ factor M' in $(\Phi^\circ)^{-1} \circ M = M' \circ \Phi^{-1}$. Any repair should be measured against that, not against a receiver.

The residual gap

After the two survivals, the blocker is no longer transport across the comparison. It is one statement:

Let T be smooth projective with quantum connection ∇ , and let $\tau^\bullet$ be a bulk parameter in the positive filtration — Iritani's $\tau^\circ = q^{-1}[Z] + O(q^{-2})$ on the ambient summand, the $O(q^{-1/(c-1)})$ tail of ς_j° together with s_j on a center summand. Show that the framed formal monodromy of $\nabla|_{\tau=\tau^\bullet}$ has the same primitive-sixth multiplicity as $\nabla|_{\tau=0}$.

Everything else in Proposition 4.7 is proved or reduced to a regular gauge, and `lem:divisor-tagging` inherits this gap and nothing else.

6 The blockwise route, and what the flatness identities already give

This section carries out the second route far enough to see what it costs. It produces three things: an exact evolution equation for the block residue under bulk motion, a proof that multiplicity-one blocks carry no framed monodromy at all, and a reduction of the residual statement to one Sylvester solvability condition per coalesced block. Conventions are Iritani–Koto's, from their Theorem 5.1: write $U = E \star_\tau$, $C_a = \phi_a \star_\tau$, and

$$\nabla_a = \partial_a + z^{-1}C_a, \quad \nabla_{z\partial_z} = z\partial_z - z^{-1}U + \mu,$$

with μ the constant grading operator. Horizontal sections satisfy $z^2\partial_z Y = (U - z\mu)Y$.

The two identities

Lemma 10 (Flatness). $[U, C_a] = 0$ and $\partial_a U = C_a + [C_a, \mu]$.

Proof. The first is commutativity and associativity of the quantum product. For the second, expand $[\nabla_a, \nabla_{z\partial_z}] = 0$. The z^{-2} term is $z^{-2}[U, C_a]$ and vanishes by the first identity; the z^{-1} term is $z^{-1}(-\partial_a U + C_a + [C_a, \mu])$, using $z\partial_z(z^{-1}C_a s) = -z^{-1}C_a s + z^{-1}C_a z\partial_z s$ and $\partial_a \mu = 0$. There is no z^0 term. $\square$

So bulk motion is not an arbitrary negative-loop gauge. The first identity says C_a commutes with every spectral projector P_i of U , hence preserves each generalized eigenspace H_i with $U|_{H_i} = u_i + N_i$; the second says the motion of U itself is expressed through C_a and μ alone.

The block frame

Let $T_a = \sum_j (\partial_a P_j) P_j$ be Kato's transport operator, so that $\partial_a P_i = [T_a, P_i]$ and $P_i T_a P_i = 0$. It is independent of z and has no pole, so it moves the blocks without touching the loop coordinate. For an operator A write $A_i = P_i A P_i$ and $A_{ij} = P_i A P_j$, and let $D_a A_i = \partial_a A_i - [T_a, A_i]$ be the covariant derivative in this frame. A short computation using $\partial_a P_i = [T_a, P_i]$ gives the general rule

$$D_a A_i = P_i (\partial_a A) P_i + P_i [A, T_a] P_i. \quad (6.1)$$

Theorem 11 (Block evolution). *Assume the eigenvalues u_j of U are pairwise distinct as scalars, so that each difference $u_i - u_j$ is invertible, and assume for the residue formula that the neighbouring blocks are semisimple, $N_j = 0$ for $j \neq i$. Then*

$$D_a U_i = C_{a,i} + [C_{a,i}, \mu_i], \quad D_a \mu_i = [C_{a,i}, S_i], \quad S_i := \sum_{j \neq i} \frac{\mu_{ij} \mu_{ji}}{u_i - u_j}.$$

Proof. For U : by (6.1) and Lemma 10, $D_a U_i = P_i (C_a + [C_a, \mu]) P_i + P_i [U, T_a] P_i$. Since C_a commutes with P_i , the first term is $C_{a,i} + [C_{a,i}, \mu_i]$; since U commutes with P_i , the second is $[U_i, P_i T_a P_i] = 0$.

For μ : $\partial_a \mu = 0$, so (6.1) gives $D_a \mu_i = P_i [\mu, T_a] P_i = \sum_{j \neq i} (\mu_{ij} X_{ji} - Y_{ij} \mu_{ji})$ with $X_{ji} = P_j (\partial_a P_i) P_i$ and $Y_{ij} = P_i (\partial_a P_j) P_j$. Differentiating $U P_i = (u_i + N_i) P_i$ and compressing by $P_j (\cdot) P_i$ gives $(\partial_a U)_{ji} + (u_j - u_i + N_j) X_{ji} = 0$, so under the semisimplicity assumption $X_{ji} = (\partial_a U)_{ji} / (u_i - u_j)$ and likewise $Y_{ij} = -(\partial_a U)_{ij} / (u_i - u_j)$. Substituting $(\partial_a U)_{ji} = C_{a,j} \mu_{ji} - \mu_{ji} C_{a,i}$ and $(\partial_a U)_{ij} = C_{a,i} \mu_{ij} - \mu_{ij} C_{a,j}$, the two terms containing $C_{a,j}$ cancel and what remains is $\sum_{j \neq i} [C_{a,i}, \mu_{ij} \mu_{ji}] / (u_i - u_j)$. $\square$

The cancellation is the point. The neighbouring blocks enter $D_a \mu_i$ only through the symmetric combination S_i , and the motion of the block residue is a commutator with the block's own quantum multiplication operator. Without the semisimplicity of the neighbours the same computation leaves the residual term $-\mu_{ij} (\Delta_{ji} + \Delta_{ij}) C_{a,j} \mu_{ji}$ with $\Delta_{ji} = (u_j - u_i + N_j)^{-1}$, which vanishes exactly when the nilpotent parts do; that correction is where a general statement would have to work harder.

Multiplicity-one blocks carry nothing

This is independent of the deformation and settles, in this normalization, the observation the earlier version of this memo only flagged.

Theorem 12 (Simple blocks are invisible). *Let $(\cdot, \cdot)$ be the Poincaré pairing. Then each μ_i is anti-self-adjoint for the restriction of the pairing to H_i , which is nondegenerate; consequently $\text{tr } \mu_i = 0$ and the spectrum of μ_i is symmetric about 0. If u_i has multiplicity one, then $\mu_i = 0$, the block's formal solution is $e^{-u_i/z}$ times a constant, its framed monodromy is the identity, and it contributes no root of unity to ν_6 .*

Proof. Quantum multiplication is Frobenius, $(a \star b, c) = (a, b \star c)$, so U is self-adjoint; distinct generalized eigenspaces of a self-adjoint operator are orthogonal, so each P_i is self-adjoint and the pairing restricts nondegenerately to H_i . The pairing vanishes unless the degrees are complementary, so μ is anti-self-adjoint; hence so is $\mu_i = P_i \mu P_i$ on H_i . An anti-self-adjoint operator has $\text{tr } \mu_i = \text{tr } \mu_i^\dagger = -\text{tr } \mu_i$ and characteristic polynomial even up to sign, giving the trace and symmetry statements. On a one-dimensional space anti-self-adjointness forces $\mu_i = 0$, and then $z^2 \partial_z Y = u_i Y$ integrates to $e^{-u_i/z}$, single-valued under the turn. $\square$

Corollary 13. ν_6 is carried entirely by the blocks of $E\star$ whose eigenvalue has multiplicity greater than one.

This matches the draft's own cubic computation, where K_0 has eigenvalues $\pm 6r, 0, 0$ and the primitive-sixth exponents come from the repeated 0: the $\pm 6r$ blocks are simple, so by Theorem 12 they contribute the identity, and the draft's indicial roots $1/6$ and $5/6$ are $\pm 1/6$ modulo $\mathbf{Z}$, exactly the symmetry about 0 that the theorem forces on a block residue.

What is left, exactly

Proposition 14 (Isomonodromy criterion for a block). *Write the block connection as $\nabla_i = z\partial_z + A_i(z)$ with $A_i(z) = -z^{-1}U_i + \mu_i$. Under the hypotheses of Theorem 11, the deformation $D_a\nabla_i$ is an infinitesimal gauge transformation by $\Theta_a(z) = -z^{-1}C_{a,i} + \beta$ if and only if*

$$[\beta, U_i] = 0 \quad \text{and} \quad [\beta, \mu_i] = [C_{a,i}, S_i].$$

The z^{-2} and z^{-1} obstructions vanish identically for that choice of leading term; only the z^0 equation remains.

Proof. Infinitesimally, $\delta A_i = [\Theta_a, A_i] - z\partial_z\Theta_a$. With $\Theta_a = z^{-1}\alpha + \beta$ this is $-z^{-2}[\alpha, U_i] + z^{-1}([\alpha, \mu_i] - [\beta, U_i] + \alpha) + [\beta, \mu_i]$. Theorem 11 gives $D_a\nabla_i = -z^{-1}(C_{a,i} + [C_{a,i}, \mu_i]) + [C_{a,i}, S_i]$. Taking $\alpha = -C_{a,i}$ kills the z^{-2} term because $C_{a,i}$ commutes with U_i , and reduces the z^{-1} equation to $[\beta, U_i] = 0$; the z^0 equation is the second displayed condition. $\square$

So the residual statement of Section 5 is now one solvability question, asked only of the coalesced blocks that Corollary 13 leaves in play: is there, in the commutant of U_i , a β with $[\beta, \mu_i] = [C_{a,i}, S_i]$? If yes in every bulk direction, the block connection is isomonodromic along the displacement, its formal invariants are constant, and Proposition 4.7 closes without any receiver.

Two remarks on the shape of the risk, so that the next attempt does not start optimistic. First, $[C_{a,i}, S_i]$ is a commutator of two self-adjoint operators — $C_{a,i}$ by the Frobenius property, S_i because $\mu_{ij}^\dagger = -\mu_{ji}$ makes each $\mu_{ij}\mu_{ji}$ self-adjoint — hence anti-self-adjoint, which is consistent with μ_i remaining anti-self-adjoint but is not by itself the solvability. Second, and this is the real danger, a coalesced block can *split* under the displacement: the splitting is governed by $D_aU_i = C_{a,i} + [C_{a,i}, \mu_i]$, and if a rank-two block carrying the exponents $\pm 1/6$ split into two simple blocks, those would carry residue 0 by Theorem 12 and ν_6 would drop by two. So the theorem that has to be proved is not general isomonodromy but exactly this: the reconstruction displacement does not split the blocks that carry ν_6 , or splits them in a way that preserves the count. That is a coalescence question in the sense of the isomonodromy-at-coalescing-eigenvalues literature, which is the next thing to read.

7 Routes, in the order I would now try them

The framing route has been removed from this list by the checks of Section 5. All below attack the residual gap stated at the end of that section, and the order has changed: the blockwise route is now first, because Section 6 has already carried it past the point where the Birkhoff route still stands.

1. **Blockwise, avoiding transport altogether.** Developed in Section 6. It has already produced the two evolution equations $D_a U_i = C_{a,i} + [C_{a,i}, \mu_i]$ and $D_a \mu_i = [C_{a,i}, S_i]$, the vanishing of every multiplicity-one block residue, and a reduction of the residual statement to one Sylvester condition per coalesced block. What remains is that condition and the coalescence question stated at the end of that section. Signs follow the displayed connection of Iritani–Koto’s Theorem 5.1 and should be rechecked against the draft’s normalization before anything is written into the manuscript.
2. **Birkhoff.** The identity $(\Phi^\circ)^{-1} \circ M = M' \circ \Phi^{-1}$ of Iritani–Koto’s Section 5.8, and its counterpart in Iritani’s Section 5.8.2, already factors the bulk transport into a z -regular factor, which Theorem 7 handles, and a factor $M' = \text{id} + O(z^{-1})$. What must be shown is that the second factor does not change the primitive-sixth multiplicity. This is a statement about a $z = \infty$ -side gauge acting on a $z = 0$ invariant, so it is not automatic, and it is close to the original difficulty in another form: any proof will have to use where M' comes from, which is the flat quantum connection — that is, the identities route 1 already exploits directly. Worth keeping as a cross-check rather than as the primary attack.
3. **The low-dimensional centers directly:** primitive-sixth vanishing after Iritani’s specialization, using the classification the draft already proves, rather than through a general transport statement.
4. **Divisor tagging on its own terms:** whether the divisor equation gives a more explicit transport than the general positive-bulk gauge of Lemma 4.5. Their Lemma 2.24 is the same mechanism in print and is worth citing; it does not close the lemma, for the reasons under check three.
5. **Normalize, then verify.** Replace “choose L so large” by the smallest weight satisfying separation, adopt the single-copy receiver of Section 3, and verify $ew(\Delta\lambda) < \varepsilon$ case by case. This is fallback machinery now.

The iterated receiver of Section 4 is not on the list: it needs a semipositivity hypothesis on the ambient variety that weak factorization does not supply. It remains worth having where it applies. Restricting Proposition 4.7 to centers satisfying the criterion is likewise available, at a cost in generality that is the author’s call.

The observation that multiplicity-one blocks contribute nothing, flagged in the second version of this memo on the strength of Dubrovin’s zero-diagonal fact in canonical coordinates, is no longer a flag: Theorem 12 proves it in this normalization, from self-adjointness of $E\star$ and anti-self-adjointness of μ for the Poincaré pairing. So every route above needs to control only the coalesced blocks.

A separate scope reduction stands on its own, unchanged: a direct quantum Künneth computation for $X \times \mathbf{P}^1$ together with a direct computation for $\mathbf{P}^4$ would leave the blowup step as the only operation formula the headline theorem needs, with the genus-eight corollary waiting. Note that this cannot be had by triviality of the bundle alone — see the end of check one.

8 Placement in the manuscript

Stated by label because these insertions renumber Section 4. Items 1 to 3 and item 5 are unconditional and can be written now; the checks of Section 5 also make item 6 partly unconditional,

since the comparison isomorphisms themselves are z -polynomial and the transport lemma applies to them with no receiver. Only the steps that pass through the pro-Laurent bulk gauge remain conditional on closing the residual gap.

1. The ground field and solution algebra go immediately before “Apply Levelt–Turrittin after a ramification $z = w^e$ ”, which already refers to a universal formal solution algebra that is never constructed.
2. Lemma 4 and Proposition 6 go after `def:framed-sixth-multiplicity`, converting that definition into a characterization, with Remark 5 stating the hypothesis that makes it available.
3. Theorem 7 becomes `lem:frame-transport`, after the paragraph beginning “The frame matters” and before `def:pro-laurent-gauge-group`.
4. In `lem:pv-base-change`, the asserted final clause and the corresponding sentence of its proof become a reference to `lem:frame-transport` — but only once the receiver repair makes that lemma applicable there, since that lemma lives over the pro-Laurent receiver.
5. At (4.1c) in `lem:formal-base-shift`, one sentence identifying M^{bulk} as the transported framed automorphism, in the module frame, with the solution-frame statement cited.
6. `prop:framed-operations` cites the lemma at its three load-bearing steps and says which field plays the role of $\mathcal{H}$ at each. Two of those steps are now unconditional: the transport across Ψ and across Φ needs only that both are z -polynomial with inverses in the same ring, which their statements give, so the field there is $F((z))$ over the Laurent coefficient field and no criterion is verified. The step that removes the positive-filtration bulk parameter is the one that waits.
7. A sentence in `prop:framed-operations` fixing the normalization of the coordinate change, citing Iritani–Koto (5.11)–(5.12) and Iritani’s Section 5.8.1, and noting that any residual constant in a logarithmic Novikov direction is a divisor substitution already covered by (4.1). This is owed regardless of which repair lands, and is check two of Section 5.
8. `lem:divisor-tagging` carries the same obstruction — it embeds the gauge of `lem:formal-base-shift`, whose negative z -order is unbounded, together with every transported operator, into one ordered Hahn field — and needs whatever repair the receiver gets.

Mechanically, a new theorem-like environment must carry exactly one semantic label and a matching row in the claim map consumed by the manuscript’s source-only gate, with an empty declaration list exactly when its coverage is recorded as absent, and the coverage snapshot string in both verification READMEs must be regenerated.

9 Corrections to the earlier versions of this memo

From the second version to this one

Both changes come from running the three checks against the sources.

1. The framing route of Section 5 was called the probable repair. It is not a repair at all: its governing theorem lives at the large-radius limit point, its blowup case is unproved and outside that theorem's hypotheses, and the gauge it supplies lands at the shifted base point the bulk gauge exists to undo.
2. The second version wrote that the unbounded negative loop order is the draft's own choice rather than something the comparison theorems impose. That is wrong, and it was the ground for the recommendation. Iritani–Koto transport along the same object, their fundamental solution M of Section 5.8, with the same per-bulk-degree z^{-1} -polynomiality the draft derives in (4.1a).

Two findings were gained in exchange, and they narrow the problem rather than restate it: the comparison isomorphisms need no receiver, and the residual gap is one displacement statement. Both are in Section 5.

From the first version to the second

For readers of the first version, these are the substantive changes.

1. The Constants Lemma was stated without the constant-coefficient hypothesis and is false without it; see Remark 5. The framed-operator characterization inherits the hypothesis.
2. The scissors were only implicit. The transport theorem has no verified instance in the application, and that now appears where the theorem does.
3. The claim that the criterion is scale-invariant, and the inference that the separating parameter L cannot be tuned, were both wrong. Tuning L down is now the first proposed route.
4. The cubic-endpoint bullet compared two different normalizations. Corrected above.
5. The finite-tensor objection to the receiver was wrong twice over: the pro-Laurent gauge lies in the Hahn factor alone, and the splitting gauge's coefficients lie in one finite extension of $H_{0,j}$. It is replaced by the gauge-relation argument.
6. The remark that a tensor product of two fields is not a domain was asserted as a general principle, which is false. The correct statement is that $\mathcal{R}_j$ need not be one.
7. The obstruction is now given in its order-free form, which answers two of the questions the first version left open: no completion helps, and the criterion is not an artifact of the ordered receiver.
8. Smaller repairs: the freeness proof (the relevant quotient group has torsion), the commutation of σ with the derivation, the inverse of M , the range of the exponents, the ramification factor in the splitting rate, and the divisor-tagging observation, which is new.

10 Questions

The first question of the second version — whether the two connections compared in Proposition 4.7 lie in the class Theorem 4.34 governs — has been answered no, three times over, in Section 5. These replace it.

1. In the commutant of U_i , is there a β with $[\beta, \mu_i] = [C_{a,i}, S_i]$, in every bulk direction and for every coalesced block? By Proposition 14 this is now the question that decides the repair.
2. Can the reconstruction displacement split a coalesced block that carries ν_6 ? The splitting is governed by $D_a U_i = C_{a,i} + [C_{a,i}, \mu_i]$, and by Theorem 12 a split into simple blocks would destroy the primitive-sixth exponents outright.
3. Failing a general answer, does the coalescence literature on isomonodromic deformations with coalescing eigenvalues cover the formal exponents, rather than only the Stokes matrices?
4. Does the factor $M' = \text{id} + O(z^{-1})$ of the sources' Birkhoff factorization preserve the primitive-sixth multiplicity? This is the same question in the other coordinates, and is worth keeping as a cross-check.
5. For the low-dimensional centers, can primitive-sixth vanishing be proved directly in the coordinates Iritani's specialization produces, given that the nef-canonical case already yields an integral- z gauge to a regular-singular connection?
6. Failing all of these, what is the right restriction on centers, and does weak factorization stay inside it?