

Calculus I Lecture Notes

Taylan Şengül

January 7, 2020

Contents

Contents	1
0 Precalculus	3
1 Limits and Continuity	5
1.1 Informal definition of limits	5
1.2 Limits at Infinity and Infinite Limits	8
1.3 Continuity	12
1.4 Formal definition of Limit	17
1.5 Review Problems	18
2 Differentiation	21
2.1 Tangent Lines and Their Slopes	21
2.2 Derivative	23
2.3 Differentiation Rules	24
2.4 Chain Rule	27
2.5 Derivatives of Trigonometric Functions	28
2.6 Higher Order Derivatives	31
2.7 Mean Value Theorem	31
2.8 Implicit Differentiation	34
2.9 Exam 1 Review	35

3	Transcendental Functions	37
3.1	Inverse Functions	37
3.2	Exponential and Logarithmic Functions	40
3.3	The Natural Logarithm and Exponential	42
3.4	The Inverse Trigonometric Functions	47
4	Applications of Derivatives	51
4.1	Related Rates	51
4.2	Indeterminate Forms	52
4.3	Extreme Values	54
4.4	Concavity and Inflections	56
4.5	Graphs of Functions	58
4.6	Extreme Value Problems	59
4.7	Linear Approximation	60
4.8	Exam 2 Review	61
5	Integration	63
5.1	The Definite Integral	63
5.2	The Fundamental Theorem of Calculus	66
5.3	The Method of Substitution	68
5.4	Areas of Plane Regions	70
5.5	Integration by Parts	73
5.6	Integrals of Rational Function	75
5.7	Inverse Substitutions	77
5.8	Indefinite Integrals Exercises	79
5.9	Improper Integrals	86
5.10	Trigonometric Integrals	88
6	Applications of Integration	93
6.1	Volumes Using Cross-Sections	93
6.2	Volumes Using Cylindrical Shells	95
6.3	Arc Length	96
6.4	Areas Of Surfaces Of Revolution	97

Chapter 0

Precalculus

Chapter 1

Limits and Continuity

1.1 Informal definition of limits

Two main problems of calculus are

1. Derivative. Find the rate of change of f .
2. Integral. Find the area under a given curve.

Both are based on the concept of limit.

We say $\lim_{x \rightarrow a} f(x) = L$ to mean that $f(x)$ is “close enough” to L for any x “close enough” to a .

Example 1.1.1. Which value is x close to when x is close to 2?

$$\lim_{x \rightarrow 2} x = 2$$

Example 1.1.2. Which value is 3 close to when x is close to 2?

$$\lim_{x \rightarrow 2} 3 = 3$$

We can generalize these examples.

Theorem 1.1.1. Let a and c be two real numbers. Then

$$\lim_{x \rightarrow a} c = c, \quad \lim_{x \rightarrow a} x = a.$$

The limit $\lim_{x \rightarrow a} f(x)$ may be different from $f(a)$ as the next example shows.

Example 1.1.3.

$$f(x) = \begin{cases} x, & \text{if } x \neq 2 \\ 1, & \text{if } x = 2 \end{cases}$$

Which value is $f(x)$ close to when x is close to (but not equal to) 2?

$\lim_{x \rightarrow 2} f(x) = \lim_{x \rightarrow 2} x = 2$ although $f(2) = 1$.

Informal definition of left and right limits

If $f(x)$ is close to L when $x < a$ and x is close enough to a then we say

$$\lim_{x \rightarrow a^-} f(x) = L$$


This is called the *left limit* of f at $x = a$.

Similarly we can define the right limit.

Theorem 1.1.2. $\lim_{x \rightarrow a} f(x) = L$ if and only if both $\lim_{x \rightarrow a^-} f(x) = L$ and $\lim_{x \rightarrow a^+} f(x) = L$.

Example 1.1.4. Find the left and right limits of the signum function

$$f(x) = \begin{cases} -1 & \text{for } x < 0 \\ 0 & \text{for } x = 0 \\ 1 & \text{for } x > 0 \end{cases}$$



Solution. The one-sided limits exist, but are not equal

$$\lim_{x \rightarrow 0^+} f(x) = 1 \text{ and } \lim_{x \rightarrow 0^-} f(x) = -1.$$

Hence $\lim_{x \rightarrow 0} f(x)$ does not exist.

Properties of Limits

Theorem 1.1.3. Suppose

$$\lim_{x \rightarrow a} f(x) = L, \quad \lim_{x \rightarrow a} g(x) = M.$$

Then

$$\lim_{x \rightarrow a} (f(x) + g(x)) = L + M, \tag{1.1}$$

$$\lim_{x \rightarrow a} (f(x) - g(x)) = L - M, \tag{1.2}$$

$$\lim_{x \rightarrow a} (f(x) \cdot g(x)) = L \cdot M \tag{1.3}$$

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{L}{M}, \quad \text{if } M \neq 0 \tag{1.4}$$

$$\lim_{x \rightarrow a} [f(x)]^n = L^n, \quad n = \text{positive integer} \tag{1.5}$$

$$\lim_{x \rightarrow a} [f(x)]^{1/n} = L^{1/n}, \quad n = \text{positive integer and } L > 0 \text{ if } n = \text{even} \tag{1.6}$$

$$\tag{1.7}$$

Proof. Proof requires the formal definition of limit. □

Using the above properties we can evaluate the following limits.

Example 1.1.5. Find $\lim_{x \rightarrow 2} x^2 + 1$ and $\lim_{x \rightarrow 2} \frac{x^2 + 1}{6 - x}$.

Solution. Using the product rule of limits and the Theorem 1.1.1,

$$\lim_{x \rightarrow 2} x^2 = \lim_{x \rightarrow 2} x \cdot \lim_{x \rightarrow 2} x = 2 \cdot 2 = 4$$

Using the sum rule of limits,

$$\lim_{x \rightarrow 2} x^2 + 1 = \lim_{x \rightarrow 2} x^2 + \lim_{x \rightarrow 2} 1 = 4 + 1 = 5$$

Using the division rule of limits,

$$\lim_{x \rightarrow 2} \frac{x^2 + 1}{6 - x} = \frac{\lim_{x \rightarrow 2} x^2 + 1}{\lim_{x \rightarrow 2} 6 - x} = \frac{5}{4}.$$

The above example is a special case of the following theorem.

Theorem 1.1.4. If $P(x)$ is a polynomial then,

$$\lim_{x \rightarrow a} P(x) = P(a)$$

If $Q(x)$ is another polynomial with $Q(a) \neq 0$ then

$$\lim_{x \rightarrow a} \frac{P(x)}{Q(x)} = \frac{P(a)}{Q(a)}.$$

The Squeeze Theorem

Theorem 1.1.5. Suppose that $f(x) \leq g(x) \leq h(x)$ and $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L$. Then $\lim_{x \rightarrow a} g(x) = L$.




Figure 1.1: The Squeeze Theorem.

Example 1.1.6. If $-x^2 \leq g(x) \leq x^2$ for $-1 \leq x \leq 1$, find $\lim_{x \rightarrow 0} g(x)$.

Example 1.1.7. Show that if $\lim_{x \rightarrow a} |f(x)| = 0$ then $\lim_{x \rightarrow a} f(x) = 0$.

Solution. Note that $-|f(x)| \leq f(x) \leq |f(x)|$ and use the Squeeze Theorem.

More examples

Example 1.1.8. Let

$$f(x) = \frac{|x - 2|}{x^2 + x - 6}.$$

Find $\lim_{x \rightarrow 2+} f(x)$, $\lim_{x \rightarrow 2-} f(x)$. Does $\lim_{x \rightarrow 2} f(x)$ exist?

Most of the limits you need to compute in this class will be $\lim_{x \rightarrow a} f(x)$ when $f(a)$ does not exist. Here is an example.

Example 1.1.9.

$$\lim_{x \rightarrow -2} \frac{x^2 + x - 2}{x^2 + 5x + 6},$$

Solution. Remember that we consider x values close to but not equal to -2 . Hence $x + 2 \neq 0$ and we can make the simplification

$$\lim_{x \rightarrow -2} \frac{x^2 + x - 2}{x^2 + 5x + 6} = \lim_{x \rightarrow -2} \frac{(x+2)(x-1)}{(x+2)(x+3)} = \lim_{x \rightarrow -2} \frac{x-1}{x+3} = \frac{-3}{1} = -3.$$

Exercises.

1. $\lim_{x \rightarrow 5} \frac{\frac{1}{x} - \frac{1}{5}}{x - 5}.$
Answer: $-\frac{1}{25}.$

2. $\lim_{x \rightarrow 4} \frac{\sqrt{x} - 2}{x^2 - 16},$
Hint: multiply and divide by the conjugate expression $\sqrt{x} + 2$. Answer: $\frac{1}{32}$

3. $\lim_{x \rightarrow -2} \frac{x^2 + 2x}{x^2 - 4}.$
Answer: $\frac{1}{2}$

4. $\lim_{h \rightarrow 0} \frac{\sqrt{4+h} - 2}{h}.$
Answer: $\frac{1}{4}$

5. $\lim_{t \rightarrow 0} \frac{t}{\sqrt{4+t} - \sqrt{4-t}}.$
Answer: 2

6. $\lim_{x \rightarrow -1} \frac{x^3 + 1}{x + 1}.$
Answer: 3

7. $\lim_{x \rightarrow 0} \frac{|3x - 1| - |3x + 1|}{x}.$
Answer: -6

8. $\lim_{x \rightarrow -2^-} \frac{x^2 - 4}{|x + 2|}.$
Answer:

9. $\lim_{y \rightarrow 1} \frac{y - 4\sqrt{y} + 3}{y^2 - 1}.$
Answer: $-\frac{1}{2}$

10. $\lim_{x \rightarrow 2} \frac{\sqrt{4 - 4x + x^2}}{x - 2}.$
Answer: 1

11. If $2 - x^2 \leq f(x) \leq 2 \cos x$ for all x , find $\lim_{x \rightarrow 0} f(x).$
Answer: 2

1.2 Limits at Infinity and Infinite Limits

Limits at Infinity

Definition 1.2.1. We will say that $\lim_{x \rightarrow \infty} f(x) = L$ if $f(x)$ is “close enough” to L whenever $x > 0$ is “large enough”.


Similarly we define $\lim_{x \rightarrow -\infty} f(x) = L$ if $f(x)$ is “close enough” to L whenever $x < 0$ is “large enough”.

If either $\lim_{x \rightarrow \infty} f(x) = L$ or $\lim_{x \rightarrow -\infty} f(x) = L$, we say that the line $y = L$ is an **horizontal asymptote** of the graph of f .

Example 1.2.1. Argue that

$$\lim_{x \rightarrow \infty} 1/x = \lim_{x \rightarrow -\infty} 1/x = 0.$$

by making a table of values of x and $1/x$.



Recall that for ordinary limits, limit of product of functions is a product of limits of functions. Same is also true for limits at infinity. Hence

$$\lim_{x \rightarrow \infty} \frac{1}{x^2} = \lim_{x \rightarrow \infty} \frac{1}{x} \cdot \lim_{x \rightarrow \infty} \frac{1}{x} = 0 \times 0 = 0.$$

Similarly

$$\lim_{x \rightarrow -\infty} \frac{1}{x^2} = 0$$

Finally, for any positive integer n

$$\lim_{x \rightarrow \infty} \frac{1}{x^n} = \lim_{x \rightarrow -\infty} \frac{1}{x^n} = 0.$$


Example 1.2.2. Let $f(x) = \frac{x}{\sqrt{x^2 + 1}}$. Find $\lim_{x \rightarrow \infty} f(x)$, $\lim_{x \rightarrow -\infty} f(x)$.

Solution.

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2 + 1}} &= \lim_{x \rightarrow \infty} \frac{x}{|x| \sqrt{1 + 1/x^2}} \\ &= \lim_{x \rightarrow \infty} \frac{x}{x \sqrt{1 + 1/x^2}} \\ &= \lim_{x \rightarrow \infty} \frac{1}{\sqrt{1 + 1/x^2}} \\ &= \frac{\lim_{x \rightarrow \infty} 1}{\lim_{x \rightarrow \infty} \sqrt{1 + 1/x^2}} \\ &= \frac{1}{\sqrt{\lim_{x \rightarrow \infty} (1 + 1/x^2)}} = \frac{1}{1} = 1. \end{aligned}$$

Similarly,

$$\lim_{x \rightarrow -\infty} \frac{x}{\sqrt{x^2 + 1}} = -1$$



Limits of Rational Functions at Infinity

Recall that a rational function is a ratio of two polynomials.

Strategy. To find limits of rational functions at infinity, divide by the highest power of x appearing in the denominator.

Example 1.2.3.

$$\lim_{x \rightarrow \pm\infty} \frac{2x^2 - x + 3}{3x^2 + 5} = \lim_{x \rightarrow \pm\infty} \frac{2 - \frac{1}{x} + \frac{3}{x^2}}{3 + \frac{5}{x}} = \frac{2}{3}.$$

Example 1.2.4.

$$\lim_{x \rightarrow \pm\infty} \frac{x - 5}{2x^2 + 4x + 1} = \lim_{x \rightarrow \pm\infty} \frac{\frac{1}{x} - \frac{5}{x^2}}{2 + \frac{4}{x} + \frac{1}{x^2}} = \frac{0}{2} = 0.$$

We can generalize the above examples.

Theorem 1.2.1. Let $P(x) = a_p x^p + a_{p-1} x^{p-1} + \cdots + a_0$ be a polynomial of degree p and $Q(x) = b_q x^q + \cdots + b_0$ be a polynomial of degree q . If $p = q$, then

$$\lim_{x \rightarrow \pm\infty} \frac{P(x)}{Q(x)} = \frac{a_p}{q_p},$$

If $p < q$, then

$$\lim_{x \rightarrow \pm\infty} \frac{P(x)}{Q(x)} = 0,$$


Example 1.2.5.

$$\lim_{x \rightarrow \infty} \sqrt{x^2 + x} - x = \lim_{x \rightarrow \infty} \frac{(\sqrt{x^2 + x} - x)(\sqrt{x^2 + x} + x)}{\sqrt{x^2 + x} + x} = \lim_{x \rightarrow \infty} \frac{x}{|x| \sqrt{1 + \frac{1}{x}} + x} = \lim_{x \rightarrow \infty} \frac{1}{\sqrt{1 + \frac{1}{x}} + 1} = \frac{1}{2}.$$

Infinite Limits

Example 1.2.6. The values of $\frac{1}{x^2}$ gets larger and larger as x approaches to 0. Thus $\lim_{x \rightarrow 0} \frac{1}{x^2}$ does not exist. Although the limit does not exist, it is useful to state why it does not exist by writing

$$\lim_{x \rightarrow 0} \frac{1}{x^2} = \infty.$$



Example 1.2.7.

$$\lim_{x \rightarrow 0^+} \frac{1}{x} = \infty.$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x} = -\infty.$$

$$\lim_{x \rightarrow 0} \frac{1}{x} \text{ does not exist.}$$

Example 1.2.8.

$$\lim_{x \rightarrow -\infty} \sqrt{x^2 + x} - x$$

Solution. Both $-x$ and $\sqrt{x^2 + x}$ grow large as $x \rightarrow -\infty$. So the limit is ∞ .**Behaviour of Polynomials at Infinity****Example 1.2.9.**

$$\lim_{x \rightarrow \infty} 4x^3 - 2x + 1 = \lim_{x \rightarrow \infty} 4x^3 = \infty.$$

$$\lim_{x \rightarrow -\infty} -3x^5 + x^3 + 1 = \lim_{x \rightarrow -\infty} -3x^5 = \infty.$$

In general,

Theorem 1.2.2. If $P(x) = a_n x^n + \cdots + a_0$ is a polynomial then

$$\lim_{x \rightarrow \pm\infty} P(x) = \lim_{x \rightarrow \pm\infty} a_n x^n.$$

If $Q(x) = b_m x^m + \cdots + b_0$ is also a polynomial then

$$\lim_{x \rightarrow \pm\infty} \frac{P(x)}{Q(x)} = \lim_{x \rightarrow \pm\infty} \frac{a_n x^n}{b_m x^m}$$

Example 1.2.10.

$$\lim_{x \rightarrow \infty} \frac{x^3 + 1}{x^2 - 2x} = \lim_{x \rightarrow \infty} \frac{x + \frac{1}{x^2}}{1 - \frac{2}{x}} = \lim_{x \rightarrow \infty} \frac{x}{1} = \infty$$

Example 1.2.11. 1. $\lim_{x \rightarrow 2} \frac{(x-2)^2}{x^2-4} = 0$

$$2. \lim_{x \rightarrow 2^+} \frac{x-3}{x^2-4} = -\infty$$

$$3. \lim_{x \rightarrow 2^-} \frac{x-3}{x^2-4} = \infty$$

$$4. \lim_{x \rightarrow 2} \frac{x-3}{x^2-4} \text{ does not exist.}$$

$$5. \lim_{x \rightarrow \infty} \frac{2x-1}{\sqrt{3x^2+x+1}},$$

6. $\lim_{x \rightarrow 1+} \frac{\sqrt{x^2 - x}}{x - x^2}$

Solution. If $x > 1$ then $x - x^2 = x(1 - x) < 0$. So

$$\lim_{x \rightarrow 1+} \frac{\sqrt{x^2 - x}}{x - x^2} = \lim_{x \rightarrow 1+} \frac{-\sqrt{x^2 - x}}{x^2 - x} = \lim_{x \rightarrow 1+} \frac{-\sqrt{x^2 - x}}{\sqrt{x^2 - x}\sqrt{x^2 - x}} = \lim_{x \rightarrow 1+} \frac{-1}{\sqrt{x^2 - x}} = -\infty$$

1.3 Continuity

Let $f(x) = \sqrt{4 - x^2}$. Domain of f is $[-2, 2]$.

- $x = -2$ is the left end point of $\text{Dom}(f)$.
- $x = 2$ is the right end point of $\text{Dom}(f)$.
- Any x with $-2 < x < 2$ is called an interior point of $\text{Dom}(f)$.

Definition 1.3.1. A function f is **continuous** at an interior point c of its domain if

$$\lim_{x \rightarrow c} f(x) = f(c)$$

f is continuous at its left endpoint c if

$$\lim_{x \rightarrow c+} f(x) = f(c)$$

f is continuous at its right endpoint c if

$$\lim_{x \rightarrow c-} f(x) = f(c)$$

The following theorem gives an alternative definition of continuity which is sometimes useful.

Theorem 1.3.1. A function f is **continuous** at an interior point c of its domain if and only if

$$\lim_{h \rightarrow 0} f(c + h) = f(c)$$

f is continuous at its left endpoint c if

$$\lim_{h \rightarrow 0+} f(c + h) = f(c)$$

f is continuous at its right endpoint c if

$$\lim_{h \rightarrow 0-} f(c + h) = f(c)$$

Proof. Let $h = x - c$. Then $x \rightarrow c$ if and only if $h \rightarrow 0$. So $\lim_{h \rightarrow 0} f(c + h) = f(c)$ is the same as $\lim_{h \rightarrow 0} f(c + h) = f(c)$. \square

Note that f is discontinuous at c if

- either $\lim_{x \rightarrow c} f(x)$ does not exist.
- or $\lim_{x \rightarrow c} f(x)$ exists but is not equal to $f(c)$.





Figure 1.2: f is discontinuous at a because of (ii) and discontinuous at b because of (i). f is continuous at c .

Definition 1.3.2. f is called a continuous function if f is continuous at every pt of its domain.

Example 1.3.1. $f(x) = \sqrt{4 - x^2}$ is continuous at every point of its domain. So it is a continuous function.



According to this definition $f(x) = \frac{1}{x}$ is also continuous!!! 0 is not in domain of f . So we say f is undefined rather than discontinuous at 0.

There are lots of continuous functions:

- polynomials,
- rational functions,
- rational powers $x^{m/n}$
- trigonometric functions
- absolute value function $|x|$

Theorem 1.3.2. If f and g are continuous at c then

- $f + g$, $f - g$, fg , are continuous at c ,
- if k is constant then kf is continuous at c ,
- $\frac{f}{g}$ is continuous at c provided that $g(c) \neq 0$.

- $f^{1/n}$ continuous at c provided that $f(c) > 0$ if n is even.

Proof. Let's prove that if f and g are continuous at c then so is $f + g$. If f and g are continuous at c then

$$\lim_{x \rightarrow c} f(x) = f(c), \quad \lim_{x \rightarrow c} g(x) = g(c),$$

By the limit rule,

$$\lim_{x \rightarrow c} (f(x) + g(x)) = \lim_{x \rightarrow c} f(x) + \lim_{x \rightarrow c} g(x) = f(c) + g(c).$$

The other proofs are similar. □

Composites of continuous functions are continuous

If g is continuous at c and f is continuous at $g(c)$ then $f \circ g$ is continuous at c . In other words,

$$\lim_{x \rightarrow c} f(g(x)) = f(\lim_{x \rightarrow c} g(x)) = f(g(c)).$$

Example 1.3.2. Find m so that

$$g(x) = \begin{cases} x - m, & \text{if } x < 3, \\ 1 - mx, & \text{if } x \geq 3 \end{cases}$$

is continuous for all x .

Continuity of Trigonometric Functions

Theorem 1.3.3. $\sin x$ and $\cos x$ are continuous at $x = 0$, i.e.


$$\lim_{x \rightarrow 0} \sin x = \sin 0 = 0, \quad \lim_{x \rightarrow 0} \cos x = \cos 0 = 1.$$

Proof.

$$|1 - \cos \theta| = |AQ| \leq |AP| \leq \theta,$$

$$|\sin \theta| = |PQ| \leq |AP| \leq \theta$$

In other words, $-\theta \leq \sin \theta \leq \theta$ and using the squeeze theorem we get $\lim_{\theta \rightarrow 0} \sin \theta = 0$. Similarly, we get $\lim_{\theta \rightarrow 0} 1 - \cos \theta = 0$ or $\lim_{\theta \rightarrow 0} \cos \theta = 1$.



Theorem 1.3.4. $\sin x$ and $\cos x$ are continuous for all x . □

Proof. By Theorem 1.3.1, we need to prove $\lim_{h \rightarrow 0} \sin(x + h) = \sin x$ for any x .

$$\lim_{h \rightarrow 0} \sin(x + h) = \lim_{h \rightarrow 0} \sin x \cos h + \cos x \sin h = \sin x \lim_{h \rightarrow 0} \cos h + \cos x \lim_{h \rightarrow 0} \sin h = \sin x.$$

Prove the continuity of $\cos x$ as an exercise. □

Extreme Value Theorem

Theorem 1.3.5. If f is continuous on the closed interval $[a, b]$ then there exist numbers p and q in the interval $[a, b]$ s.t.


$$f(p) \leq f(x) \leq f(q)$$

for all x in $[a, b]$. $f(p)$ is called the **absolute minimum value** and $f(q)$ is called the **absolute maximum value**.


Extreme value theorem is an existence theorem. It only guarantees the existence of p and q but does not tell how to actually find them.

We say a function f is **bounded** if there exists M and N such that $M \leq f(x) \leq N$ for all x in the domain of f . Extreme value theorem says that continuous functions on closed intervals must be bounded.


Example 1.3.3. The conclusions of the theorem may fail if the function f is not continuous or the interval is not closed.




(a) The function $f(x) = 1/x$ on the open interval $(0, 1)$ is continuous but unbounded and has no minimum and no maximum.



(b) The function $f(x) = x$ on $(0, 1)$ is discontinuous, bounded and has no minimum and no maximum.



(a) This function is defined on the closed interval $[0, 1]$, discontinuous, has a minimum but no maximum.



(b) This function is defined on the closed interval $[0, 1]$, discontinuous, bounded, has no minimum and no maximum.

Intermediate Value Theorem

Theorem 1.3.6 (Intermediate Value Theorem). If f is continuous on $[a, b]$ and if s is between $f(a)$ and $f(b)$ then there exists c in $[a, b]$ s.t. $f(c) = s$.




Figure 1.5: Illustration of the intermediate value theorem.

Example 1.3.4. *If a child grows from 1 m to 1.5 m between the ages of two and six years, then, at some time between two and six years of age, the child's height must have been 1.23 m.*

In particular, a continuous function on a closed interval takes every value between its minimum m and maximum M . Hence its range is a closed interval $[m, M]$.

Example 1.3.5. *Show that the equation $x^3 - x - 1 = 0$ has a solution in the interval $[1, 2]$.*

Solution. $f(x) = x^3 - x - 1$ is a polynomial and hence continuous. $f(1) = -1$ and $f(2) = 5$. Since 0 lies between -1 and 5 , the intermediate value theorem assures us that there must be a number c in $[1, 2]$ such that $f(c) = 0$.

Bisection Algorithm

Intermediate Value Theorem is also an existence theorem. It does not say how to find c in its statement. Let's try to better estimate the root of previous example. Write $f(x) = x^3 - x - 1$ and try to find a smaller interval where a root lies of

$$f(x) = 0.$$

We know that a root lies in $[1, 2]$, if say that the root is 1.5 the maximum error will be 0.5.

Now $f(1.5) = 0.875 > 0$. So a root lies in $[1, 1.5]$, and if we say the root is 1.25 then the maximum error will be 0.25.

If this is not sufficient then compute $f(1.25) = -0.2969$, now if we say the root is 1.375 then the error is less than 0.125.

Next step is $f(1.375) = 0.2246$. So a root must lie in $[1.25, 1.375]$. The error is less than 0.0625 if we say the root is 1.315.

Going this way, we find the approximations, 1.3438, 1.3282, 1.3204. Hence the root must lie in $[1.3204, 1.3282]$. So the first two decimal digits of the root are 1.32.

In engineering, you almost never get exact results. All you can do is lower your error below an acceptable threshold.

Optional Issues

Is there a function which is continuous only at a single point? Yes!

Example 1.3.6.

$$f(x) = \begin{cases} x, & \text{if } x \text{ is a rational number} \\ 0, & \text{otherwise} \end{cases}$$

is continuous only at $x = 0$.

This also answers the following question

If a function is continuous at point, is it continuous in some open interval around that point? NO!

1.4 Formal definition of Limit

The informal description of the limit uses phrases like “close enough” and “really very small”. “Fortunately” there is a good definition, i.e. one which is unambiguous and can be used to settle any dispute about the question of whether $\lim_{x \rightarrow a} f(x)$ equals some number L or not.

In this section we assume that f is defined in an open interval containing a except possibly at $x = a$.

Definition 1.4.1. We say that

$$\lim_{x \rightarrow a} f(x) = L$$

if for every $\epsilon > 0$ there exists a $\delta > 0$ such that

$$0 < |x - a| < \delta \text{ implies } |f(x) - L| < \epsilon. \quad (1.8)$$

Why the absolute values? Recall that the quantity $|x - y|$ is the distance between the points x and y on the number line.

What are ϵ and δ ? The quantity ϵ is how close you would like $f(x)$ to be to its limit L ; the quantity δ is how close you have to choose x to a to achieve this. To prove that $\lim_{x \rightarrow a} f(x) = L$ you must assume that someone has given you an unknown $\epsilon > 0$, and then find a positive δ for which (1.8) holds. The δ you find will depend on ϵ .

When we first discussed the limit, say $\lim_{x \rightarrow 5} 2x + 1$, we made a table,

x	$f(x) = 2x + 1$
5.1	11.2
5.01	11.02
5.001	11.002
4.9	10.8
4.99	10.98
4.999	10.998

This table can be written also in this form.

$ x - 5 $	$ f(x) - 11 $
0.1	0.2
0.01	0.02
0.001	0.002

It looks like for any $\epsilon > 0$, if $|x - 5| < \frac{\epsilon}{2}$ then $|f(x) - 11| < \epsilon$.

1.5 Review Problems

Example 1.5.1. Evaluate the limits if they exist. If they do not exist, state whether they are ∞ , $-\infty$ or just does not exist.

1. $\lim_{x \rightarrow 2} \frac{x^2 + 1}{1 - x^2},$

2. $\lim_{x \rightarrow 1} \frac{x^2}{1 - x^2},$

3. $\lim_{x \rightarrow \infty} \frac{\cos x}{x},$ (Hint: Use Sandwich Theorem)

4. $\lim_{x \rightarrow -\infty} \frac{2x^3 + 2x - 1}{-3x^3 + x^2},$

5. $\lim_{x \rightarrow -\infty} x + \sqrt{x^2 - 4x + 1},$

Solution.

$$\begin{aligned} \lim_{x \rightarrow -\infty} x + \sqrt{x^2 - 4x + 1} &= \lim_{x \rightarrow -\infty} x + |x| \sqrt{1 - \frac{4}{x} + \frac{1}{x^2}} = \lim_{x \rightarrow -\infty} x \left(1 - \sqrt{1 - \frac{4}{x} + \frac{1}{x^2}} \right) \\ &= \lim_{x \rightarrow -\infty} x \left(1 - \sqrt{1 - \frac{4}{x} + \frac{1}{x^2}} \right) \frac{\left(1 + \sqrt{1 - \frac{4}{x} + \frac{1}{x^2}} \right)}{\left(1 + \sqrt{1 - \frac{4}{x} + \frac{1}{x^2}} \right)} \\ &= \lim_{x \rightarrow -\infty} x \left(1 - \left(1 - \frac{4}{x} + \frac{1}{x^2} \right) \right) \lim_{x \rightarrow -\infty} \frac{1}{\left(1 + \sqrt{1 - \frac{4}{x} + \frac{1}{x^2}} \right)} \\ &= \lim_{x \rightarrow -\infty} x \left(\frac{4}{x} - \frac{1}{x^2} \right) \frac{1}{2} = \lim_{x \rightarrow -\infty} \left(4 - \frac{1}{x} \right) \frac{1}{2} = 2. \end{aligned}$$

6. $\lim_{x \rightarrow 0} \frac{x}{|x - 1| - |x + 1|}.$

7. $\lim_{x \rightarrow 5} \frac{x - 5}{x^2 - 25}$

8. $\lim_{x \rightarrow -5} \frac{x^2 + 3x - 10}{x + 5}$

9. $\lim_{x \rightarrow 1} \frac{x^{-1} - 1}{x - 1}$

10. $\lim_{u \rightarrow 1} \frac{u^4 - 1}{u^3 - 1}$

11. $\lim_{x \rightarrow 9} \frac{\sqrt{x} - 3}{x - 9}$

12. $\lim_{x \rightarrow 0} \frac{\frac{1}{x-1} + \frac{1}{x+1}}{x}$

13. $\lim_{v \rightarrow 2} \frac{v^3 - 8}{v^4 - 16}$

14. If $2 - x^2 \leq g(x) \leq 2 \cos x$, for all x , find $\lim_{x \rightarrow 0} g(x)$

15. If $\lim_{x \rightarrow 2} \frac{f(x) - 5}{x - 2} = 3$, find $\lim_{x \rightarrow 2} f(x)$

16. If $\lim_{x \rightarrow 2} \frac{f(x) - 5}{x - 2} = 4$, find $\lim_{x \rightarrow 2} f(x)$

17. $\lim_{h \rightarrow 0^+} \frac{\sqrt{h^2 + 4h + 5} - \sqrt{5}}{h}$

18. $\lim_{x \rightarrow -2^-} (x + 3) \frac{|x + 2|}{x + 2}$

19. Define $g(3)$ in a way that extends $g(x) = (x^2 - 9)/(x - 3)$ to be a continuous at $x = 3$.

20. For what value of b is

$$g(x) = \begin{cases} ax + 2b, & x \leq 0 \\ x^2 + 3a - b, & 0 < x \leq 2 \\ 3x - 5, & x > 2 \end{cases}$$

continuous at every x ?

21. Explain why the equation $\cos x = x$ has at least one solution.

22. If $f(x) = x^3 - 8x + 10$, show that there is a value c for which $f(c) = 1000$.

23. Suppose that a function f is continuous on the closed interval $[0, 1]$ and that $0 \leq f(x) \leq 1$ for every x in $[0, 1]$. Show that there must exist a number c in $[0, 1]$ such that $f(c) = c$ (c is called a fixed point of f).
(Hint: Consider the function $g(x) = f(x) - x$ and try to find a zero of $g(x)$).

24. Show that the function $F(x) = (x - a)^2(x - b)^2 + x$ takes on the value $(a + b)/2$ for some value of x .

Chapter 2


Differentiation

2.1 Tangent Lines and Their Slopes

Problem: Find a straight line L that is tangent to a curve C at a point P .

“For simplicity, restrict ourselves to curves which are graphs of functions.”

How do we define the tangent line to a curve?



The slope of the line PQ is

$$\frac{f(x_0 + h) - f(x_0)}{h}.$$

Definition 2.1.1. Suppose f is cts at $x = x_0$ and

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h} = m$$

If the limit exists, then the line with equation

$$y = m(x - x_0) + f(x_0)$$

is called **the tangent line** to the graph of $y = f(x)$ at $P = (x_0, f(x_0))$. If the limit does not exist and $m = \infty$ or $m = -\infty$ then the tangent line is the vertical line $x = x_0$. If the limit does not exist and is not $\pm\infty$ then there is no tangent line at P .

Example 2.1.1. Find an equation of the tangent line to the curve $y = x^2$ at $(1, 1)$.

Solution. The slope is

$$m = \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h} = 2.$$


And an equation is $y = 2(x - 1) + 1$.

Example 2.1.2. Find an equation of the tangent line to the curve $y = x^{1/3} = \sqrt[3]{x}$ at the origin.

Solution. The slope of the tangent line is

$$m = \lim_{h \rightarrow 0} \frac{h^{1/3}}{h} = \infty.$$


So the tangent line is a vertical line $x = 0$ (in other words the y -axis).



Tangent lines to curves such as circles and parabolas do not cross these curves, they just touch at a single point. However, for graphs of functions tangent lines may cross the curve such as above. In fact at inflection points (which we will define later) they always do! For example the tangent line to the graph of $f(x) = x^3$ at $x = 0$ is the y -axis.

Example 2.1.3. Does $f(x) = x^{2/3}$ have a tangent line at $(0, 0)$?


Solution. The limit of the difference quotient is undefined at 0 since the right limit is ∞ while the left limit is $-\infty$. Hence the graph has no tangent line at $(0, 0)$.



"We say that this curve has a *cusp* at the origin. A *cusp* is an infinitely sharp point. If you were traveling along the curve, you would have to stop and turn 180° at the origin."

Example 2.1.4. Does $f(x) = |x|$ have a tangent line at $(0, 0)$?

Solution. The difference quotient is $\frac{|h|}{h}$ which has right limit 1 and left limit -1 at $h = 0$.



2.2 Derivative

Definition 2.2.1. The **derivative** of a function f at x is

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}.$$

whenever the limit exists. If $f'(x)$ exists, f is called **differentiable** at x .

$f'(x)$ is the slope of the tangent line to the graph of f at $(x, f(x))$.

We will regard f' as a function whose domain is those x at which f is differentiable.

Another way of defining derivative is

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Two limits are equivalent. This can be seen by letting $x = x_0 + h$.

Example 2.2.1. Show that the derivative of the linear function $f(x) = ax + b$ is $f'(x) = a$. In particular the derivative of a constant function is zero.

Example 2.2.2. Use the definition of the derivative to calculate the derivatives of a) $f(x) = x^2$, b) $f(x) = \frac{1}{x}$, c) $f(x) = \sqrt{x}$.

The previous three formulas are special cases of the following **Power Rule for Derivative**:

$$f(x) = x^r \implies f'(x) = r x^{r-1}$$

whenever x^{r-1} makes sense.

Proof of the Power Rule for positive integers. Let $f(x) = x^n$ and n a positive integer. Then

$$\begin{aligned} f'(x_0) &= \lim_{x \rightarrow x_0} \frac{x^n - x_0^n}{x - x_0} = \lim_{x \rightarrow x_0} \frac{(x - x_0)(x^{n-1} + x^{n-2}x_0 + \cdots + x x_0^{n-1} + x_0^{n-1})}{x - x_0} \\ &= \lim_{x \rightarrow x_0} (x^{n-1} + x^{n-2}x_0 + \cdots + x x_0^{n-1} + x_0^{n-1}) = n x_0^{n-1} \end{aligned}$$

We will prove the general version later.

Example 2.2.3.

$$f(x) = x^{5/3} \implies f'(x) = x^{2/3},$$

for all x . How about $f'(-1/8)$?

$$f(x) = \frac{1}{\sqrt{x}} \implies f'(x) = -\frac{1}{2}x^{-3/2}$$

for $x > 0$.

Example 2.2.4. Differentiate the absolute value function $f(x) = |x|$ to get

$$f'(x) = \operatorname{sgn}(x) = \begin{cases} -1, & \text{if } x < 0 \\ 1, & \text{if } x > 0 \end{cases}$$

Note that f is not differentiable at 0.

Example 2.2.5. How should the function $f(x) = x\operatorname{sgn}(x)$ be defined at $x = 0$ so that it is continuous there? Is it then differentiable there?

Notations for Derivative

Let $y = f(x)$. We denote the derivative by

$$y' = f'(x) = \frac{dy}{dx} = \frac{d}{dx}f(x).$$

If we want to evaluate the derivative at point x_0

$$y' |_{x=x_0} = f'(x_0) = \frac{dy}{dx} |_{x=x_0} = \frac{d}{dx}f(x) |_{x=x_0}.$$

The notations y' and $f'(x)$ are *Lagrange notations* for the derivative. The notations $\frac{dy}{dx}$ and $\frac{d}{dx}f(x)$ are called *Leibniz notations* for the derivative.

The Leibniz notation is suggested by the definition of the derivative. Let $\Delta y = f(x+h) - f(x)$ be the increment in y and $\Delta x = x+h - x = h$ be the increment in x . Then

$$\frac{dy}{dx} = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x}$$

2.3 Differentiation Rules

Differentiability is stronger than continuity.

Theorem 2.3.1. If f is differentiable at x then f is cts at x .

Proof.

$$\lim_{h \rightarrow 0} (f(x+h) - f(x)) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \lim_{h \rightarrow 0} h = f'(x)0 = 0$$

This means

$$0 = \lim_{h \rightarrow 0} f(x+h) - \lim_{h \rightarrow 0} f(x) = \lim_{h \rightarrow 0} f(x+h) - f(x)$$

Hence

$$\lim_{h \rightarrow 0} f(x+h) = f(x)$$

□

Theorem 2.3.2. *If f and g are differentiable at x then*

$$(f+g)'(x) = f'(x) + g'(x),$$

$$(f-g)'(x) = f'(x) - g'(x),$$

and for any constant c

$$(cf)'(x) = cf'(x).$$

Proof. Let's prove the derivative of sums is sum of derivatives. The others are similar.

$$\begin{aligned} (f+g)'(x) &= \lim_{h \rightarrow 0} \frac{(f+g)(x+h) - (f+g)(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h) + g(x+h) - f(x) - g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} + \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} = f'(x) + g'(x), \end{aligned}$$

□

The sum rule extends to any number of functions.

$$(f_1 + \cdots + f_n)'(x) = f_1'(x) + \cdots + f_n'(x).$$

Example 2.3.1. *Take the derivative of*

$$f(x) = 5\sqrt{x} + \frac{3}{x} - 19$$

It is NOT true that derivative of product of functions is a product of their derivatives. Usually $(fg)'(x) \neq f'(x)g(x)$.

Theorem 2.3.3. *If f and g are differentiable at x then*

$$(fg)'(x) = f'(x)g(x) + f(x)g'(x).$$

Proof.

$$\begin{aligned} (fg)'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} = \lim_{h \rightarrow 0} \frac{(f(x+h) - f(x))g(x+h) + f(x)(g(x+h) - g(x))}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \lim_{h \rightarrow 0} g(x+h) + \lim_{h \rightarrow 0} f(x) \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \end{aligned}$$

□

Example 2.3.2. *Find the derivative of $f(x) = (x^2 + x + 1)(2x + \frac{1}{x})$.*

The product rule can be extended to any number of functions

$$(f_1 f_2 f_3)' = f_1' f_2 f_3 + f_1 f_2' f_3 + f_1 f_2 f_3'$$

$$(f_1 \cdots f_n)' = f_1' f_2 \cdots f_n + f_1 f_2' f_3 \cdots f_n + \cdots + f_1 \cdots f_{n-1} f_n'$$

Theorem 2.3.4. If f is differentiable at x and $f(x) \neq 0$ then $1/f$ is diff at x , and

$$\left(\frac{1}{f}\right)'(x) = \frac{-f'(x)}{f(x)^2}.$$

Proof.

$$\frac{d}{dx} \frac{1}{f(x)} = \lim_{h \rightarrow 0} \frac{\frac{1}{f(x+h)} - \frac{1}{f(x)}}{h} = \lim_{h \rightarrow 0} \frac{-1}{f(x+h)f(x)} \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

The result follows by limit rules and continuity of f . □

Example 2.3.3. Differentiate $y = \frac{x^5}{x^{2/3} + 1}$.

Theorem 2.3.5. If f and g are differentiable at x and $g(x) \neq 0$ then

$$\left(\frac{f}{g}\right)'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{g^2(x)}$$

Proof. Using the product rule and reciprocal rule,

$$\left(\frac{f}{g}\right)'(x) = \left(\frac{1}{g}(x)f(x)\right)' = \frac{f'(x)g(x) - f(x)g'(x)}{g^2(x)}$$

□

Example 2.3.4. Find the derivative of $f(x) = \frac{a+bx}{m+cx}$.

Example 2.3.5. Find an equation of the tangent line to $y = \frac{2}{3-4\sqrt{x}}$ at the point $(1, -2)$.

Solution. Let us define $g(x) = 3 - 4\sqrt{x}$. Then $g'(x) = -4 \frac{1}{2\sqrt{x}} = -\frac{2}{\sqrt{x}}$ and

$$y' = 2 \frac{-g'(x)}{g(x)^2} = 2 \frac{\frac{2}{\sqrt{x}}}{(3-4\sqrt{x})^2} = \frac{4}{\sqrt{x}(3-4\sqrt{x})^2}$$

Hence $y'(1) = 4$. And the equation of the tangent line is $y = 4(x-1) - 2$.

Example 2.3.6. Find the x -coordinates of points on the curve $y = \frac{x+1}{x+2}$ where the tangent line is parallel to the line $y = 4x$.

Solution. Solving $y' = 4$, we find $x = -3/2$ and $x = -5/2$.

Example 2.3.7. If $f(2) = 2$ and $f'(2) = 3$, calculate

$$\frac{d}{dx} \left(\frac{x^2}{f(x)} \right) \Big|_{x=2}$$

Solution. Answer is

$$\frac{2 \cdot 2 f(2) - 2^2 f'(2)}{f(2)^2} = \frac{8 - 12}{4} = -1.$$

2.4 Chain Rule

The following theorem is known as the chain rule.

Theorem 2.4.1. *If $f(u)$ is differentiable at $u = g(x)$ and $g(x)$ is differentiable at x , then*

$$(f \circ g)'(x) = f'(g(x))g'(x)$$

Proof. The proof in the case $g(x) \neq g(a)$ for x sufficiently close to a .

$$(f \circ g)'(a) = \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} = \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \cdot \frac{g(x) - g(a)}{x - a} = f'(g(a))g'(a)$$

If there is always an $x \neq a$, x near a such that $g(x) = g(a)$ then the above proof fails due to division by zero. You can find the proof in this case in Wikipedia. \square

In Leibniz notation, if $y = f(u)$ where $u = g(x)$ then

$$y = f(g(x)) = (f \circ g)(x)$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$

where $\frac{dy}{du}$ is evaluated at $u = g(x)$.

Example 2.4.1. *Find the derivative of $y = \sqrt{x^2 + 1}$.*

Solution. Here $y = f(g(x))$ where $f(u) = \sqrt{u}$ and $u = x^2 + 1$.

$$\frac{dy}{dx} = f'(g(x))g'(x) = \frac{1}{2\sqrt{g(x)}}g'(x) = \frac{1}{2\sqrt{x^2 + 1}}2x = \frac{x}{\sqrt{x^2 + 1}}.$$

Example 2.4.2. *Differentiate $y = (x^3 - 1)^{1000}$.*

Solution. Let $u = (x^3 - 1)$ then $y = u^{1000}$. $y' = 1000u^{999}u' = 1000(x^3 - 1)^{999}3x^2$.

Example 2.4.3.

$$\frac{d}{dx}|x| = \frac{d}{dx}\sqrt{x^2} = \frac{1}{2\sqrt{x^2}}2x = \frac{x}{\sqrt{x^2}} = \frac{x}{|x|}, \quad x \neq 0$$

This function is called sign function or signum function.

$$\text{sgn}(x) = \frac{x}{|x|} = \begin{cases} 1, & x > 0 \\ -1, & x < 0 \end{cases}.$$

Example 2.4.4. *Express in terms of f and f' .*

a) $\frac{d}{dx}f(x^2),$

b) $\frac{d}{dx}(f(\pi - 2f(x)))^4.$

Solution. For (a)

$$\frac{d}{dx}f(x^2) = f'(x^2)2x$$

For (b)

$$\frac{d}{dx}[f(\pi - 2f(x))]^4 = 4[f(\pi - 2f(x))]^3 f'(\pi - 2f(x))(-2f'(x)).$$

Example 2.4.5. For

$$f(x) = \left(1 + \sqrt{2x+1}\right)^{-4/3}$$

evaluate $f'(0)$.

Solution.

$$\begin{aligned} f'(x) &= \frac{-4}{3}(1 + \sqrt{2x+1})^{-7/3} \frac{d}{dx} \sqrt{2x+1} = \frac{-4}{3}(1 + \sqrt{2x+1})^{-7/3} \frac{1}{2\sqrt{2x+1}} \frac{d}{dx}(2x+1) \\ &= \frac{-4}{3}(1 + \sqrt{2x+1})^{-7/3} \frac{1}{2\sqrt{2x+1}} 2 \end{aligned}$$

Hence

$$f'(0) = -\frac{1}{2^{1/3}3}.$$

Example 2.4.6. Find an equation of the tangent line to the graph of

$$y = (1 + x^{2/3})^{3/2}$$

at $x = -1$.

Solution.

$$y' = \frac{3}{2}(1 + x^{2/3})^{1/2} \frac{2}{3}x^{-1/3}.$$

$$y'(-1) = \frac{3}{2}(1 + 1)^{1/2} \frac{2}{3}(-1) = -\sqrt{2}.$$

2.5 Derivatives of Trigonometric Functions

The radian measure of an angle is defined to be the length of the arc of a unit circle corresponding to that angle.

$$\text{angle in degrees} = \text{angle in radians} \cdot \frac{180^\circ}{\pi}.$$

In calculus all angles are measured in radians. By an angle of $\pi/3$ we mean $\pi/3$ radians or 60° not $(\pi/3)^\circ \approx 1.04^\circ$.

Theorem 2.5.1. $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1.$

Proof.

Suppose $0 < \theta < \frac{\pi}{2}$.

Area of OQP triangle is $\frac{1}{2} \sin \theta \cos \theta$.

Area of OAP arc is $\frac{\theta}{2\pi} \pi 1^2$.

Area of OAT triangle is $\frac{1}{2} \tan \theta = \frac{\sin \theta}{2 \cos \theta}$.

$$\frac{1}{2} \sin \theta \cos \theta \leq \frac{\theta}{2} \leq \frac{\sin \theta}{2 \cos \theta}$$

Multiply by $\frac{2}{\sin \theta} > 0$

$$\cos \theta \leq \frac{\theta}{\sin \theta} \leq \frac{1}{\cos \theta}$$

Take reciprocal to get

$$\cos \theta \leq \frac{\sin \theta}{\theta} \leq \frac{1}{\cos \theta},$$

for $0 < \theta < \frac{\pi}{2}$.

Use the squeeze theorem to show that

$$\lim_{\theta \rightarrow 0^+} \frac{\sin \theta}{\theta} = 1$$

Similarly, we can show that (2.1) holds for $-\frac{\pi}{2} < \theta < 0$ and hence

$$\lim_{\theta \rightarrow 0^-} \frac{\sin \theta}{\theta} = 1$$

□


Example 2.5.1. Show that $\lim_{h \rightarrow 0} \frac{\cos h - 1}{h} = 0$.

Solution.

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} &= \lim_{h \rightarrow 0} \frac{(\cos h - 1)(\cos h + 1)}{h(\cos h + 1)} = \lim_{h \rightarrow 0} \frac{\cos^2 h - 1}{h(\cos h + 1)} \\ &= \lim_{h \rightarrow 0} \frac{-\sin^2 h}{h(\cos h + 1)} = -\lim_{h \rightarrow 0} \frac{\sin h}{h} \frac{\sin h}{\cos h + 1} = -1 \cdot 0 = 0 \end{aligned}$$

Example 2.5.2. Find the limit of

- $\lim_{x \rightarrow 0} \frac{\sin x}{\sin 2x}$
- $\lim_{x \rightarrow 0} \frac{x \sin x}{2 - 2 \cos x}$
- $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2}$
- $\lim_{x \rightarrow 0} \frac{\tan 2x}{x}$



Theorem 2.5.2. $\sin x$ is differentiable for every x and

$$\frac{d}{dx} \sin x = \cos x$$

Proof.

$$\begin{aligned} \frac{d}{dx} \sin x &= \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h} = \lim_{h \rightarrow 0} \frac{\sin x \cos h + \cos x \sin h - \sin x}{h} \\ &= \lim_{h \rightarrow 0} \frac{\sin x(\cos h - 1)}{h} + \lim_{h \rightarrow 0} \frac{\cos x \sin h}{h} = \sin x \lim_{h \rightarrow 0} \frac{(\cos h - 1)}{h} + \cos x \lim_{h \rightarrow 0} \frac{\sin h}{h} = \cos x \end{aligned}$$

□

Theorem 2.5.3. $\cos x$ is differentiable for every x and

$$\frac{d}{dx} \cos x = -\sin x.$$

Proof.

$$\frac{d}{dx} \cos x = \frac{d}{dx} \sin\left(\frac{\pi}{2} - x\right) = -\cos\left(\frac{\pi}{2} - x\right) = -\sin x.$$

□

Example 2.5.3. Evaluate the derivative of

a) $\sin(\pi x) + \cos(3x),$

b) $x^2 \cos(\sqrt{x}),$

c) $\frac{\cos x}{1 - \sin x}$

The derivatives of the other trigonometric functions

$$\tan x = \frac{\sin x}{\cos x}, \quad \sec x = \frac{1}{\cos x}, \quad \cot x = \frac{\cos x}{\sin x}, \quad \csc x = \frac{1}{\sin x}.$$

Since \cos and \sin are everywhere differentiable, the above functions are differentiable everywhere except where their denominators are zero. The derivatives of these functions can be derived by using quotient and reciprocal rules.

$$\frac{d}{dx} \tan x = \sec^2 x, \quad \frac{d}{dx} \sec x = \sec x \tan x, \quad \frac{d}{dx} \cot x = -\csc^2 x, \quad \frac{d}{dx} \csc x = -\csc x \cot x.$$

Example 2.5.4. Verify the derivative formulas for $\tan x$ and $\sec x$.

Example 2.5.5. Find the derivative of $y = \sin(\cos(\tan t))$.

Example 2.5.6. Find the points on the curve $y = \tan(2x)$, $-\pi/4 < x < \pi/4$, where the normal is parallel to the line $y = -x/8$.

Exercises

1. $\lim_{x \rightarrow \infty} \frac{\sin x}{x}$. (Hint: Use Sandwich Theorem)
2. $\lim_{t \rightarrow \infty} \frac{2 - t + \sin t}{t + \cos t}$ (Hint: Divide both sides by t and use the previous exercise)

2.6 Higher Order Derivatives

Derivative of derivative is called **second derivative**. If $y = f(x)$ then

$$y'' = f''(x) = \frac{d}{dx} \frac{d}{dx} y = \frac{d^2}{dx^2} y = \frac{d^2}{dx^2} f(x).$$

Similar notations can be used for third, fourth, etc. derivatives. For n -th derivative, we write

$$y^{(n)} = f^{(n)}(x) = \frac{d^n y}{dx^n}$$

Example 2.6.1. Calculate all the derivatives of $y = x^3$.

Example 2.6.2. Calculate all the derivatives of $y = x^n$ where n is a positive integer.

Solution.

$$y^{(k)} = \begin{cases} \frac{n!}{(n-k)!} x^{n-k} & \text{if } 0 \leq k \leq n \\ 0 & \text{if } k > n \end{cases}$$

Example 2.6.3. Show that if A , B and k are constants, then the function $y = A \cos(kt) + B \sin(kt)$ is a solution of the second order differential equation

$$\frac{d^2 y}{dx^2} + k^2 y = 0.$$

Example 2.6.4. If $y = \tan kx$ show that $y'' = 2k^2 y(1 + y^2)$.

Example 2.6.5. If f and g are twice differentiable functions, show that

$$(fg)'' = f''g + 2f'g' + fg''.$$

What do you think about the general formula for $\frac{d^n}{dx^n}(fg)$?

2.7 Mean Value Theorem

The Mean Value Theorem is the midwife of calculus - not very important or glamorous by itself, but often helping to deliver other theorems that are of major significance.

– E. Purcell and D. Varberg

Suppose you drive in 2 hours from city A to city B which are 200km apart. That means your average speed was 100km/h. Even if you did not travel constant speed, there was at least one instant where your speed was exactly 100km/h. This is called **mean value theorem**.

Theorem 2.7.1 (The Mean-Value Theorem). Suppose that f is continuous on the interval $[a, b]$ and that it is differentiable on the open interval (a, b) . Then there exists a point c in the open interval (a, b) s.t.

$$\frac{f(b) - f(a)}{b - a} = f'(c).$$




Figure 2.1: Mean Value Theorem says that the slope of the secant line joining two points on the graph of $f(x)$ is equal to the slope of the tangent line at some point $x = c$ between a and b .

Let $f(t)$ denote the distance from city A. Then $f(0) = 0$ and $f(2) = 200$. Mean Value Theorem says there is a time $t = c$ s.t. $f'(c) = 100$.

Example 2.7.1. Let $f(x) = |x|$ on $[-1, 1]$. Show that there is no $c \in [-1, 1]$ satisfying the conclusion of the Mean Value Theorem. Why?

The Mean Value Theorem is an *existence theorem* like Intermediate Value Theorem. In particular

- We don't know how to find c .
- We don't know how many different c can be found satisfying Mean Value Theorem (there is at least one).




Figure 2.2: There may be more than one c satisfying the conclusion of the Mean Value Theorem.

Example 2.7.2. Show that $\sin x < x$ for all $x > 0$.

Solution. For $x > 2\pi$, we have $\sin x < 1 < 2\pi < x$. Now assume $0 < x < 2\pi$. By the Mean Value Theorem there exists c , $0 < c < x$ such that

$$\frac{\sin x - \sin 0}{x - 0} = \cos c.$$

Hence $\sin x = x \cos c$. Since $0 < c < 2\pi$, $\cos c < 1$. Since also $x > 0$, we have $x \cos c < x$. So $\sin x = x \cos c < x$.

Example 2.7.3. Show that $\sqrt{1+x} < 1 + \frac{x}{2}$ for all $x > 0$.

Solution. Let $f(x) = \sqrt{1+x}$. Then $f'(c) < \frac{1}{2}$ for $c > 0$. Use Mean Value Theorem.

Example 2.7.4. Determine all the numbers c which satisfy the conclusions of the Mean Value Theorem for

$$f(x) = x^3 + 2x^2 - x, \quad x \in [-1, 2]$$

Solution. Solve

$$3c^2 + 4c - 1 = f'(c) = \frac{f(2) - f(-1)}{2 - (-1)} = \frac{14 - 2}{3} = 4$$

Solutions of $3c^2 + 4c - 5 = 0$ are

$$c_{\pm} = \frac{-4 \pm \sqrt{76}}{6}.$$

Notice that only $\frac{-4 + \sqrt{76}}{6}$ lies in $[-1, 2]$.

Example 2.7.5. Suppose f is continuous and differentiable on $[3, 9]$. Suppose $f(3) = -4$, and $f'(x) \leq 10$ for all x . What is the largest value possible for $f(9)$?

Solution. By Mean Value Theorem, there exists $c \in (3, 9)$ such that

$$f(9) - f(3) = f'(c)(9 - 3) \leq 10 \times 6 = 60.$$

So $f(9) \leq 60 + f(3) = 56$.

Definition 2.7.1. Suppose f is defined on an interval I . If for all x_1, x_2 in I s.t. $x_2 > x_1$,

If	Then on I , f is
$f(x_2) > f(x_1)$	increasing
$f(x_2) < f(x_1)$	decreasing
$f(x_2) \geq f(x_1)$	non-decreasing
$f(x_2) \leq f(x_1)$	non-increasing

Theorem 2.7.2. Suppose f is differentiable on an open interval I . If for all $x \in I$,


If	Then on I , f is
$f'(x) > 0$	increasing
$f'(x) < 0$	decreasing
$f'(x) \geq 0$	non-decreasing
$f'(x) \leq 0$	non-increasing

Proof. Let's prove the first statement. Let $x_2 > x_1$ in I . By the Mean Value Theorem, there exists c , $x_1 < c < x_2$, such that $f(x_2) - f(x_1) = f'(c)(x_2 - x_1)$. Since $f'(c) > 0$ and $x_2 - x_1 > 0$, we have $f(x_2) > f(x_1)$. So f is increasing. \square

Example 2.7.6. On what intervals is $f(x) = x^3 - 12x + 1$ increasing or decreasing?

Solution.

$f'(x) = 3(x-2)(x+2)$. So f is decreasing on $(-2, 2)$ and increasing otherwise.



We know that if f is a constant function then its derivative is zero. The converse is also true.

Theorem 2.7.3. If $f'(x) = 0$ on an interval I then $f(x)$ is constant on I .

Proof. Choose x_0 in I . Let $C = f(x_0)$. If x is any other point in I then by Mean Value Theorem, $f(x) - f(x_0) = f'(c)(x - x_0) = 0$. \square

Challenge Problem. Suppose that $f, g : \mathbb{R} \rightarrow \mathbb{R}$ are differentiable functions such that the limits $L = \lim_{x \rightarrow -\infty} f(x)$ and $M = \lim_{x \rightarrow -\infty} g(x)$ exist and $L \leq M$. Is it possible that $f(1) = 4$, and $g(1) = 2$? **Hint:** Consider $h(x) = g(x) - f(x)$ and use MVT.

2.8 Implicit Differentiation

We learned to find the slope of a curve that is the graph of a function. But not all curves are graphs of functions, for example the circle $x^2 + y^2 = 1$.

Curves are graphs of equations in two variables

$$F(x, y) = 0.$$

For the circle $F(x, y) = x^2 + y^2 - 1$.

Example 2.8.1. Find the slope of the circle $x^2 + y^2 = 25$ at the point $(3, -4)$.

Solution. *1st method.* Solve the equation $x^2 + y^2 = 1$ for y . There are two solutions $y_{1,2} = \pm\sqrt{25 - x^2}$. The point lies on the graph of y_2 . Take derivative of y_2 .

2nd method. To differentiate with respect to x treat y as a function of x and use Chain Rule.

$$\frac{d}{dx}(x^2 + y(x)^2) = \frac{d}{dx}0 = 0.$$

This gives

$$2x + 2y(x) \frac{dy(x)}{dx} = 0$$

or

$$\frac{dy}{dx} = -\frac{2x}{2y}$$

Plug in $x = 3$, $y = -4$ to find $\frac{dy}{dx} = 3/4$.

This second method is known as the **implicit differentiation**.

Example 2.8.2. Find an equation of the tangent line to the curve $x \sin(xy - y^2) = 0$ at $(1, 1)$

Example 2.8.3. Find y'' in terms of x and y if $xy + y^2 = 2x$.

The General Power Rule for Derivative

So far, we proved the following rule

$$\frac{dx^r}{dx} = rx^{r-1}$$

for integer exponents r and a few special exponents such as $r = 1/2$. Using the implicit differentiation, we can give a proof for any rational exponent $r = m/n$ where $n \neq 0$.

If $y = x^{m/n}$ then $y^n = x^m$. Differentiating implicitly

$$ny^{n-1} \frac{dy}{dx} = \frac{dy^n}{dx} = \frac{dx^m}{dx} = mx^{m-1}$$

Hence

$$\frac{dy}{dx} = \frac{m}{n} x^{m-1} y^{1-n} = rx^{m-1} x^{(1-n)m/n} = rx^{m-1+r-m} = rx^r.$$

2.9 Exam 1 Review

Section	Exercises
1.2	7-36, 37-42, 43-46, 49-60, 74, 75
1.3	1-10, 11-34
1.4	17, 18, 29, 31
1.5	7-10
2.1	1-12, 13-17, 18-24
2.2	11-24 (ignore differentials), 30-33, 34-39, 40-49
2.3	1-50
2.4	1-16, 30-34, 36-39
2.5	1-36, 39-42, 45-46
2.6	1-12
2.8	1-3, 5-7, 8-15
2.9	1-8, 9-16

Table 2.1: Exam 1 Review Problems from Adams & Essex Calculus: A Complete Course 7th Edition

Sample Exam 1

1. Find the following limits if they exist.

a)

$$\lim_{x \rightarrow \infty} \left(\frac{x^2}{x+1} - \frac{x^2}{x-1} \right)$$

b)

$$\lim_{x \rightarrow 0} \frac{x}{|x-1| - |x+1|}$$

c)

$$\lim_{x \rightarrow -\infty} \left(x + \sqrt{x^2 - 4x + 1} \right)$$

2. Show that $f(x) = x^3 + x - 1$ has a zero between $x = 0$ and $x = 1$.
3. Find the slope of the tangent line to the curve

$$\tan(xy^2) = \frac{2xy}{\pi}$$

at the point $(-\pi, 1/2)$.

4. Calculate the derivatives.

a)

$$y = \frac{1 + \sqrt{x}}{x^4 + 1}$$

b)

$$y = (\sin(\sqrt{x}) + 1)^3 + 7x \cos x$$

5. Is the function $y = |x - 2|$ differentiable at $x = 2$? Show your work using the limit definition of the derivative.

Chapter 3

Transcendental Functions

3.1 Inverse Functions

Definition 3.1.1. f is called **one-to-one** if $f(x_1) \neq f(x_2)$ whenever $x_1 \neq x_2$ or equivalently

$$f(x_1) = f(x_2) \implies x_1 = x_2$$




Figure 3.1: A function which is not 1-1.

Horizontal Line Test. Let $f: \mathbb{R} \rightarrow \mathbb{R}$. By definition of a function any vertical line intersects the graph at one point. f is 1-1 if its graph is never intersected by any horizontal line more than once.

Theorem 3.1.1. Increasing or decreasing functions are 1-1. Thus if $f'(x) > 0$ for all x in an interval I , then f is 1-1 on I . Similarly if $f'(x) < 0$ for all x in I then f is 1-1 on I .

Definition 3.1.2. If f is one-to-one then it has an inverse function f^{-1} defined as follows: If x is in the range of f then it is in the domain of f^{-1} and

$$f^{-1}(x) = y \iff x = f(y).$$

If f is not 1-1 then it is not invertible.

Given $y = f(x)$, to find the inverse function, we solve x in terms of y .

Example 3.1.1. Show that $f(x) = 2x - 1$ is one-to-one and find its inverse $f^{-1}(x)$.

Solution. Since $f'(x) = 2 > 0$, f is increasing on \mathbb{R} and therefore one-to-one for all x . Solve $y = f(x) = 2x - 1$ for x , to get

$$x = \frac{y+1}{2}$$

Then $x = f^{-1}(y) = \frac{y+1}{2}$ or

$$f^{-1}(x) = \frac{x+1}{2}.$$

Usually, we can not solve $y = f(x)$ for x . For example $y = x + x^3$ is 1-1 (check) but it is not possible solve it for x .

Properties of inverse functions

1. The domain of f^{-1} is the range of f .
2. The range of f^{-1} is the domain of f .
3. $f(f^{-1}(x)) = x$ for all x in the domain of f^{-1} .

Proof. If $f^{-1}(x) = y$ then $x = f(y)$ and $f(f^{-1}(x)) = f(y) = x$. □

4. $f^{-1}(f(x)) = x$ for all x in the domain of f .
5. $(f^{-1})^{-1}(x) = f(x)$ for all x in the domain of f . (The inverse of inverse of f is f .)

Proof.

$$(f^{-1})^{-1}(x) = y \iff f^{-1}(y) = x \iff y = f(x).$$

□

6. The graph of f^{-1} is the reflection of the graph of f in the line $x = y$. (Because if (a, b) is a point on the graph of $y = f(x)$ then (b, a) is a point on the graph of $y = f^{-1}(x)$).




Figure 3.2: The graph of the inverse function is a reflection along $y = x$.

Inverting Non One-to-one Functions by Restricting the Domain of Definition

The function $f(x) = x^2$ defined on all real numbers is not one-to-one because $(-a)^2 = a^2$ for any a . Hence f is not invertible.

In fact f is not invertible on any interval around $x = 0$.

Notice that $f'(0) = 0$.

Let us define a new function F by restricting the domain of f ,

$$F(x) = x^2, \quad x \geq 0.$$

Then $F^{-1}(x) = \sqrt{x}$.

Figure 3.3: The restriction of x^2 to $[0, \infty)$ and its inverse.

Conversely, since the range of the 1-1 function \sqrt{x} is $[0, \infty)$, the domain of its inverse $g(x) = x^2$ is $x \geq 0$.

Derivatives of Inverse Functions


Let $y = f^{-1}(x)$. Then $f(y) = x$. Considering y as a function of x and using implicit differentiation,

$$\frac{d}{dx}f(y) = \frac{d}{dx}x \implies f'(y)\frac{dy}{dx} = 1 \implies \frac{dy}{dx} = \frac{1}{f'(y)}$$

In short if $y = f^{-1}(x)$ then

$$(f^{-1})'(x) = \frac{1}{f'(y)} = \frac{1}{f'(f^{-1}(x))}$$

This formula says if $f'(y) \neq 0$ then f^{-1} is differentiable at x .



In Leibniz notation, $\frac{dy}{dx} = (f^{-1})'(x)$ while $\frac{dx}{dy} = f'(y)$, the above formula reads

$$\frac{dy}{dx} \frac{dx}{dy} = 1.$$

For example, if $y = x^2$, $x \geq 0$, then $x = \sqrt{y}$ and $\frac{dy}{dx} = 2x$ and $\frac{dx}{dy} = \frac{1}{2\sqrt{y}}$. So

$$\frac{dy}{dx} \frac{dx}{dy} = 2x \frac{1}{2\sqrt{y}} = 2x \frac{1}{2x} = 1.$$

Example 3.1.2. Show that $f(x) = x^3 + x$ is one-to-one on the whole real line and find $(f^{-1})'(10)$. Hint: $2^3 + 2 = 10$.

Solution. First $f'(x) = 3x^2 + 1 > 0$. Hence f is 1-1. Let $y = f^{-1}(x)$. Then $f(y) = x$, that is

$$y^3 + y = x$$

Take $\frac{d}{dx}$ of both sides to get

$$3y^2 y' + y' = 1 \implies y' = (f^{-1})'(x) = \frac{1}{3y^2 + 1}$$

Now if $y = f^{-1}(10)$ then $y^3 + y = 10$ and $y = 2$. So $(f^{-1})'(10) = \frac{1}{13}$.

Example 3.1.3. If $f(x) = 3x + x^3$, show that f has an inverse and find the slope of $y = f^{-1}(x)$ at $x = 0$.

3.2 Exponential and Logarithmic Functions

An **exponential function** is a function of the form $f(x) = a^x$ where the **base** a is a positive constant and the **exponent** x is the variable. Let's define this function.

- $a^0 = 1$.
- $a^n = a \cdot a \cdots a$ (n -times) if $n = 1, 2, 3, \dots$
- $a^{-n} = \frac{1}{a^n}$ if $n = 1, 2, 3, \dots$
- $a^{m/n} = \sqrt[n]{a^m}$ if $n = 1, 2, \dots$ and $m = \pm 1, \pm 2, \dots$

How should we define a^x if x is not rational? What does 2^π mean? We will define a^x for irrational x in the next section. For now, let us regard a^x as a limit as discussed in the next problem.

Example 3.2.1. Since the irrational number $\pi = 3.141592\dots$ is the limit of the sequence of rational numbers

$$r_1 = 3 \quad r_2 = 3.1 \quad r_3 = 3.14 \quad \dots$$

we can calculate 2^π as the limit of the sequence

$$2^3 = 8 \quad 2^{3.1} = 8.5741877\dots \quad 2^{3.14} = 8.8152409\dots$$

This gives

$$2^\pi = \lim_{n \rightarrow \infty} 2^{r_n} = 8.824977\dots$$

If x is irrational, then we define a^x as the limit values a^r for rational numbers r approaching x

$$a^x = \lim_{\substack{r \rightarrow x \\ r \text{ is rational}}} a^r.$$

Laws of Exponents

If $a > 0$ and $b > 0$ and x, y are real numbers then

1. $a^0 = 1$,
2. $a^{x+y} = a^x a^y$,
3. $a^{-x} = \frac{1}{a^x}$,

$$4. a^{x-y} = \frac{a^x}{a^y},$$

$$5. (a^x)^y = a^{xy},$$

$$6. (ab)^x = a^x b^x.$$


If $a > 1$ then

$$\lim_{x \rightarrow \infty} a^x = \infty, \quad \lim_{x \rightarrow -\infty} a^x = 0.$$

If $0 < a < 1$ then

$$\lim_{x \rightarrow \infty} a^x = 0, \quad \lim_{x \rightarrow -\infty} a^x = \infty.$$

The domain of a^x is $(-\infty, \infty)$ and its range is $(0, \infty)$.



Logarithm

If $a > 0$ and $a \neq 1$ then the function a^x is 1-1 (1^x has no inverse). The inverse function of a^x is $\log_a x$, called the **logarithm of x base a** .

$$y = \log_a x \iff x = a^y$$

Since a^x has domain $(-\infty, \infty)$, and range $(0, \infty)$, $\log_a x$ has domain $(0, \infty)$ and range $(-\infty, \infty)$.

Since a^x and $\log_a x$ are inverse functions

$$\log_a a^x = x \quad \forall x, \quad a^{\log_a x} = x, \quad x > 0$$




Figure 3.4: The graph of logarithmic function is a reflection of the graph of the exponential function in the line $y = x$.

Laws of Logarithm If $x > 0$, $y > 0$, $a > 0$, $b > 0$, $a \neq 1$, $b \neq 1$, then

1. $\log_a 1 = 0$
2. $\log_a(xy) = \log_a x + \log_a y$
3. $\log_a\left(\frac{1}{x}\right) = -\log_a x$
4. $\log_a\left(\frac{x}{y}\right) = \log_a x - \log_a y$
5. $\log_a x^y = y \log_a x$
6. $\log_a x = \frac{\log_b x}{\log_b a}$

Example 3.2.2. Prove $\log_a(xy) = \log_a x + \log_a y$ using laws of exponent.

Solution. Take $u = \log_a x$, $v = \log_a y$ then $x = a^u$, $y = a^v$ and

$$xy = a^{u+v} \iff u + v = \log(xy)$$

Example 3.2.3. Simplify

1. $\log_2 10 + \log_2 12 - \log_2 15$.

$$\log_2 10 + \log_2 12 - \log_2 15 = \log_2 \frac{10 \times 12}{15} = \log_2 8 = 3$$

2. $\log_{a^2} a^3$.

$$\log_{a^2} a^3 = \frac{\log_a a^3}{\log_a a^2} = \frac{3}{2}$$

3. $3^{\log_9 4}$.

$$3^{\log_9 4} = 3^{\frac{1}{2} \log_3 4} = 3^{\log_3 2} = 2$$

Example 3.2.4. Solve

$$3^{x-1} = 2^x,$$

in terms of $a = \log 2$ and $b = \log 3$.

Solution. Take logarithm base 3 of both sides.

$$(x-1)\log_3 3 = x\log_3 2 \iff x-1 = x\log_3 2 \iff x = \frac{1}{1-\log_3 2} = \frac{1}{1-a/b}$$

Numerically $x \approx 2.70951$.


3.3 The Natural Logarithm and Exponential

$f(x)$	$f'(x)$
$x^3/3$	x^2
$x^2/2$	x
x	1
x^0	0
$-x^{-1}$	x^{-2}
$-x^{-2}/2$	x^{-3}

Table 3.1: What is the mysterious function whose derivative is x^{-1} ?

Definition 3.3.1. For $x > 0$, let A_x be the area bounded by the curve $y = 1/t$, the t -axis and the vertical lines $t = 1$ and $t = x$. The **natural logarithm** function is defined by

$$\ln x = \begin{cases} A_x & x \geq 1 \\ -A_x & 0 < x < 1 \end{cases}$$



- Domain of $\ln x$ is $(0, \infty)$,
- $\ln 1 = 0$,
- $\ln x > 0$ if $x > 1$,
- $\ln x < 0$ if $0 < x < 1$,

Theorem 3.3.1. If $x > 0$ then $\frac{d}{dx} \ln x = \frac{1}{x}$

For $h > 0$, $\ln(x+h) - \ln x$ is the area under $1/t$ between $t = x$ and $t = x+h$. Thus

$$\frac{h}{x+h} < \ln(x+h) - \ln x < \frac{h}{x}$$


Thus

Proof.
$$\frac{1}{x+h} < \frac{\ln(x+h) - \ln x}{h} < \frac{1}{x}$$

Now use Squeeze Theorem to get

$$\lim_{h \rightarrow 0+} \frac{\ln(x+h) - \ln x}{h} = \frac{1}{x}$$

Similar argument holds for $h < 0$.



□

Theorem 3.3.2. If $x \neq 0$ then

$$\frac{d}{dx} \ln |x| = \frac{1}{x},$$

and

$$\int \frac{1}{x} dx = \ln|x| + C.$$

Proof. If $x < 0$, then by Chain Rule,

$$\frac{d}{dx} \ln|x| = \frac{d}{dx} \ln(-x) = \frac{1}{-x}(-1) = \frac{1}{x}.$$

□

This also shows that $\ln x$ is an **increasing** function for all $x > 0$.

Example 3.3.1. Find the derivatives of

1. $y = \ln|\cos x|$

2. $y = \ln(x + \sqrt{x^2 + 1})$

Solution. For (1),

$$y' = \frac{1}{\cos x}(-\sin x) = -\tan x.$$

For (2),

$$y' = \frac{1}{\sqrt{x^2 + 1}}.$$

The natural logarithm function $\ln x$ satisfies all the rules that the regular logarithms satisfy, that's why we call it natural log after all!

Theorem 3.3.3. 1. $\ln(xy) = \ln x + \ln y$.

2. $\ln(1/x) = -\ln x$.

3. $\ln(x/y) = \ln x - \ln y$.

4. $\ln x^r = r \ln x$.

Proof. For (i), if y is constant, then for all $x > 0$

$$\frac{d}{dx}(\ln(xy) - \ln x) = \frac{y}{xy} - \frac{1}{x} = 0$$

Thus for each $y > 0$, $\ln(xy) - \ln x = C$ (a constant depending on y) for $x > 0$. Setting $x = 1$ we get $C = \ln y$. The others can be done similarly (homework). □

Also note that

$$\ln 2^n = n \ln 2 \rightarrow \infty \text{ as } n \rightarrow \infty.$$

$$\ln 2^{-n} = -n \ln 2 \rightarrow -\infty \text{ as } n \rightarrow \infty.$$

This, combined with $\ln x$ is increasing shows that

$$\lim_{x \rightarrow \infty} \ln x = \infty, \quad \lim_{x \rightarrow 0^+} \ln x = -\infty.$$

Thus domain of $\ln x$ is $(0, \infty)$ and the range of $\ln x$ is $(-\infty, \infty)$.

The Exponential Function

Let $f(x) = \ln x$. Since $f'(x) = 1/x > 0 \implies f$ is increasing $\implies f$ is 1-1 $\implies f$ has an inverse. Call its inverse $\exp x$. Thus

$$\exp x = y \iff x = \ln y$$

- $\exp 0 = 1$ (since $\ln 1 = 0$),
- Domain of \exp is $(-\infty, \infty)$ (since range of \ln is $(-\infty, \infty)$),
- Range of $\exp =$ Domain of $\ln = (0, \infty)$,
- Cancellation identities

$$\exp \ln x = x, \quad x > 0$$

$$\ln \exp x = x, \quad -\infty < x < \infty.$$

Definition 3.3.2. $e = \exp(1) \approx 2.718\dots$

Thus $\ln e = 1$. Hence e is the number for which the area bounded by $y = 1/x$, the x -axis and the lines $x = 1$, $x = e$ is 1.

$$e^x = \exp(\ln(e^x)) = \exp(x \ln e) = \exp(x).$$

Since \exp is actually an exponential function, its inverse must be a logarithm

$$\ln x = \log_e x$$

The derivative of $y = e^x$ is calculated by implicit differentiation:

$$y = e^x \iff x = \ln y \iff 1 = \frac{y'}{y} \iff y' = y = e^x$$

This is a remarkable property:

$$\frac{d}{dx} e^x = e^x, \quad \int e^x dx = e^x + C$$

Example 3.3.2. Find the derivatives of

1. $e^{x^2-3x},$

2. $\sqrt{1+e^{2x}}$

General Exponentials and Logarithms

Definition 3.3.3. If $a > 0$ then for all real x , we define

$$a^x = e^{x \ln a}$$

This coincides with our previous definition that a^x is the limit of a^{r_n} where r_n are rational numbers tending to x .

Example 3.3.3. $2^\pi = e^{\pi \ln 2} \approx 8.825$.

Derivative of $y = a^x$.

$$\frac{d}{dx} a^x = \frac{d}{dx} e^{x \ln a} = e^{x \ln a} \ln a = a^x \ln a.$$

Example 3.3.4. Show that the graph of $f(x) = x^\pi - \pi^x$ has negative slope at $x = \pi$.

Solution. $f'(\pi) = \pi^\pi(1 - \ln \pi)$. Note that $\ln \pi > \ln e = 1$

Definition 3.3.4. Let $y = a^x$. Then $\frac{dy}{dx} = a^x \ln a$ which is negative if $0 < a < 1$ and positive if $a > 1$. Thus a^x is 1-1 and has an inverse function. We define its inverse as $\log_a x$.

Derivative of $y = \log_a x$.

$$\frac{d}{dx} \log_a x = \frac{d}{dx} \frac{\ln x}{\ln a} = \frac{1}{x \ln a}.$$

Logarithmic Differentiation

Example 3.3.5. Let $y = x^x$, $x > 0$. Find y' .

Solution. Neither the power rule $d/dx(x^a) = ax^{a-1}$ nor the exponential rule $d/dx(a^x) = \ln a a^x$ works.

$$\ln y = x \ln x \implies \frac{y'}{y} = 1 \ln x + x \frac{1}{x} \implies y' = x^x (\ln x + 1)$$

This technique is called **logarithmic differentiation** and is used to differentiate functions of the form $y = (f(x))^{g(x)}$ ($f(x) > 0$).

Example 3.3.6. Find dy/dt if $y = (\sin t)^{\ln t}$ where $0 < t < \pi$.

Solution.

$$y' = (\sin t)^{\ln t} \left(\frac{\ln \sin t}{t} + \ln t \cot t \right).$$

Example 3.3.7. If $y = \frac{(x+1)(x+2)(x+3)}{\sqrt{x+4}}$, find y' .

Solution. Since $(x+1)$ is not necessarily positive, $\ln(x+1)$ may or may not be defined. So we take the absolute value and then logarithm.

$$\ln |y| = \ln |x+1| + \ln |x+2| + \ln |x+3| - \frac{1}{2} \ln |x+4|$$

$$\frac{y'}{y} = \frac{1}{x+1} + \dots$$

Hyperbolic Functions

The hyperbolic sine function \sinh (pronounced 'zinch' (rhymes with 'pinch')) is

$$\sinh x = \frac{e^x - e^{-x}}{2}$$

and the hyperbolic cosine function \cosh (pronounced 'kosh' (rhymes with 'gosh')) is

$$\cosh x = \frac{e^x + e^{-x}}{2}$$

One can also define \tanh , \coth , etc.

These functions have many remarkable properties some of which resemble trigonometric functions. Show that

$$\begin{aligned} \cosh 0 &= 1, & \sinh 0 &= 0 \\ 2 \sinh x \cosh x &= \sinh 2x \\ \frac{d \sinh x}{dx} &= \cosh x, & \frac{d \cosh x}{dx} &= \sinh x. \end{aligned}$$

The reason these functions are called hyperbolic is

$$\cosh^2 x - \sinh^2 x = 1$$

So the parametric curve $(\cosh x, \sinh x)$ defines the hyperbola $x^2 - y^2 = 1$.

3.4 The Inverse Trigonometric Functions

The six trigonometric functions are periodic and hence not 1-1. However we can restrict their domains in such a way that the restricted functions are 1-1.

The $\arcsin x$ or $\arcsin x$ is the inverse of the $\sin x$ restricted to $[-\pi/2, \pi/2]$,

$$\begin{aligned} \sin(\arcsin y) &= y, & -1 \leq y \leq 1, \\ \arcsin(\sin x) &= x, & -\pi/2 \leq x \leq \pi/2. \end{aligned}$$




Figure 3.5: $f(x) = \arcsin x$ is a partial inverse of the sine function.

Example 3.4.1. *Simplify*

1. $\arcsin \frac{1}{2} = \frac{\pi}{6}$,
2. $\arcsin \frac{-\sqrt{2}}{2} = -\frac{\pi}{4}$,
3. $\arcsin 2$ is undefined since 2 is not in the range of sine.

Example 3.4.2. *Simplify*

1. $\sin(\arcsin 0.7) = 0.7$,
2. $\arcsin(\sin 3\pi/4) = \pi/4$,
3. $\cos(\arcsin 0.6) = 0.8$.

Solution. Let $\theta = \arcsin 0.6$. By the Pythagorean Theorem, $\cos \theta = 0.8$.

4. Similarly $\cos(\arcsin x) = \sqrt{1 - x^2}$.

Theorem 3.4.1.

$$\frac{d}{dx} \arcsin x = \frac{1}{\sqrt{1 - x^2}}.$$

Proof. Let $y = \arcsin x$ so that $x = \sin y$. Then

$$\frac{dy}{dx} = \frac{1}{\frac{dx}{dy}} = \frac{1}{\cos y} = \frac{1}{\sqrt{1 - x^2}}.$$

□

The Arctan Function

Define the $y = \arctan x$ to be the inverse of $y = \tan x$ on $(-\pi/2, \pi/2)$.

$$\begin{aligned} \tan(\arctan x) &= x, & -\infty < x < \infty, \\ \arctan(\tan x) &= x, & -\pi/2 < x < \pi/2. \end{aligned}$$




Figure 3.6: $f(x) = \arctan x$.

Example 3.4.3. 1. $\tan(\arctan 3) = 3$,

2. $\arctan(\tan \frac{3\pi}{4}) = \arctan -1 = -\frac{\pi}{4}$

3. $\cos(\arctan x) = \frac{1}{\sqrt{1+x^2}}$

Theorem 3.4.2. $\frac{d \arctan(x)}{dx} = \frac{1}{1+x^2}$

Proof. Let $y = \arctan x$ so that $x = \tan y$,

$$\frac{d \arctan(x)}{dx} = \frac{1}{\frac{dx}{dy}} = \frac{1}{\sec^2 y} = \frac{1}{1 + \tan^2 y} = \frac{1}{1 + x^2}.$$

□

Example 3.4.4. Find the slope of the curve $\arctan\left(\frac{2x}{y}\right) = \frac{\pi x}{y^2}$ at the point $(1, 2)$.

Solution. Taking $\frac{d}{dx}$ of both sides

$$\frac{1}{1 + \left(\frac{2x}{y}\right)^2} 2 \left(\frac{y - xy'}{y^2} \right) = \pi \left(\frac{y^2 - 2xyy'}{y^4} \right)$$

Plugging $x = 1, y = 2$,

$$2 - y' = \pi(1 - y') \implies y' = \frac{\pi - 2}{\pi - 1}.$$

Other inverse trigonometric functions

$\cos x$ is 1-1 on $[0, \pi]$ so we define $y = \arccos x$ as the inverse of $y = \cos x$ restricted to $[0, \pi]$.

$$y = \arccos x \iff x = \cos y \quad 0 \leq y \leq \pi.$$

Theorem 3.4.3.

$$\frac{d}{dx} \arccos x = -\frac{1}{\sqrt{1-x^2}}$$

For the derivative,

$$\frac{dy}{dx} = \frac{1}{\frac{dx}{dy}} = \frac{1}{-\sin y} = -\frac{1}{\sqrt{1-x^2}}$$

Note that

$$\frac{d}{dx} \arccos x = -\frac{d}{dx} \arcsin x$$

The inverse and the derivative of other trigonometric functions can be defined similarly.

Quiz Problems

Example 3.4.5. Simplify

1. $\cos(\arctan \frac{1}{2}) = \frac{2}{\sqrt{5}}$

2. $\tan(\arccos x) = \frac{\sqrt{1-x^2}}{x}$

Example 3.4.6. Show that $\frac{d}{dx} (\arcsin x^2)^{1/2} = \frac{x}{\sqrt{1-x^4} \sqrt{\arcsin x^2}}.$

Chapter 4

Applications of Derivatives

4.1 Related Rates

Example 4.1.1. How fast is the area of a rectangle changing if one side is 10cm long and is increasing at a rate of 2cm/s and the other side is 8cm long and is decreasing at a rate of 3cm/s?

Solution. The area A , and the lengths of sides x and y are functions of time t . Also $A = xy$. We are given $\frac{dx}{dt} = 2$, $\frac{dy}{dt} = -3$ when $x = 10$, $y = 8$.

Then

$$\frac{dA}{dt} = \frac{dx}{dt}y + x\frac{dy}{dt}$$

gives $\frac{dA}{dt} = -14$.

In the previous problem, notice that the average changes

$$\begin{aligned}\frac{A(1) - A(0)}{1 - 0} &= 12 \cdot 5 - 10 \cdot 8 = -20, \\ \frac{A(.5) - A(0)}{.5 - 0} &= 2(11 \cdot 6.5 - 10 \cdot 8) = -17 \\ \frac{A(.1) - A(0)}{.1 - 0} &= 10(10.2 \cdot 7.7 - 10 \cdot 8) = -14.6\end{aligned}$$

converge to the instantaneous rate we found.

This is possible because $A'(t) = -14 - 6t$. Hence A changes in a non-constant fashion even though x and y changes constantly. What we are computing in this problem is $A'(0) = -14$. And this result holds even if $\frac{dx}{dt}$ and $\frac{dy}{dt}$ are not constant. (maybe velocity of x is not constant and it accelerates according to $x(t) = 10 + 2t + t^2$)

Example 4.1.2. How fast is the surface area of a ball changing when the volume of the ball is $32\pi/3 \text{ cm}^3$ and is increasing at $2\text{cm}^3/\text{s}$? (The surface area of the ball is $A = 4\pi r^2$ and the volume is $V = \frac{4}{3}\pi r^3$. Note that $V(r) = \int_0^r A(r)dr$)

Solution. When $V = 32\pi/3$, $r = 2$, $\frac{dV}{dt} = 2$

$$\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

gives $\frac{dr}{dt} = \frac{1}{8\pi}$.
Now

$$\frac{dA}{dt} = 8\pi r \frac{dr}{dt} = 2$$

Example 4.1.3. A point is moving to the right along the first quadrant portion of the curve $x^2 y^3 = 72$. When the point has coordinates $(3, 2)$, its horizontal velocity is 2 units/s. What is its vertical velocity?

Solution. Taking d/dt of both sides

$$2x \frac{dx}{dt} y^3 + x^2 3y^2 \frac{dy}{dt} = 0$$

At $x = 3$, $y = 2$, $\frac{dx}{dt} = 2$,

$$\frac{dy}{dt} = -\frac{8}{3}.$$

Exercises.

- Find the rate of change of the area of a square whose side is 6 cm long, if the side length is increasing at 2 cm/min.
- Air is being pumped into a spherical balloon. The volume of the balloon is increasing at a rate

of $20 \text{ cm}^3/\text{s}$ when the radius is 30 cm. How fast is the radius increasing at that time? (The volume of a ball of radius r is $V = \frac{4}{3}\pi r^3$.)

- The area of a circle is decreasing at a rate of $2 \text{ cm}^2/\text{min}$. How fast is the radius of the circle changing when the area is 100 cm^2 ?

Answer:

4.2 Indeterminate Forms

To evaluate the limit $\lim_{x \rightarrow 0} \frac{\sin x}{x}$ we can not plug in $x = 0$. We call $\sin x/x$ an **indeterminate form** of $[0/0]$ at $x = 0$.

The limit of an indeterminate form $[0/0]$ can be any number.

$$\lim_{x \rightarrow 0} \frac{x}{x} = 1, \quad \lim_{x \rightarrow 0} \frac{x}{x^3} = \infty, \quad \lim_{x \rightarrow 0} \frac{x^3}{x^2} = 0.$$

There are other types of indeterminate forms $[\infty/\infty]$, $[0 \cdot \infty]$, $[\infty - \infty]$, $[0^\infty]$, $[\infty^0]$, $[1^\infty]$.

Theorem 4.2.1 (l'Hopital's Rules). Let f and g are differentiable on an interval containing a . Suppose that $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ are either both 0 or both $\pm\infty$. If $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$ exists then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}.$$

The results hold true if $\lim_{x \rightarrow a}$ is replaced by $\lim_{x \rightarrow a+}$ and $\lim_{x \rightarrow a-}$ or if $a = \pm\infty$.

Proof. Proof follows from generalized mean value theorem which we did not cover.

Let's give a proof for the following special case. Suppose $f(a) = g(a) = 0$, $g'(a) \neq 0$ and f, g have continuous derivatives at $x = a$. Then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}}{\lim_{x \rightarrow a} \frac{g(x) - g(a)}{x - a}} = \frac{f'(a)}{g'(a)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

□

Note that in applying l'Hopital's rule we calculate the quotient of the derivatives, not the derivative of the quotients.

Example 4.2.1. Evaluate

$$\lim_{x \rightarrow 1} \frac{\ln x}{x^2 - 1}$$

Solution.

$$\lim_{x \rightarrow 1} \frac{\ln x}{x^2 - 1} \quad \left[\frac{0}{0} \right]$$

$$\lim_{x \rightarrow 1} \frac{\ln x}{x^2 - 1} = \lim_{x \rightarrow 1} \frac{\frac{1}{x}}{2x} = \lim_{x \rightarrow 1} \frac{1}{2x^2} = \frac{1}{2}.$$

If one application of the l'Hopital's rule again gives an indeterminate form, we can apply it again.

Example 4.2.2. Evaluate

$$\lim_{x \rightarrow 0} \frac{2 \sin x - \sin(2x)}{2e^x - 2 - 2x - x^2}$$

Solution. Applying l'Hopital's rule three times we get the answer 3.

Example 4.2.3.

$$\lim_{x \rightarrow 1^+} \frac{x}{\ln x}$$

Solution. If you apply the l'Hopital's rule, you get the wrong answer of 1. This is not an indeterminate form, and you can't use l'Hopital's rule. The real answer is ∞ .

Example 4.2.4.

$$\lim_{x \rightarrow 0^+} \frac{1}{x} - \frac{1}{\sin x}$$

Solution. This is an indeterminate form of type $[\infty - \infty]$ which can be brought to the form $[0/0]$.

$$\lim_{x \rightarrow 0^+} \frac{1}{x} - \frac{1}{\sin x} = \lim_{x \rightarrow 0^+} \frac{\sin x - x}{x \sin x} = \lim_{x \rightarrow 0^+} \frac{\cos x - 1}{\sin x + x \cos x} = \lim_{x \rightarrow 0^+} \frac{-\sin x}{\cos x + \cos x - x \sin x} = \frac{0}{-2} = 0.$$

where we use l'Hopital's rule twice.

To deal with indeterminate forms of types $[0^0]$, $[\infty^0]$ and $[1^\infty]$, we take logarithms.

Example 4.2.5.

$$\lim_{x \rightarrow 0^+} x^x.$$

Solution. This is of the form $[0^0]$. Let $y = x^x$. Then

$$\lim_{x \rightarrow 0^+} \ln y = \lim_{x \rightarrow 0^+} x \ln x = \lim_{x \rightarrow 0^+} \frac{\ln x}{1/x} = \lim_{x \rightarrow 0^+} \frac{1/x}{-1/x^2} = 0$$

Since \ln is a continuous function

$$\begin{aligned} \ln \lim_{x \rightarrow 0^+} y &= \lim_{x \rightarrow 0^+} \ln y = 0, \\ \lim_{x \rightarrow 0^+} x^x &= e^0 = 1. \end{aligned}$$

Example 4.2.6. Evaluate

$$\lim_{x \rightarrow \infty} \left(1 + \sin \frac{3}{x}\right)^x$$

Solution. This is of the form $[1^\infty]$. Again first evaluate the limit of the logarithm. $y = \left(1 + \sin \frac{3}{x}\right)^x$.

$$\lim_{x \rightarrow \infty} \ln y = \lim_{x \rightarrow \infty} \frac{\ln \left(1 + \sin \frac{3}{x}\right)}{1/x} = 3$$

Hence

$$\lim_{x \rightarrow \infty} \left(1 + \sin \frac{3}{x}\right)^x = e^3.$$

Exercises.

1. $\lim_{x \rightarrow \infty} \frac{x^2}{e^x}.$

Answer: 0

2. $\lim_{x \rightarrow 0} \frac{x - \sin x}{x^3}$

Answer: 1/6

3. $\lim_{x \rightarrow 0} \frac{x - \sin x}{x - \tan x}$

Answer: $-\frac{1}{2}$

4. $\lim_{x \rightarrow 0^+} x^{\sqrt{x}}$

Answer: 1

5. $\lim_{x \rightarrow 1} \frac{\ln(ex) - 1}{\sin \pi x}$

Answer: $-\frac{1}{\pi}$

6. $\lim_{x \rightarrow 0^+} \frac{\csc x}{\ln x}$

Answer: ∞

7. $\lim_{x \rightarrow 0} (1 + \tan x)^{1/x}$

Answer: 1

4.3 Extreme Values

A function has an **absolute maximum value** $f(x_0)$ if $f(x) < f(x_0)$ holds for every x in its domain.

Similarly, define **absolute minimum value**.

If it has an absolute min/max, then that value may be achieved at more than one point. For example the function $\cos x$ attains its absolute max at $x = 2n\pi$ for any integer n .

A function may or may not have an absolute min/max value. For example the function $f(x) = x$, $0 < x < 1$ does not have an absolute maximum or minimum.

Recall from the section on continuous functions that,

A continuous function defined on a closed and bounded interval must have an absolute maximum and an absolute minimum.

Maximum and minimum values of a function are collectively referred to as **extreme values**.

Function f has a **local maximum** value $f(x_0)$ if there exists $h > 0$ such that $f(x) \leq f(x_0)$ whenever x is in the domain of f and $|x - x_0| < h$.

Similarly we define **local minimum**.

We define **critical points** of f where $f'(x) = 0$, **singular points** of f where x is in domain of f and $f'(x)$ does not exist.

Following theorem says where the extreme values are located.

Theorem 4.3.1. *If the function f is defined on an interval I and has a local max or local min at $x = x_0$ then x_0 **must** be either a critical point, a singular point or an endpoint of the interval.*

Proof. If $f(x_0)$ is a local extrema and x_0 is not an endpoint or singular point, then $f'(x_0) = 0$. Otherwise, either $f'(x_0) > 0$ which means f is increasing at x_0 or $f'(x_0) < 0$ which means f is decreasing at x_0 so that $f(x_0)$ is neither a local min nor local max. \square

This theorem does not say f must have a local min/max at every singular, critical or endpoint. For example for $f(x) = x^3$, $f'(0) = 0$ but $f(0)$ is not an extremum value.


Example 4.3.1. Find the maximum and minimum values of the function $g(x) = x^3 - 3x^2 - 9x + 2$ on the interval $-2 \leq x \leq 2$.

Solution. g is a continuous function defined on a closed and bounded interval so it must have an absolute minimum and absolute maximum.

Since g is a polynomial, it can't have singular points.

$g'(x) = 3(x^2 - 2x - 3) = 3(x+1)(x-3)$. $g'(x) = 0$ if $x = -1$ or $x = 3$. $x = 3$ is not in the domain, so we ignore it.


We check the values of $g(x)$ at endpoints and critical points, $g(-2) = 0$, $g(-1) = 7$, $g(2) = -20$. The maximum value is 7, the minimum value is -20.



Example 4.3.2. Find the maximum and minimum values of $h(x) = 3x^{2/3} - 2x$ on the interval $[-1, 1]$.

Solution. $h'(x) = 2(x^{-1/3} - 1)$. $h'(0)$ is undefined, 0 is a singular point of h . h has a critical point at $x = 1$ which is also an endpoint.

$h(-1) = 5$, $h(0) = 0$, $h(1) = 1$. h has maximum value 5 and minimum value 0.



The first derivative test

By investigating the sign of the first derivative we can determine whether an extrema is a local minimum or local maximum.

Example 4.3.3. Find the local and absolute extreme values of $f(x) = x^4 - 2x^2 - 3$ on the interval $[-2, 2]$. Sketch the graph of f .


Solution. $f'(x) = 2x(x^2 - 1) = 4x(x - 1)(x + 1)$. The critical points are 0, -1, 1. There are no singular points.

x	-2	-1	0	1	2
f'		-	+	-	+
f	max	\searrow	min	\nearrow	max

$$f(-2) = f(2) = 5, f(-1) = f(1) = -4, f(0) = -3.$$

Since f is continuous and defined on a closed and bounded interval, it must have an absolute min/max.

So 5 is the absolute maximum and -4 is the absolute minimum.




Example 4.3.4. Locate all extreme values of $f(x) = x\sqrt{2-x^2}$. Determine whether any of these extreme values are absolute. Sketch the graph.

Solution. Note that f has domain $[-\sqrt{2}, \sqrt{2}]$. $f'(x) = -2\frac{x^2-1}{\sqrt{2-x^2}}$. Critical points are ± 1 . Singular points are $\pm\sqrt{2}$ and endpoints are also $\pm\sqrt{2}$.

$f(\pm\sqrt{2}) = 0$, $f(-1) = -1$, $f(1) = 1$. Since f is continuous on a closed bounded interval it must have maximum value 1 and minimum value -1.

x	$-\sqrt{2}$	-1	1	$\sqrt{2}$
f'		-	+	-
f	max	\searrow	min	\searrow



4.4 Concavity and Inflections

We say f is **concave up** on an interval I if f' is increasing on I and **concave down** on I if f' decreasing on I .

Note that if f is concave up then f lies above its tangents and below its chords while if f is concave down then f lies below its tangents and above its chords.

If f changes its concavity at x_0 then we call x_0 and **inflection point**.

Theorem 4.4.1. Assume f is twice differentiable.

- If $f'' > 0$ on an interval I then f is concave up on I ,
- If $f'' < 0$ on an interval I then f is concave down on I ,

4.5 Graphs of Functions

Definition 4.5.1. The graph of $y = f(x)$ has a **vertical asymptote** at $x = a$ if either $\lim_{x \rightarrow a^-} f(x) = \pm\infty$ or $\lim_{x \rightarrow a^+} f(x) = \pm\infty$.

Definition 4.5.2. The graph of $y = f(x)$ has a **horizontal asymptote** at $y = L$ if either $\lim_{x \rightarrow \infty} f(x) = L$ or $\lim_{x \rightarrow -\infty} f(x) = L$.


Example 4.5.1. Find the vertical and the horizontal asymptotes of $f(x) = \frac{1}{x^2 - x}$.

Solution. The vertical asymptotes are $x = 0, x = 1$.

$$\lim_{x \rightarrow 0^-} \frac{1}{x^2 - x} = \infty, \quad \lim_{x \rightarrow 0^+} \frac{1}{x^2 - x} = -\infty,$$

$$\lim_{x \rightarrow 1^-} \frac{1}{x^2 - x} = -\infty, \quad \lim_{x \rightarrow 1^+} \frac{1}{x^2 - x} = \infty,$$

The function has a horizontal asymptote, $\lim_{x \rightarrow \infty} \frac{1}{x^2 - x} = \lim_{x \rightarrow -\infty} \frac{1}{x^2 - x} = 0$. This is a two-sided horizontal asymptote.




Example 4.5.2. $f(x) = e^x$ has a left horizontal asymptote $y = 0$, $\lim_{x \rightarrow -\infty} e^x = 0$.

Example 4.5.3. $f(x) = \tan^{-1} x$ has a two one sided limits, $\lim_{x \rightarrow \infty} \tan^{-1} x = \pi/2$ and $\lim_{x \rightarrow -\infty} \tan^{-1} x = -\pi/2$.

Definition 4.5.3. The straight line $y = ax + b$ ($a \neq 0$) is an **oblique asymptote** of the graph $y = f(x)$ if either $\lim_{x \rightarrow \infty} (f(x) - (ax + b)) = 0$ or $\lim_{x \rightarrow -\infty} (f(x) - (ax + b)) = 0$.

Example 4.5.4. Let $f(x) = \frac{x^2 + 1}{x} = x + \frac{1}{x}$. Then $\lim_{x \rightarrow \pm\infty} (f(x) - x) = 0$. Hence f has a two-sided oblique asymptote.



Asymptotes of rational function

Let $f(x) = \frac{P_m(x)}{Q_n(x)}$, where P_m and Q_n are polynomials of degree m and n respectively. Suppose that P_m and Q_n have no common linear factors. The graph of f has

1. a vertical asymptote at every position at every x for which $Q_n(x) = 0$.
2. a two-sided horizontal asymptote $y = 0$ only if $m < n$.

3. a two-sided horizontal asymptote $y = L$ only if $m = n$. L is the ratio of the coefficients of the highest degree terms in P_m and Q_n .
4. a two sided oblique asymptote only if $m = n + 1$.

Example 4.5.5. Find the oblique asymptote of $y = \frac{x^3}{x^2+x+1}$.

Solution. By polynomial division, we get $y = x - 1 + \frac{1}{x^2+x+1}$. $y = x - 1$ is the oblique asymptote.

Checklist For Curve Sketching

1. Examine $f(x)$ to find the domain, intercepts, asymptotes and even/odd symmetries.
2. Find points where $f'(x) = 0$ (critical points of f) and where $f'(x)$ is undefined (singular points of f).
3. Find points where $f''(x) = 0$ (critical points of f) and where $f''(x)$ is undefined (singular points of f).
4. Make a table to investigate the signs of $f'(x)$ and $f''(x)$ to find the intervals where f is increasing or decreasing and the intervals where f is concave up and down. Find also the extreme points and inflection points of the graph.

4.6 Extreme Value Problems


Example 4.6.1. Find the area of the largest rectangle that can be inscribed in a semicircle of radius R if one side of the rectangle lies along the diameter of the semicircle.

Solution. $(x/2)^2 + y^2 = R^2$. So

$$A = xy = x\sqrt{R^2 - (x/2)^2}.$$

$$\frac{dA}{dx} = \frac{2R^2 - x^2}{\sqrt{4R^2 - x^2}}$$

The derivative is zero when $x = \sqrt{2}R$. Use the first derivative test to see that this gives max area $A = R^2$.



Example 4.6.2. Find the shortest distance from the origin to the curve $x^2y^4 = 1$.

Solution. The distance is $\sqrt{x^2 + y^2}$. Instead of minimizing distance, an easier way is to minimize its square $x^2 + y^2$. Solving $x^2y^4 = 1$ for y and plugging into distance,

$$D(y) = \frac{1}{y^4} + y^2$$

Then $D'(y) = -\frac{4}{y^5} + 2y$. Solving $D'(y) = 0$ for y we get $y = 2^{1/6}$. Use the first derivative test to check that $D(2^{1/6}) = \frac{3}{\sqrt[3]{8}}$ is a minimum.

Example 4.6.3. A manufacturer has 100 tons of metal that he can sell now with a profit of \$5 a ton. For each week that he delays shipment, he can produce another 10 tons of metal. However, for each week he waits, the profit drops 25 cents a ton. If he can sell the metal at any time, when is the best time to sell so that his profit is maximized?

Solution. Let x be the number of weeks to wait.

Ship	Amount of metal	Profit per ton	Total profit
now	100	5	500
in x weeks	$100 + 10x$	$5 - 0.25x$	$500 + 25x - 0.25x^2$

$P(x) = 500 + 25x - 2.5x^2$. Solve $P'(x) = 0$ to find $x = 5$. And maximum profit is \$562.50.

Exercises.

Answer: $\frac{10R}{\sqrt{5}}$

1. Find the shortest distance from the point $(8, 1)$ to the curve $y = 1 + x^{3/2}$.

Answer: $\sqrt{44}$

2. Find the largest possible perimeter of the rectangle that can be inscribed in a semicircle of radius R if one side of the rectangle lies along the diameter of the semicircle.

3. Among all rectangles of perimeter P , show that the square has the greatest area.


4. Among all rectangles of given area A , show that the square has the least perimeter.

5. Find the equation of the straight line of maximum slope tangent to the curve $y = 1 + 2x - x^3$.

Answer: $y = 1 + 2x$

4.7 Linear Approximation

The best line approximating the graph of $y = f(x)$ near $(a, f(a))$ is the tangent line through $(a, f(a))$.



The linearization of the function f about a is the function L defined by

$$L(x) = f(a) + f'(a)(x - a)$$

We say that L approximates f near $x = a$ and write $f(x) \approx L(x)$.

Example 4.7.1. Using the linearization, approximate $\sqrt{26}$. (Hint: use the linearization of \sqrt{x} at $x = 25$.)

Solution. $f'(x) = \frac{1}{2\sqrt{x}}$. $f'(25) = \frac{1}{10}$. So

$$L(x) = 5 + \frac{1}{10}(x - 25).$$

Hence $f(26) \approx L(26) = 5.1$.

Example 4.7.2. Approximate $\cos \pi/5 = \cos 36^\circ$ using the linearization of $\cos x$ at $x = \pi/6$.

Solution. $L(x) = \cos \frac{\pi}{6} - \sin \frac{\pi}{6} (x - \frac{\pi}{6}) = \frac{\sqrt{3}}{2} - \frac{1}{2} (x - \frac{\pi}{6})$.

$$\cos 36^\circ \approx L(\pi/5) = \frac{\sqrt{3}}{2} - \frac{1}{2} \frac{\pi}{30} \approx 0.81367$$

Error Estimation

The error in the linear approximation is

$$\frac{f''(s)}{2} (x - a)^2$$

where s is some number between a and x . (The proof depends on the generalized mean value theorem.)

Since we do not know s , we have to choose $f''(s)$ to be largest (in absolute value) possible value, to get the maximum error.

So for the previous example, $f''(x) = -\sin x$, $a = \pi/6$, $x = \pi/5$, $\pi/6 < s < \pi/5$. Note that $f''(s) \leq 1$. So the error is smaller that $\frac{1}{2} (x - a)^2 = \frac{\pi^2}{1800} < 0.00549$. So $0.81367 - 0.00549 < \cos \pi/5 < 0.81367 + 0.00549$.

Exercises.

Answer: $7 + \frac{1}{14}$.

1. Sketch $y = \frac{1}{\sqrt{x}}$ and its linearization about $x = 4$.

3. Approximate $\sin 46^\circ$ using linearization.

2. Approximate $\sqrt{50}$ using linearization.

Answer: $\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2} \frac{\pi}{180}$. (Note: $f(x) = \sin x^\circ = \sin \frac{\pi x}{180}$)

4.8 Exam 2 Review

Section	Exercises
3.1	27-29
3.3	11-16, 19-48, 55-66
3.5	1-12, 19-32
4.1	1-15
4.3	1-24
4.4	1-17, 18-39
4.5	1-22
4.6	7-39
4.7	1-32
4.9	1-10, 15-22

Table 4.1: Exam 2 Review Problems from Adams & Essex Calculus: A Complete Course 7th Edition

Sample Exam 2

1. Find the dimensions of the right triangle with hypotenuse $h = 2$ and maximum area.

2. Find the min and max values of $f(x) = 2x^3 - 15x^2 + 24x + 19$ on the interval $[0, 5]$
3. Let $f(x) = x^4 - 3x^2 + 2$.
- Find the intervals on which f is increasing or decreasing. Find all local extrema for f .
 - Find the intervals on which f is concave up or down. Find all inflection points for f .
 - Sketch a graph of $f(x)$ using parts (a) and (b).
4. Find all the horizontal and vertical asymptotes for

$$y = \frac{x^2 + 3}{1 - 3x^2}.$$

5. Let $f(x) = 2x^2 + x^3$, $x > 0$. Show that f is invertible and find $(f^{-1})'(16)$.
6. Find dy/dx by implicit differentiation if

$$y^2 e^x + y \ln x = 2.$$

7. Let $y = (1/x)^{\ln x}$. Find dy/dx .
8. Find $\cos(\sin^{-1} 0.7)$.

Chapter 5

Integration

5.1 The Definite Integral

Our main goal in this section is to find the area between the graph of a function and the x-axis.

Idea is to approximate this region with rectangles.

Let's start with an easy example.

Example 5.1.1. Find the area of the region lying under the straight line $y = x + 1$, above the x-axis and between the lines $x = 0$ and $x = 2$.

Solution. Two ways of approximating the area. With “smaller” rectangles and “larger” rectangles.

With “smaller” rectangles. Divide the interval $[0, 2]$ into n equal pieces, call $x_0 = 0$, $x_1 = 2/n$, $x_2 = 4/n$, ..., $x_n = 2$.

$$L_n = f(x_0)(x_1 - x_0) + f(x_1)(x_2 - x_1) + \cdots + f(x_{n-1})(x_n - x_{n-1})$$

$$L_1 = f(0)(2 - 0) = 2,$$

$$L_2 = f(0)(1 - 0) + f(1)(2 - 1) = 1 + 2 = 3,$$

$$L_4 = (f(0) + f(1/2) + f(1) + f(3/2))(1/2) = 7/2,$$

$$L_n = \frac{2(2n - 1)}{n}$$

Thus

$$\lim_{n \rightarrow \infty} L_n = 4$$

which is the area.

We can repeat this with “larger” rectangles. In this case

$$U_n = \frac{2(2n + 1)}{n}$$

And again, $\lim_{n \rightarrow \infty} U_n = 4$.

This procedure can be used to find the areas under more exotic curves.

Definition 5.1.1. Suppose $f : [a, b] \rightarrow \mathbb{R}$. If $a = x_0 < x_1 < \cdots < x_n = b$ then the set $P = \{x_0, x_1, \dots, x_n\}$ is called a **partition** of the interval $[a, b]$. Let $f(l_i)$ be the smallest value and $f(u_i)$ be the largest value of $f(x)$ on $[x_i, x_{i+1}]$. Then we define the **lower Riemann sum**

$$L(f, P) = f(l_0)(x_1 - x_0) + f(l_1)(x_2 - x_1) + \cdots + f(l_{n-1})(x_n - x_{n-1})$$

and the **upper Riemann sum**

$$U(f, P) = f(u_0)(x_1 - x_0) + f(u_1)(x_2 - x_1) + \cdots + f(u_{n-1})(x_n - x_{n-1})$$

Suppose that there exists exactly one number I such that for every partition P of $[a, b]$,

$$L(f, P) \leq I \leq U(f, P)$$

Then we say f is **integrable** on $[a, b]$ and we call I , the definite integral of f on $[a, b]$ and write

$$I = \int_a^b f(x) dx.$$

Definition 5.1.2. Let R be the region bounded by the graph of $f(x)$, the x -axis and the lines $x = a$ and $x = b$. If $f(x) \geq 0$ on $[a, b]$ then we define

$$\text{Area}(R) = \int_a^b f(x) dx$$

If $f(x) \leq 0$ on $[a, b]$ then

$$\text{Area}(R) = - \int_a^b f(x) dx$$

In general $\int_a^b f(x)$ is the area of the part of R lying above the x -axis minus the area of the part below the x -axis.




Figure 5.1: $\int_a^b f(x) dx = A_1 - A_2$

Here the variable x is a dummy variable. Replacing x by any other symbol does not change value of the integral. The function f is known as **integrand**. dx is differential of x and if an integrand depends on more than one variable it tells which one is the variable of integration.

In the first example, we showed that

$$\int_0^2 (x+1) dx = 4.$$

Which functions are integrable?

Theorem 5.1.1. *If f is continuous on $[a, b]$ then f is integrable.*

Piecewise continuous functions are also integrable.

Example 5.1.2. *Not every function is integrable. Define $f : [0, 1] \rightarrow \mathbb{R}$ such that $f(x) = 0$ if x is a rational number and $f(x) = 1$ if x is an irrational number. Let P be a partition of $[0, 1]$. Then the smallest value $f(l_i)$ in the subinterval $[x_i, x_{i+1}]$ is 0 since every such interval contains a rational, while the largest value $f(u_i)$ is 1 since every such interval also contains an irrational. Thus $L(f, P) = 0$ and $U(f, P) = 1$. This is true for any partition P . Hence there are infinitely many numbers between $L(f, P)$ and $U(f, P)$ for any partition P . Hence f is not integrable on $[0, 1]$.*

Properties of Definite Integral

The following properties are easy consequences of the definition of definite integral.

Theorem 5.1.2. *Let f and g be integrable on an interval containing the points a , b and c .*

1. $\int_a^a f(x) dx = 0$.
2. We can define $\int_a^b f(x) dx$ when $a > b$. In this case the partition points are $x_0 = a > x_1 > \cdots > x_n = b$. Hence $\int_a^b f(x) dx = -\int_b^a f(x) dx$.
3. If c is a constant $\int_a^b c f(x) dx = c \int_a^b f(x) dx$.
4. $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$
5. $\int_a^b f(x) dx + \int_b^c f(x) dx = \int_a^c f(x) dx$.
6. If $a \leq b$ and $f(x) \leq g(x)$ then $\int_a^b f(x) dx \leq \int_a^b g(x) dx$.
7. If f is an odd function then $\int_{-a}^a f(x) dx = 0$.

Example 5.1.3. $\int_{-3}^3 (\sin x^3)^5 dx = 0$ since $f(x) = (\sin x^3)^5$ is odd. Verify.

8. If f is an even function then $\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$.

Example 5.1.4. Show that $\int_a^b c dx = c(b - a)$ and $\int_a^b x dx = \frac{(b^2 - a^2)}{2}$ interpreting the integrals as areas.

Example 5.1.5. Using the properties of the integral, compute

$$\int_{-2}^2 (3 + 5x) dx$$

Example 5.1.6. Compute $\int_{-3}^3 \sqrt{9 - x^2} dx$.

Solution. This is the area of the semicircle with radius 3 and center $(0, 0)$. The answer is $\frac{9\pi}{2}$.

5.2 The Fundamental Theorem of Calculus

In this section we develop the relation between the integral and the derivative.

Antiderivative

We will call $F(x)$ as an antiderivative of $f(x)$ if $F'(x) = f(x)$. For example x is an antiderivative of 1. Note that $x + 1$ is also an antiderivative of 1. So antiderivatives are not unique.

If F and G are antiderivatives of f on an interval, so that $F'(x) = G'(x) = f(x)$ then

$$\frac{d}{dx}(F(x) - G(x)) = 0.$$

But Theorem 2.7.3 tells that $F(x) - G(x)$ must be a constant. Hence if $F(x)$ is an antiderivative of $f(x)$ then for any C , $F(x) + C$ is also an antiderivative of $f(x)$. Also any antiderivative of $f(x)$ is of the form $F(x) + C$ for some c .

Definition 5.2.1. The *indefinite integral* of $f(x)$ on interval I is

$$\int f(x) dx = F(x) + C$$

provided $F'(x) = f(x)$ on I .

The Fundamental Theorem of Calculus

Theorem 5.2.1.

PART I. Suppose f is continuous.

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

PART II. Suppose f is differentiable.

$$\int_a^b f'(x) dx = f(b) - f(a)$$

Intuitively, the second part of fundamental theorem of calculus states that the total change (right hand side) is the sum of all the little changes (right hand side). Recall that $f'(x)dx$ is a tiny change in the value of f . If you add up (integrate) all these tiny changes, you get the total change $f(b) - f(a)$.

Proof. For the first part, let

$$F(x) = \int_a^x f(t) dt.$$

Then

$$F'(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{1}{h} \left(\int_a^{x+h} f(t) dt - \int_a^x f(t) dt \right) = \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) dt$$

Let $m(h)$ be the minimum, $M(h)$ be the maximum of f on $[x, x+h]$. Then $m(h) \leq f(t) \leq M(h)$ on $x \leq t \leq x+h$. Thus

$$m(h)h = \int_x^{x+h} m(h) dx \leq \int_x^{x+h} f(t) dt \leq \int_x^{x+h} M(h) dx = M(h)h.$$

Or

$$m(h) \leq \frac{1}{h} \int_x^{x+h} f(t) dt \leq M(h)$$

Since $\lim_{h \rightarrow 0} m(h) = \lim_{h \rightarrow 0} M(h) = f(x)$, by Sandwich Theorem,

$$F'(x) = \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) dt = f(x).$$

Proof of the second part. Let

$$F(x) = \int_a^x f'(t) dt.$$

Then by part I, $F'(x) = f'(x)$. We have seen that the only function whose derivative is zero on an interval is the constant function. Thus $F'(x) - f'(x) = 0$. Hence $F(x) - f(x) = c$, a constant. Since $0 = F(a)$, $c = -f(a)$. And

$$\int_a^b f'(t) dt = F(b) = f(b) + c = f(b) - f(a).$$

□

Second part gives a method to evaluate definite integrals. To compute $\int_a^b f(x) dx$, find a function $F(x)$ whose derivative is $f(x)$. Then the value of $\int_a^b f(x) dx = F(b) - F(a)$.

We will use the evaluation symbol

$$F(x) \Big|_a^b = F(b) - F(a).$$

Example 5.2.1. Evaluate

$$1. \int_0^a x^2 dx = \frac{a^3}{3}$$

$$2. \int_{-1}^2 (x^2 - 3x + 2) dx = \frac{9}{2}$$

Example 5.2.2. Find the derivatives of the following functions.

$$1. F(x) = \int_x^3 e^{-t^2} dt$$

$$2. G(x) = \int_{-4}^{5x} e^{-t^2} dt$$

$$3. H(x) = \int_{x^2}^{x^3} e^{-t^2} dt$$

Solution. By the Fundamental Theorem of Calculus Part I,

$$F(x) = - \int_3^x e^{-t^2} dt \implies F'(x) = -e^{-x^2}.$$

Let $g(x) = \int_{-4}^x e^{-t^2} dt$. Then $G(x) = g(5x)$ and

$$G'(x) = g'(5x)5 = 5e^{-(5x)^2}$$

$H(x) = \int_{x^2}^a e^{-t^2} dt + \int_a^{x^3} e^{-t^2} dt$. Then

$$H'(x) = e^{-x^6} 3x^2 - e^{-x^4} 2x.$$

In general

$$\frac{d}{dx} \int_{f(x)}^{g(x)} h(t) dt = h(g(x))g'(x) - h(f(x))f'(x).$$

Exercises.

1. $\frac{d}{dx} \int_2^x \frac{\sin t}{t} dt$

Answer: $\frac{\sin x}{x}$

2. $\frac{d}{dx} \int_{\sin x}^{\cos x} \frac{1}{1-x^2} dx$

Answer: $-\frac{1}{\sin x} - \frac{1}{\cos x}$

3. $\frac{d}{dx} F(\sqrt{x})$, if $F(t) = \int_0^t \cos(x^2) dx$

Answer: $\frac{1}{2\sqrt{x}} \cos(x)$

4. Find $H'(2)$ if $H(x) = 3x \int_4^{x^2} e^{\sqrt{t}} dt$

Answer: $6e^2$

5.3 The Method of Substitution

The following should be memorized.

- $\int x^n dx = \frac{1}{n+1} x^{n+1} + C$, if $n \neq -1$

- $\int 1 dx = x + C$

- $\int x dx = \frac{1}{2} x^2 + C$

- $\int x^2 dx = \frac{1}{3} x^3 + C$

- $\int \sqrt{x} dx = \frac{2}{3} x^{3/2} + C$

- $\int \frac{1}{x} dx = \ln|x| + C$

- $\int \sin x dx = -\cos x + C$

- $\int \cos x dx = \sin x + C$

- $\int \sec^2 x dx = \tan x + C$

- $\int \csc^2 x dx = -\cot x + C$

- $\int \sec x \tan x dx = \sec x + C$

- $\int \csc x \cot x dx = -\csc x + C$

- $\int \frac{1}{\sqrt{1-x^2}} dx = \arcsin x + C$

- $\int \frac{1}{1+x^2} dx = \arctan x + C$

- $\int e^x dx = e^x + C$

- $\int a^x dx = \frac{1}{\ln a} a^x + C$

Example 5.3.1.

1. $\int (x^3 - 3x^2 + 6x - 9) dx = \frac{x^4}{4} - x^3 + 3x^2 - 9x + C$

2. $\int (5x^{3/4} - \frac{1}{\sqrt{x}}) dx$

$$3. \int \frac{(x+1)^3}{x} dx$$

The Chain Rule says

$$\frac{d}{dx} f(g(x)) = f'(g(x)) g'(x).$$

So we have,

$$\int f'(g(x)) g'(x) dx = f(g(x)) + C$$

To see this another way, let $u = g(x)$. Then $du/dx = g'(x)$. In differential form $du = g'(x) dx$

$$\int f'(g(x)) g'(x) dx = \int f'(u) du = f(u) + C = f(g(x)) + C$$

Example 5.3.2. Compute the following integrals.

$$1. I = \int x \sin(2x^2) dx.$$

Let $2x^2 = u$ then $4x dx = du$.

$$I = \frac{1}{4} \int \sin u du = -\frac{\cos u}{4} + C = -\frac{\cos 2x^2}{4} + C$$

$$2. I = \int \sec^2(3x+2) dx$$

Let $3x+2 = u$ then $3dx = du$.

$$I = \int \sec^2 u \frac{du}{3} = \frac{\tan u}{3} + C = \frac{1}{3} \tan(3x+2) + C$$

$$3. I = \int \frac{x}{(x-4)^3} dx$$

Let $x-4 = u$.

$$I = \int \frac{u+4}{u^3} du = \int (u^{-2} + 4u^{-3}) du = -u^{-1} - 2u^{-2} = \frac{-1}{x-4} - \frac{2}{(x-4)^2} + C$$

$$4. I = \int \tan^2 \theta \sec^2 \theta d\theta.$$

Let $\tan \theta = u$. Then $\sec^2 \theta d\theta = du$.

$$I = \int u^2 du = \frac{u^3}{3} + C = \frac{\tan^3 \theta}{3} + C$$

$$5. I = \int \sqrt{\frac{x^4}{x^3-1}} dx = \int \frac{x^2}{\sqrt{x^3-1}} dx.$$

Let $x^3-1 = u$. Then $x^2 dx = \frac{du}{3}$.

$$I = \int \frac{du/3}{\sqrt{u}} = \frac{1}{3} \frac{u^{1/2}}{1/2} + C = \frac{2}{3} \sqrt{x^3-1} + C$$

6. Let $y = x \int_2^{x^2} \sin(t^3) dt$. Find y'

$$y' = \int_2^{x^2} \sin(t^3) dt + x \sin(x^6) 2x$$

7. $I = \int \sec x dx$.

There is an interesting trick to evaluate this integral!

$$I = \int \sec x \frac{(\sec x + \tan x)}{\sec x + \tan x} dx = \int \frac{\sec^2 x + \sec x \tan x}{\sec x + \tan x} dx$$

Let $u = \sec x + \tan x$, then

$$I = \int \frac{du}{u} = \ln |u| + C = \ln |\sec x + \tan x| + C.$$

Exercises.

1. $\int x e^{x^2} dx$

Answer: $\frac{1}{2} e^{x^2} + C$

2. $\int_0^{2\pi} \sin^2 x \cos^2 x dx$

Answer: $\frac{\pi}{4}$

3. $\int_e^{e^2} \frac{dt}{t \ln t}$,

Answer: $\ln 2$

4. $\int \frac{dx}{e^x + 1}$

Answer: $x - \ln(1 + e^x) + C$.

5. $\int \frac{x^2}{2 + x^6}$

Answer: $\frac{1}{3\sqrt{2}} \arctan(x^3/\sqrt{2}) + C$

6. $\int \sec^5 x \tan x dx$

Answer: $\frac{1}{5} \sec^5 x + C$


5.4 Areas of Plane Regions

Example 5.4.1. Find the area of the region lying above the x -axis and under the curve $y = 3x - x^2$.

Solution. The points where the graph intersects the x -axis are $y = 0$ which gives $x = 0, x =$.

The area is

$$\int_0^3 (3x - x^2) dx = \left(\frac{3}{2}x^2 - \frac{1}{3}x^3 \right) \Big|_0^3 = \frac{9}{2}.$$




Example 5.4.2. Find the area of the region lying above the line $y = 1$ and below the curve $y = \frac{5}{x^2+1}$.

Solution. The curves $y = 1$ and $y = \frac{5}{x^2+1}$ intersect at $x = \pm 2$. The area

The area is

$$\int_{-2}^2 \frac{5}{x^2+1} dx - \int_{-2}^2 1 dx = 10 \arctan 2 - 4.$$



Suppose $f(x) \leq g(x)$ for $a \leq x \leq b$. Then the area of the region between these two curves and the lines $x = a$ and $x = b$ is

$$\text{Area} = \int_a^b (g(x) - f(x)) dx.$$


Example 5.4.3. Find the area of the bounded region lying between the curves $y = x^2 - 2x$ and $y = 4 - x^2$.

Solution. The two curves intersect at

$$x^2 - 2x = 4 - x^2 \implies 2x^2 - 2x - 4 = 0 \implies (x-2)(x+1) = 0$$

So the intersection points are $x = -1$ and $x = 2$. The area of the region is

$$\text{Area} = \int_{-1}^2 (4 - x^2) - (x^2 - 2x) dx = 9$$




Example 5.4.4. Find the area of the region bounded by $y = \sqrt{x}$ and $y = x^2$.

Solution. The curves intersect at

$$\sqrt{x} = x^2 \implies x = x^4 \implies x(1 - x^3) = 0.$$

Hence the intersection points are $x = 0$ and $x = 1$.

$$\text{Area} = \int_0^1 (\sqrt{x} - x^2) dx = \frac{2}{3} - \frac{1}{3} = \frac{1}{3}.$$



Example 5.4.5. Find the area of the region lying to the right of the parabola $x = y^2 - 12$ and to the left of the straight line $y = x$.


Solution. The curves intersect at

$$y^2 - 12 = y \implies (y - 4)(y + 3) = 0$$

The intersection points are $y = 4$ and $y = -3$.

$$\text{Area} = \int_{-3}^4 (y - (y^2 - 12)) dy = \frac{343}{6}$$

An alternative way is to make the transformation $x \rightarrow y$, $y \rightarrow x$. The problem becomes finding the area between $y = x^2 - 12$ and $y = x$.



Exercises.

- Find the area bounded by the curves $y = x^2 - 2x$, and $y = 3x - x^2$.

Answer: $\frac{125}{24}$

- Find the area bounded by the curves $y = x^3$, and $y = x$ in the region $x \geq 0$.

Answer: $\frac{1}{4}$.

- Find the area bounded by $y = \sin x$ and $y = \cos x$ between two consecutive intersections of these curves.

Answer: $2\sqrt{2}$

- Find the area bounded by the curves $y = \frac{1}{x}$ and $2x + 2y = 5$.

Answer: $\frac{15}{8} - \ln 4$

- Find the area bounded by the curves $x - y = 7$ and $x = 2y^2 - y + 3$.

Answer: 9

5.5 Integration by Parts

Integrating both sides of

$$\frac{d}{dx}(u(x)v(x)) = \frac{du}{dx}v + u\frac{dv}{dx}$$

we get

$$\int \frac{d}{dx}(u(x)v(x))dx = \int \frac{du}{dx}vdx + \int u\frac{dv}{dx}dx$$

Since $\int \frac{d}{dx}(u(x)v(x))dx = u(x)v(x)$, in differential notation, we get

$$uv = \int duv + \int u dv$$

Another way to write this is

$$\int u dv = uv - \int v du$$

This is one of the most powerful method to integrate, known as the **integration by parts**.

Example 5.5.1. $\int xe^x dx$.

Solution. Let $u = x$ and $dv = e^x dx$. Then $du = dx$ and $v = e^x$.

$$\int xe^x dx = xe^x - \int e^x dx = xe^x - e^x + C$$

Example 5.5.2. $\int \ln x dx$.

Solution. Let $u = \ln x$ and $dv = dx$. Then $du = dx/x$ and $v = x$.

$$\int \ln x dx = x \ln x - \int x \frac{dx}{x} = x \ln x - x + C$$

Example 5.5.3. $I = \int x^2 \sin x dx$

Solution. We have to integrate by parts twice. Let $u = x^2$ and $dv = \sin x dx$. Then $du = 2x dx$ and $v = -\cos x$.

$$I = x^2(-\cos x) - \int (-\cos x)2x dx = -x^2 \cos x + \int 2x \cos x dx$$

Now let $u = 2x$ and $dv = \cos x dx$. Then $du = 2 dx$ and $v = \sin x$. And

$$\int 2x \cos x dx = 2x \sin x - \int 2 \sin x dx$$

Hence

$$I = -x^2 \cos x + 2x \sin x + 2 \cos x + C$$

Example 5.5.4. $I = \int x \arctan x dx$

Solution. Let $u = \arctan x$, $dv = x dx$. Then $du = dx/(1+x^2)$ and $v = x^2/2$.

$$I = \frac{1}{2}x^2 \arctan x - \frac{1}{2} \int \frac{x^2}{1+x^2} dx = \frac{1}{2}x^2 \arctan x - \frac{1}{2} \int \left(1 - \frac{1}{1+x^2}\right) dx$$

And

$$I = \frac{1}{2}x^2 \arctan x - \frac{1}{2}(x - \arctan x) + C$$

Example 5.5.5. Find $I = \int e^x \sin x dx$.

Solution. There is a circular argument here. We will integrate by parts twice to return the same integral. Let $u = \sin x$ and $dv = e^x dx$. Then $du = \cos x dx$, $v = e^x$.

$$\int e^x \sin x dx = e^x \sin x - \int \cos x e^x dx$$

Now let $u = \cos x$ and $dv = e^x dx$.

$$\int \cos x e^x dx = \cos x e^x - \int (-\sin x) e^x = \cos x e^x + I$$

So

$$I = e^x \sin x - e^x \cos x - I$$

Hence

$$2I = e^x(\sin x - \cos x) + C \implies I = \frac{e^x}{2}(\sin x - \cos x) + C.$$

Example 5.5.6. $I = \int \sec^3 x dx$.

Solution. Let $u = \sec x$ and $dv = \sec^2 x dx$. Then $du = \sec x \tan x dx$ and $v = \tan x$

$$I = \sec x \tan x - \int \sec x \tan^2 x dx$$

Using $\tan^2 x = \sec^2 x - 1$,

$$I = \sec x \tan x + \int \sec x dx - I$$

Using $\int \sec x dx = \ln|\sec x + \tan x|$, (see the section on “The Method of Substitution”) we get

$$I = \frac{1}{2} \sec x \tan x + \frac{1}{2} \ln|\sec x + \tan x| + C$$

Exercises.

1. $\int x \cos x dx,$

Answer: $\cos x + x \sin x + C$

2. $\int x e^{\sqrt{x}} dx$

Hint: use $x = u^2$. Answer: $2e^{\sqrt{x}}(-6 + 6\sqrt{x} - 3x + x^{3/2}) + C.$

3. $\int x^2 \arctan(x) dx$

Answer: $\frac{1}{6}(-x^2 + 2x^3 \arctan(x) + \ln(1+x^2)) + C$

5. $\int \arctan x dx$

4. $\int_1^e \sin(\ln x) dx$

Answer: $\frac{1}{2}(1 - e \cos 1 + e \sin 1)$

Hint: use $\arctan x = u$, $dx = dv$. Answer: $x \arctan x - \frac{1}{2} \ln(1+x^2) + c$

5.6 Integrals of Rational Function

In this section we are concerned with integrals of the form

$$\int \frac{P(x)}{Q(x)} dx$$

where $P(x)$ and $Q(x)$ are both polynomials.

We will look at methods to deal with such integrals when $\deg(P(x)) < \deg(Q(x))$.

The case $\deg(Q(x)) = 1$ and $\deg(P(x)) = 0$

Example 5.6.1. $\int \frac{1}{ax+b} dx = \frac{1}{a} \ln ax+b + C.$

Solution. Let $u = ax+b$ then $du = adx$ and the integral becomes $\frac{1}{a} \int \frac{du}{u}$.

The case $\deg(Q(x)) = 2$ and $\deg(P(x)) = 0$

First let's look at two examples where $Q(x)$ does not have real roots.

Example 5.6.2. $\int \frac{dx}{x^2+a^2} = \frac{1}{a} \tan^{-1} \frac{x}{a} + C.$

Solution. Let $x = a \tan \theta$ (We will talk about these types of transformations in the next section in detail!), then $dx = a \sec^2 \theta d\theta$ and $x^2 + a^2 = a^2(\sec^2 \theta + 1) = a^2 \tan^2 \theta$.

If $Q(x) = ax^2 + bx + c$ has no real roots, we have to **complete to squares**.

Example 5.6.3. $\int \frac{dx}{x^2+3x+3}$

Solution. Notice that x^2+3x+3 has no real roots. So we complete to squares

$$x^2+3x+3 = (x+\frac{3}{2})^2 + \frac{3}{4}$$

Letting $u = (x+3/2)$ and $du = dx$,

$$\int \frac{dx}{(x+\frac{3}{2})^2 + \frac{3}{4}} = \int \frac{du}{u^2 + \frac{3}{4}} = \frac{2}{\sqrt{3}} \tan^{-1} \frac{2(x+\frac{3}{2})}{\sqrt{3}} + C$$

The last part follows from the last example.

If $Q(x)$ has real roots then we use **partial fractions**.

Partial Fractions

Let us still assume that $\deg(P(x)) < \deg(Q(x))$. The Fundamental Theorem of Algebra tells that every polynomial can be factored (over the real numbers) into a product of real linear factors $(x - a_i)$ and real quadratic factors $x^2 + b_i x + c_i$ having no real roots.

$$Q(x) = (x - a_1)^{m_1} (x - a_2)^{m_2} \cdots (x - a_j)^{m_j} (x^2 + b_1 x + c_1)^{n_1} \cdots (x^2 + b_k x + c_k)^{n_k}$$

To each factor of the form $(x - a)^m$, the partial fraction decomposition contains a sum

$$\frac{A_1}{(x - a)} + \frac{A_2}{(x - a)^2} + \cdots + \frac{A_m}{(x - a)^m}$$

To each factor of the form $(x^2 + bx + c)^n$, the partial fraction decomposition contains a sum

$$\frac{B_1 x + C_1}{(x^2 + bx + c)} + \frac{B_2 x + C_2}{(x^2 + bx + c)^2} + \cdots + \frac{B_n x + C_n}{(x^2 + bx + c)^n}$$

Example 5.6.4. $\int \frac{(x+4)}{x^2-5x+6} dx$

Solution.

$$\begin{aligned} \frac{x+4}{x^2-5x+6} &= \frac{A}{x-2} + \frac{B}{x-3} \\ x+4 &= A(x-3) + B(x-2) \end{aligned}$$

Plugging $x = 2$ gives $A = -6$ and plugging $x = 3$ gives $B = 7$. So

$$\int \frac{(x+4)}{x^2-5x+6} dx = -6 \int \frac{dx}{x-2} + 7 \int \frac{dx}{x-3} = -6 \ln(x-2) + 7 \ln(x-3) + C$$

Example 5.6.5. $\int \frac{2+3x+x^2}{x(x^2+1)} dx.$

Solution. The partial fraction decomposition is

$$\frac{2+3x+x^2}{x(x^2+1)} = \frac{A}{x} + \frac{Bx+C}{x^2+1} \implies A(x^2+1) + x(Bx+C) = 2+3x+x^2$$

Since this equation holds for every x , we have $A + B = 1$ (coefficient of x^2 term), $C = 3$ (coefficient of x term), $A = 2$ (coefficient of constant term) and $B = -1$.

$$\int \frac{2+3x+x^2}{x(x^2+1)} dx = \int \frac{2}{x} dx + \int \frac{-x+3}{x^2+1} dx = 2 \ln x - \frac{1}{2} \ln(x^2+1) + 3 \tan^{-1} x + C.$$

Example 5.6.6. Evaluate $\int \frac{1}{x(x-1)^2} dx.$

Solution.

$$\begin{aligned} \frac{1}{x(x-1)^2} &= \frac{A}{x} + \frac{B}{(x-1)} + \frac{C}{(x-1)^2} \\ 1 &= A(x-1)^2 + Bx(x-1) + Cx \end{aligned}$$

Letting $x = 0$ gives $A = 1$, $x = 1$ gives $C = 1$. The coefficient of x^2 is $A + B$ which must be zero. So $B = -1$.

$$\int \frac{1}{x(x-1)^2} dx = \int \frac{1}{x} dx - \int \frac{1}{x-1} dx + \int \frac{1}{(x-1)^2} dx = \ln|x| - \ln|x-1| - \frac{1}{x-1} + C.$$

The last integral can be found by letting $u = x - 1$.

The Case $\deg(P(x)) \geq \deg(Q(x))$

If $\deg(P(x)) \geq \deg(Q(x))$ then we divide $P(x)$ to $Q(x)$ and get a rational function with the degree of numerator less than the degree of denominator.

Example 5.6.7. Evaluate $\int \frac{x^3 + 3x^2}{x^2 + 1} dx$

Solution.

$$\begin{aligned}\frac{x^3 + 3x^2}{x^2 + 1} &= x + 3 - \frac{x + 3}{x^2 + 1} \\ \int \frac{x^3 + 3x^2}{x^2 + 1} dx &= \int \frac{x}{x^2 + 1} dx + \int \frac{3}{x^2 + 1} dx \\ \frac{x^3 + 3x^2}{x^2 + 1} &= \frac{x^2}{2} + 3x - \frac{1}{2} \ln(x^2 + 1) - 3 \tan^{-1} x + C\end{aligned}$$

Exercises.

Answer: $\frac{1}{6}(-\ln|1-x| + 2\ln|2-x| + \ln|1+x| - 2\ln|2+x|) + C$

1. $\int \frac{2dx}{5-4x} dx,$

Answer: $-\frac{1}{2} \ln|5-4x| + C$

5. $\int \frac{dx}{x^4 - 4x^3}$

Answer: $\frac{1}{8x^2} + \frac{1}{16x} + \frac{1}{64} \ln|4-x| - \frac{1}{64} \ln|x| + C$

2. $\int \frac{2x+2}{x^2+4} dx$

Answer: $\arctan(x/2) + \ln|4+x^2| + C$

6. $\int \frac{1}{x^3 + 9x} dx$

Answer: $\frac{1}{9} \ln|x| - \frac{1}{18} \ln|9+x^2| + C$

3. $\int \frac{x^2}{x^2 + x - 2} dx$

Answer: $x + \frac{1}{3} \ln|1-x| - \frac{4}{3} \ln|2+x| + C.$

7. $\int \frac{1}{e^{2x} - 4e^x + 4} dx$

Answer: $\frac{1}{4} \left(-\frac{2}{-2+e^x} + x - \ln|2-e^x| \right) + C$

5.7 Inverse Substitutions**The Inverse Sine Substitution**

If an integral involves $\sqrt{a^2 - x^2}$, try the substitution $x = a \sin \theta$ or $\theta = \sin^{-1} \frac{x}{a}$.

We can assume $a > 0$. Notice that $\sqrt{a^2 - x^2}$ makes sense only when $-a \leq x \leq a$ which corresponds to $-\pi/2 \leq \theta \leq \pi/2$ so that $\cos \theta \geq 0$. Hence

$$\sqrt{a^2 - x^2} = \sqrt{a^2(1 - \sin^2 \theta)} = a\sqrt{\cos^2 \theta} = a|\cos \theta| = a \cos \theta.$$

Example 5.7.1. Evaluate $I = \int \frac{dx}{(5-x^2)^{3/2}}$.

Solution. Let $x = \sqrt{5} \sin \theta$, $dx = \sqrt{5} \cos \theta d\theta$.

$$(5-x^2)^{3/2} = (5-5\sin^2 \theta)^{3/2} = 5^{3/2} |\cos \theta|^3 = 5^{3/2} \cos^3 \theta.$$

since $\cos \theta \geq 0$. So

$$I = \int \frac{\sqrt{5} \cos \theta d\theta}{5^{3/2} \cos^3 \theta} = \frac{1}{5} \int \sec^2 \theta d\theta = \frac{1}{5} \tan \theta + C = \frac{1}{5} \frac{x}{\sqrt{5-x^2}} + C$$

The last equality can be found using $\sin \theta = \frac{x}{\sqrt{5}}$.

Example 5.7.2.

$$\begin{aligned} \int \frac{dx}{\sqrt{8x-x^2}} &= \int \frac{dx}{\sqrt{16-(x-4)^2}} \\ &= \int \frac{du}{\sqrt{a^2-u^2}} \\ &= \sin^{-1} \left(\frac{u}{a} \right) + C \\ &= \sin^{-1} \left(\frac{x-4}{4} \right) + C \end{aligned}$$

The inverse Tangent Substitution

If an integral involves $\sqrt{a^2+x^2}$ or $\frac{1}{x^2+a^2}$, try the substitution $x = a \tan \theta$ or $\theta = \tan^{-1} \frac{x}{a}$.

Since x can take any real value, we have $-\pi/2 < \theta < \pi/2$ so that $\sec \theta > 0$. Assuming $a > 0$,

$$\sqrt{a^2+x^2} = \sqrt{a^2(1+\tan^2 \theta)} = a\sqrt{\sec^2 \theta} = a|\sec \theta| = a \sec \theta.$$

Example 5.7.3. Evaluate $I = \int \frac{dx}{\sqrt{4+x^2}}$.

Solution. Let $x = 2 \tan \theta$, $dx = 2 \sec^2 \theta d\theta$.

$$\sqrt{4+x^2} = 2\sqrt{\sec^2 \theta} = 2|\sec \theta| = 2 \sec \theta$$

since $\sec \theta > 0$. Using $\tan \theta = x/2$ we can find $\sec \theta = \frac{\sqrt{4+x^2}}{2}$ and

$$I = \int \sec \theta d\theta = \ln |\sec \theta + \tan \theta| + C = \ln \left| \frac{\sqrt{4+x^2}}{2} + \frac{x}{2} \right| + C$$

You are not responsible for the inverse secant transformation which can be used to solve integrals involving $\sqrt{x^2-a^2}$.

Exercises.

$$1. \int \frac{x^2}{\sqrt{1-4x^2}} dx,$$

Answer: $-\frac{1}{8}x\sqrt{1-4x^2} + \frac{1}{16}\arcsin(2x) + C$

2. $\int \frac{1}{x\sqrt{9-x^2}} dx$

Answer: $\frac{1}{3}\ln|x| - \frac{1}{3}\left|3 + \sqrt{9-x^2}\right| + C.$

3. $\int \frac{1}{x^2+2x+10} dx$

Answer: $\frac{1}{3}\arctan\left(\frac{1+x}{3}\right) + C$

5.8 Indefinite Integrals Exercises

Example 5.8.1. $\int \frac{x}{x^2+9} dx = \frac{1}{2} \int \frac{2x}{x^2+9} dx = \frac{1}{2} \ln(x^2+9) + c$

Example 5.8.2. $\int (e^{-x} - e^{-4x}) dx = -e^{-x} + \frac{1}{4}e^{-4x} + c$

Example 5.8.3. $\int \frac{3-\cos x}{3x-\sin x} dx = \ln|3x-\sin x| + c$

Example 5.8.4.

$$\begin{aligned} \int \left(3x^4 + \frac{1}{x} + \sqrt[3]{x^2}\right) dx &= \int 3x^4 dx + \int \frac{1}{x} dx + \int \sqrt[3]{x^2} dx \\ &= 3\frac{x^5}{5} + \ln|x| + \int x^{2/3} dx \\ &= 3\frac{x^5}{5} + \ln|x| + \frac{x^{2/3+1}}{2/3+1} + c \\ &= 3\frac{x^5}{5} + \ln|x| + \frac{3}{5}\sqrt[3]{x^5} + c \end{aligned}$$

Example 5.8.5.

$$\begin{aligned} \int 4x^3(1+2x^4)^4 dx &= \frac{1}{2} \int 8x^3(1+2x^4)^4 dx \\ &= \frac{1}{10}(1+2x^4)^5 + c \end{aligned}$$

Example 5.8.6.

$$\int \frac{dx}{x \ln^3 x} = \int \frac{1}{x} \ln^{-3} x dx = -\frac{1}{2 \ln^2 x} + c$$

Example 5.8.7.

$$\int x^2 e^{x^3} dx = \frac{1}{3} \int 3x^2 e^{x^3} dx = \frac{1}{3} e^{x^3} + c$$

Example 5.8.8.

$$\int \frac{x^3}{1+x^8} dx = \int \frac{x^3}{1+(x^4)^2} dx = \frac{1}{4} \int \frac{4x^3}{1+(x^4)^2} dx = \frac{1}{4} \arctan x^4 + c$$

Example 5.8.9.

$$\begin{aligned}
 \int \frac{x^3 + x + 1}{x^2 + 1} dx &= \int \frac{x(x^2 + 1)}{x^2 + 1} dx + \int \frac{1}{x^2 + 1} dx \\
 &= \int x dx + \int \frac{1}{x^2 + 1} dx \\
 &= \frac{x^2}{2} + \arctan x + c
 \end{aligned}$$

Example 5.8.10.

$$\begin{aligned}
 \int \cos^3 x dx &= \int \cos^2 x \cdot \cos x dx \\
 &= \int (1 - \sin^2 x) \cdot \cos x dx \\
 &= \int \cos x dx - \int \sin^2 x \cdot \cos x dx \\
 &= \sin x - \frac{\sin^3 x}{3} + c
 \end{aligned}$$

Example 5.8.11.

$$\int \frac{\sin x}{\cos x - 4} dx = -\ln |\cos x - 4| + c = \ln(4 - \cos x) + c$$

Example 5.8.12.

$$\int \frac{x^2}{(x^3 + 5)^4} dx = \frac{1}{3} \int 3x^2 (x^3 + 5)^{-4} dx = -\frac{1}{9} \frac{1}{(x^3 + 5)^3} + c$$

Example 5.8.13.

$$\int \frac{1}{x(1 + \log^2 x)} dx = \arctan(\log(x)) + c$$

Example 5.8.14.

$$\int \frac{1}{\tan^4 x \cos^2 x} dx = \int \frac{1}{\cos^2 x} \tan^{-4} x dx = -\frac{1}{3 \tan^3 x} + c$$

Example 5.8.15.

$$\begin{aligned}
 \int \frac{\cos x}{\sqrt{3 - \sin^2 x}} dx &= \int \frac{\cos x}{\sqrt{3} \sqrt{1 - \frac{\sin^2 x}{3}}} dx \\
 &= \int \frac{\frac{\cos x}{\sqrt{3}}}{\sqrt{1 - \left(\frac{\sin x}{\sqrt{3}}\right)^2}} dx = \arcsin\left(\frac{\sin x}{\sqrt{3}}\right) + c
 \end{aligned}$$

Example 5.8.16.

$$\int (2x + 3)^3 dx = \frac{(2x + 3)^4}{8} + c$$

Example 5.8.17.

$$\int \frac{1}{2 - x} dx = -\log |2 - x| + c$$

Example 5.8.18.

$$\begin{aligned}
 \int \frac{x^2}{\sqrt{x^3+2}} dx &= \frac{1}{3} \int 3x^2 (x^3+2)^{-\frac{1}{2}} dx \\
 &= \frac{1}{3} \frac{(x^3+2)^{-\frac{1}{2}+1}}{-\frac{1}{2}+1} \\
 &= \frac{2}{3} (x^3+2)^{\frac{1}{2}} \\
 &= \frac{2}{3} \sqrt{x^3+2} + c
 \end{aligned}$$

Example 5.8.19.

$$\begin{aligned}
 \int \frac{\sqrt{x} + \sqrt[3]{x}}{\sqrt[4]{x}} dx &= \int \frac{x^{1/2} + x^{1/3}}{x^{1/4}} dx \\
 &= \int (x^{1/2-1/4} + x^{1/3-1/4}) dx \\
 &= \int (x^{1/4} + x^{1/12}) dx \\
 &= \frac{x^{1/4+1}}{1/4+1} + \frac{12}{1/12+1} + c \\
 &= \frac{4}{5} \left(x^{5/4} + \frac{12}{13} x^{13/12} \right) + c
 \end{aligned}$$

Example 5.8.20.

$$\int \frac{e^{2x}}{3+e^{2x}} dx = \frac{1}{2} \log(3+e^{2x}) + c$$

Example 5.8.21.

$$\int \frac{e^{2+\sqrt{x}}}{\sqrt{x}} dx = 2 \int \frac{e^{2+\sqrt{x}}}{2\sqrt{x}} dx = 2e^{2+\sqrt{x}} + c$$

Example 5.8.22.

$$\int \frac{x}{\sqrt{x^2+a^2}} dx = \int \frac{2x}{2\sqrt{x^2+a^2}} dx = \sqrt{x^2+a^2} + c$$

Example 5.8.23.

$$\int \frac{e^x}{\sqrt{2e^x+1}} dx = \sqrt{2e^x+1} + c$$

Example 5.8.24.

$$\int \frac{e^x}{4+e^{2x}} dx = \frac{1}{4} \int \frac{e^x}{1+\left(\frac{e^x}{2}\right)^2} dx = \frac{1}{2} \arctan \frac{e^x}{2} + c$$

Example 5.8.25.

$$\int \frac{\cos \log x}{x} dx = \sin \log x + c$$

Example 5.8.26.

$$\int \sqrt[4]{(x-2)^3} dx = \int (x-2)^{3/4} dx = \frac{4}{7} (x-2)^{7/4} + c$$

Example 5.8.27.

$$\begin{aligned}
\int \sqrt{x} \log x \, dx &= \frac{x^{1/2+1}}{1/2+1} \log x - \int \frac{x^{1/2+1}}{1/2+1} \frac{1}{x} \, dx \\
&= \frac{x^{3/2}}{3/2} \log x - \int \frac{x^{1/2}}{3/2} \, dx \\
&= \frac{2}{3} x^{3/2} \log x - \frac{2}{3} \int x^{1/2} \, dx \\
&= \frac{2}{3} x^{3/2} \log x - \frac{2}{3} \cdot \frac{2}{3} x^{3/2} + c \\
&= \frac{2}{3} x^{3/2} \left[\log x - \frac{2}{3} \right] + c
\end{aligned}$$

Example 5.8.28.

$$\begin{aligned}
\int e^{2x} \sin(3x) \, dx &= \frac{1}{2} e^{2x} \sin(3x) - \int \frac{1}{2} e^{2x} \cdot 3 \cos(3x) \, dx = \\
&= \frac{1}{2} e^{2x} \sin(3x) - \frac{3}{2} \left(\frac{1}{2} e^{2x} \cos(3x) - \int \frac{1}{2} e^{2x} \cdot (-3) \sin(3x) \, dx \right) = \\
&= \frac{1}{2} e^{2x} \sin(3x) - \frac{3}{4} e^{2x} \cos(3x) - \frac{9}{4} \int e^{2x} \sin(3x) \, dx \\
\left(1 + \frac{9}{4} \right) \int e^{2x} \sin(3x) \, dx &= e^{2x} \left(\frac{1}{2} \sin(3x) - \frac{3}{4} \cos(3x) \right) \\
\int e^{2x} \sin(3x) \, dx &= \frac{1}{13} e^{2x} (2 \sin(3x) - 3 \cos(3x)) + c
\end{aligned}$$

Example 5.8.29.

$$\begin{aligned}
\int \arcsin x \, dx &= x \arcsin x - \int \frac{x}{\sqrt{1-x^2}} \, dx \\
&= x \arcsin x + \int \frac{-2x}{2\sqrt{1-x^2}} \, dx \\
&= x \arcsin x + \sqrt{1-x^2} + c
\end{aligned}$$

Example 5.8.30.

$$\begin{aligned}
\int \log^2 x \, dx &= x \log^2 x - \int x \log x \frac{1}{x} \, dx \\
&= x \log^2 x - \int \log x \, dx \\
&= x \log^2 x - 2x \log x + 2x + c
\end{aligned}$$

Example 5.8.31.

$$\begin{aligned}
\int (x+2)^2 e^x \, dx &= (x+2)^2 e^x - \int 2(x+2) e^x \, dx \\
&= (x+2)^2 e^x - \left(2(x+2) e^x - \int 2e^x \, dx \right) \\
&= (x+2)^2 e^x - 2(x+2) e^x + 2e^x + c
\end{aligned}$$

Example 5.8.32.

$$\begin{aligned}
\int \arctan x \, dx &= x \arctan x - \int \frac{x}{1+x^2} \, dx \\
&= x \arctan x - \frac{1}{2} \int \frac{2x}{1+x^2} \, dx \\
&= x \arctan x - \frac{1}{2} \log(1+x^2) + c
\end{aligned}$$

Example 5.8.33.

$$\begin{aligned}
\int x \arctan x \, dx &= \frac{1}{2} x^2 \arctan x - \int \frac{1}{2} x^2 \frac{1}{1+x^2} \, dx \\
&= \frac{1}{2} x^2 \arctan x - \frac{1}{2} \int \frac{x^2+1-1}{1+x^2} \, dx \\
&= \frac{1}{2} x^2 \arctan x - \frac{1}{2} \int \left(1 - \frac{1}{1+x^2} \right) \, dx \\
&= \frac{1}{2} x^2 \arctan x - \frac{1}{2} x + \frac{1}{2} \arctan x + c \\
&= \frac{x^2+1}{2} \arctan x - \frac{1}{2} x + c
\end{aligned}$$

Example 5.8.34.

$$\begin{aligned}
\int \log(1+x^2) \, dx &= x \log(1+x^2) - \int \frac{2x^2}{1+x^2} \, dx \\
&= x \log(1+x^2) - 2 \int \frac{x^2+1-1}{1+x^2} \, dx \\
&= x \log(1+x^2) - 2 \int \left(1 - \frac{1}{1+x^2} \right) \, dx \\
&= x \log(1+x^2) - 2x + 2 \arctan x + c
\end{aligned}$$

Example 5.8.35.

$$\begin{aligned}
\int \frac{x}{x^2+4x+3} \, dx &= \int \frac{x}{(x+1)(x+3)} \, dx \\
\frac{x}{(x+1)(x+3)} &= \frac{A}{(x+1)} + \frac{B}{(x+3)} \\
\int \frac{x}{x^2+4x+3} \, dx &= -\frac{1}{2} \int \frac{1}{(x+1)} \, dx + \frac{3}{2} \int \frac{1}{(x+3)} \, dx \\
&= -\frac{1}{2} \log(x+1) + \frac{3}{2} \log(x+3) + c
\end{aligned}$$

Example 5.8.36.

$$\begin{aligned}
\int \frac{x}{(x^2+1)(x-1)} \, dx &= \frac{1}{2} \int \frac{1}{(x-1)} \, dx - \frac{1}{2} \int \frac{x-1}{(x^2+1)} \, dx \\
&= \frac{1}{2} \int \frac{1}{(x-1)} \, dx - \frac{1}{4} \int \frac{2x}{(x^2+1)} \, dx + \frac{1}{2} \int \frac{1}{(x^2+1)} \, dx \\
&= \frac{1}{2} \log|x-1| - \frac{1}{4} \log(x^2+1) + \frac{1}{2} \arctan x + c
\end{aligned}$$

Example 5.8.37.

$$\begin{aligned}
\int \frac{x}{(x-1)^2} dx &= \int \frac{x+1-1}{(x-1)^2} dx \\
&= \int \left(\frac{1}{x-1} + \frac{1}{(x-1)^2} \right) dx \\
&= \log|x-1| - \frac{1}{x-1} + c
\end{aligned}$$

Example 5.8.38.

$$\begin{aligned}
\frac{3x^2-x}{(x+1)^2(x+2)} &= \frac{A}{(x+2)} + \frac{B}{(x+1)} + \frac{D}{(x+1)^2} \\
\int \frac{3x^2-x}{(x+1)^2(x+2)} dx &= \int \frac{14}{x+2} dx - \int \frac{11}{x+1} dx + \int \frac{4}{(x+1)^2} dx \\
&= 14 \log|x+2| - 11 \log|x+1| - \frac{4}{x+1} + c
\end{aligned}$$

Example 5.8.39.

$$\begin{aligned}
\frac{x^2-2x-1}{x^2-4x+4} &= 1 + \frac{2x-5}{x^2-4x+4} \\
\frac{2x-5}{x^2-4x+4} &= \frac{2x-4-1}{x^2-4x+4} = \frac{2x-4}{x^2-4x+4} - \frac{1}{x^2-4x+4} = \frac{2(x-2)}{(x-2)^2} - \frac{1}{(x-2)^2} = \frac{2}{x-2} - \frac{1}{(x-2)^2} \\
\int \frac{x^2-2x-1}{x^2-4x+4} dx &= x + \frac{1}{x-2} + 2 \log|x-2| + c
\end{aligned}$$

Example 5.8.40.

$$\begin{aligned}
\frac{1}{x^2(x^2+1)} &= \frac{1}{x^2} - \frac{1}{x^2+1} \\
\int \frac{1}{x^2(x^2+1)} dx &= -\frac{1}{x} - \arctan x + c
\end{aligned}$$

Example 5.8.41.

$$\begin{aligned}
\frac{1}{(1-x^2)^2} &= \frac{1}{4(x+1)} + \frac{1}{4(x+1)^2} - \frac{1}{4(x-1)} + \frac{1}{4(x-1)^2} \\
\int \frac{1}{(1-x^2)^2} dx &= \frac{1}{4} \log|x+1| - \frac{1}{4} \log|x-1| - \frac{1}{4(x+1)} - \frac{1}{4(x-1)} + c
\end{aligned}$$

Example 5.8.42.

$$\begin{aligned}
\frac{1}{x^4-1} &= \frac{1}{(x-1)(x+1)(1+x^2)} = \frac{1}{4(x-1)} - \frac{1}{4(x+1)} + \frac{1}{2(1+x^2)} \\
\int \frac{dx}{x^4-1} &= \frac{1}{4} \log|1-x| - \frac{1}{4} \log|x+1| - \frac{1}{2} \arctan x + c
\end{aligned}$$

Example 5.8.43.

$$\int \frac{1+x+\sqrt{x}}{1+x\sqrt{x}} dx = 2 \int \frac{t^3+t^2+t}{t^3+1} dt$$

by putting $\sqrt{x} = t$.

$$\begin{aligned}\frac{t^3+t^2+t}{t^3+1} &= 1 + \frac{t^2+t-1}{(t+1)(t^2-t+1)} \\ &= 1 - \frac{1}{3(t+1)} + \frac{4t-2}{3(t^2-t+1)} \\ &= 1 - \frac{1}{3} \frac{1}{1+t} + \frac{2}{3} \frac{2t-1}{t^2-t+1} \\ \int \frac{t^3+t^2+t}{t^3+1} dt &= \int \left(1 - \frac{1}{3} \frac{1}{1+t} + \frac{2}{3} \frac{2t-1}{t^2-t+1} \right) dt \\ &= t - \frac{1}{3} \log|1+t| + \frac{2}{3} \log(t^2-t+1) + c \\ \int \frac{1+x+\sqrt{x}}{1+x\sqrt{x}} dx &= 2\sqrt{x} - \frac{2}{3} \log(1+\sqrt{x}) + \frac{4}{3} \log(x-\sqrt{x}+1) + c\end{aligned}$$

Example 5.8.44.

$$\int \frac{\tan^2 x + 1}{\tan x + 1} dx = \int \frac{1}{t} dt = \ln|t| + c = \ln|\tan x + 1| + c$$

by $t = \tan x + 1$

Example 5.8.45.

$$\begin{aligned}\int \frac{1}{x(\log^2 x - 1)} dx &= \int \frac{1}{t^2 - 1} dt \\ &= \int \frac{1}{(t-1)(t+1)} dt\end{aligned}$$

by $t = \log x$.

$$\begin{aligned}\int \frac{1}{(t-1)(t+1)} dt &= \frac{1}{2} \int \frac{1}{t-1} dt - \frac{1}{2} \int \frac{1}{t+1} dt = \frac{1}{2} \log|t-1| - \frac{1}{2} \log|t+1| + c \\ \int \frac{1}{x(\log^2 x - 1)} dx &= \frac{1}{2} \log|\log x - 1| - \frac{1}{2} \log|\log x + 1| + c\end{aligned}$$

Example 5.8.46. Which one is correct?

$$I = \int \sec^2 x \tan x dx \quad u = \sec x \quad du = \sec x \tan x \quad I = \int 2u du = u^2 = \sec^2 x + C$$

$$I = \int \sec^2 x \tan x dx \quad u = \tan x \quad du = \sec^2 x \quad I = \int 2u du = u^2 = \tan^2 x + C$$

Since $\tan^2 x + C = \sec^2 x - 1 + C = \sec^2 x + C'$, both are correct.

Example 5.8.47.

$$\begin{aligned}\int_0^{\pi/4} \frac{dx}{1-\sin x} &= \int_0^{\pi/4} \frac{1}{1-\sin x} \cdot \frac{1+\sin x}{1+\sin x} dx \\ &= \int_0^{\pi/4} \frac{1+\sin x}{1-\sin^2 x} dx \\ &= \int_0^{\pi/4} \frac{1+\sin x}{\cos^2 x} dx \\ &= \int_0^{\pi/4} (\sec^2 x + \sec x \tan x) dx \\ &= [\tan x + \sec x]_0^{\pi/4} = (1 + \sqrt{2} - (0 + 1)) = \sqrt{2}\end{aligned}\tag{5.1}$$

5.9 Improper Integrals

Consider $I = \int_a^b f(x) dx$, where f is continuous on (a, b) .

If $a = -\infty$ or $b = \infty$ then we say I is an **improper integral of type I**.

If f is unbounded as x approaches to a or b then we say I is an **improper integral of type II**.

Improper Integrals of Type I

Definition 5.9.1. If f is continuous on $[a, \infty)$

$$\int_a^\infty f(x) dx = \lim_{R \rightarrow \infty} \int_a^R f(x) dx$$

Similarly if f is continuous on $(-\infty, b]$


$$\int_{-\infty}^b f(x) dx = \lim_{R \rightarrow -\infty} \int_R^b f(x) dx$$

In either cases, if the limit is finite, we say the integral **converges** and if the limit does not exist, we say the integral **diverges**. If the limit is ∞ or $-\infty$ we say the integral diverges to ∞ or $-\infty$.

Example 5.9.1. Find the area of the region lying under the curve $y = \frac{1}{x^2}$ and above the x -axis to the right of $x = 1$.

Solution. The area is $A = \int_1^\infty \frac{dx}{x^2}$ which is an improper integral of type-I.


$$\begin{aligned} A &= \int_1^\infty \frac{dx}{x^2} = \lim_{R \rightarrow \infty} \int_1^R \frac{dx}{x^2} \\ &= \lim_{R \rightarrow \infty} \left(-\frac{1}{x} \right) \Big|_1^R = \lim_{R \rightarrow \infty} \left(-\frac{1}{R} + 1 \right) = 1 \end{aligned}$$



Example 5.9.2. Find the area of the region lying under the curve $y = \frac{1}{x}$ and above the x -axis to the right of $x = 1$.

Solution. The area is

$$\begin{aligned} A &= \int_1^\infty \frac{dx}{x} = \lim_{R \rightarrow \infty} \int_1^R \frac{dx}{x} \\ &= \lim_{R \rightarrow \infty} \ln x \Big|_1^R = \lim_{R \rightarrow \infty} \ln R = \infty \end{aligned}$$



Definition 5.9.2. For integrals of the form $\int_{-\infty}^{\infty} f(x) dx$, we define

$$\int_{-\infty}^{\infty} f(x) dx = \int_{-\infty}^0 f(x) dx + \int_0^{\infty} f(x) dx$$

The integral on the left converges if and only if both integrals on the right converges.

Note that $\int_{-\infty}^{\infty} f(x) dx$ and $\lim_{R \rightarrow \infty} \int_{-R}^R f(x) dx$ may or may not be equal. For example $\int_{-\infty}^{\infty} x dx$ is divergent according to our definition since $\int_0^{\infty} x dx = \infty$ is divergent. But $\lim_{R \rightarrow \infty} \int_{-R}^R x dx = 0$.

Example 5.9.3. Evaluate $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$.


Solution.

$$I = \int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \int_{-\infty}^0 \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{1}{1+x^2} dx = 2 \int_0^{\infty} \frac{1}{1+x^2} dx$$

since the integrand is an even function.

$$\int_0^{\infty} \frac{1}{1+x^2} dx = \lim_{R \rightarrow \infty} \int_0^R \frac{dx}{1+x^2} = \lim_{R \rightarrow \infty} \arctan R = \frac{\pi}{2}$$

So the answer is $I = \pi$.



Improper Integrals of Type-II

Definition 5.9.3. If f is continuous on the interval $(a, b]$ and is possibly unbounded near a then

$$\int_a^b f(x) dx = \lim_{c \rightarrow a^+} \int_c^b f(x) dx$$


Similarly, if f is continuous on the interval $[a, b)$ and is possibly unbounded near b then

$$\int_a^b f(x) dx = \lim_{c \rightarrow b^-} \int_a^c f(x) dx$$

Example 5.9.4. Find the area of the region lying under $y = 1/\sqrt{x}$, above the x -axis, between $x = 0$ and $x = 1$.

Solution. The area is

$$\int_0^1 \frac{1}{\sqrt{x}} dx = \lim_{c \rightarrow 0^+} \int_c^1 \frac{1}{\sqrt{x}} dx = \lim_{c \rightarrow 0^+} 2x^{1/2} \Big|_c^1 = \lim_{c \rightarrow 0^+} (2 - 2\sqrt{c}) = 2$$



Example 5.9.5. Evaluate $\int_0^2 \frac{dx}{\sqrt{2x-x^2}}$

Solution. By the substitution $u = x - 1$


$$I = \int_0^2 \frac{dx}{\sqrt{2x-x^2}} = \int_0^2 \frac{dx}{\sqrt{1-(x-1)^2}} = \int_{-1}^1 \frac{du}{\sqrt{1-u^2}}$$

By the even symmetry,

$$I = 2 \int_0^1 \frac{du}{\sqrt{1-u^2}}$$

This is an improper integral of Type-II as the integrand is unbounded at $u = 1$.

$$I = 2 \lim_{c \rightarrow 1^-} \int_0^c \frac{du}{\sqrt{1-u^2}} = 2 \lim_{c \rightarrow 1^-} \arcsin u \Big|_0^c = 2 \lim_{c \rightarrow 1^-} \arcsin c = 2 \arcsin 1 = \pi$$



Exercises. Evaluate the following integrals or show they diverge.

1. $\int_0^\infty \cos x dx$

Answer: the integral diverges.

2. $\int_0^\infty e^{-2x} dx$

Answer: $1/2$.

3. $\int_0^\infty x e^{-x} dx$

Answer: 1.

4. $\int_0^{\pi/2} \tan x dx$

Answer: the integral diverges to ∞ .

5. $\int_0^1 \frac{dx}{x}$

Answer: the integral diverges to ∞ .

6. $\int_0^1 \ln x dx$

Answer: -1 .

7. $\int_{-\infty}^\infty x e^{-x^2} dx$

Answer: 0.

5.10 Trigonometric Integrals

Products of Powers of Sines and Cosines

$$\int \sin^m x \cos^n x dx$$

Case 1. If m is odd. Write

$$\sin^m x = \sin^{2k+1} x = (\sin^2 x)^k \sin x = (1 - \cos^2 x)^k \sin x$$

$$\int \sin^m x \cos^n x dx = \int (1 - \cos^2 x)^k \cos^n x \sin x dx = - \int (1 - u^2)^k u^n du$$

where $u = \cos x$.

Example 5.10.1.

$$\begin{aligned}
\int \sin^3 x \cos^2 x dx &= \int \sin^2 x \cos^2 x \sin x dx \\
&= \int (1 - \cos^2 x) (\cos^2 x) (-d(\cos x)) \\
&= \int (1 - u^2) (u^2) (-du) \\
&= \int (u^4 - u^2) du \\
&= \frac{u^5}{5} - \frac{u^3}{3} + C = \frac{\cos^5 x}{5} - \frac{\cos^3 x}{3} + C
\end{aligned}$$

Case 2. If n is odd. In this case

$$\cos^n x = \cos^{2k+1} x = (\cos^2 x)^k \cos x = (1 - \sin^2 x)^k \cos x$$

and

$$\int \sin^m x \cos^n x dx = \int \sin^m x (1 - \sin^2 x)^k \cos x dx = \int u^m (1 - u^2)^k du$$

with $u = \sin x$.**Example 5.10.2.**

$$\begin{aligned}
\int \cos^5 x dx &= \int \cos^4 x \cos x dx = \int (1 - \sin^2 x)^2 d(\sin x) \\
&= \int (1 - u^2)^2 du \\
&= \int (1 - 2u^2 + u^4) du \\
&= u - \frac{2}{3}u^3 + \frac{1}{5}u^5 + C = \sin x - \frac{2}{3}\sin^3 x + \frac{1}{5}\sin^5 x + C
\end{aligned}$$

Case 3. If m and n are both even. Substitute

$$\sin^2 x = \frac{1 - \cos 2x}{2}, \quad \cos^2 x = \frac{1 + \cos 2x}{2}$$

to reduce the integrand to one in lower powers of $\cos 2x$.**Example 5.10.3.**

$$\begin{aligned}
\int \sin^2 x \cos^4 x dx &= \int \left(\frac{1 - \cos 2x}{2} \right) \left(\frac{1 + \cos 2x}{2} \right)^2 dx \\
&= \frac{1}{8} \int (1 - \cos 2x) (1 + 2\cos 2x + \cos^2 2x) dx \\
&= \frac{1}{8} \int (1 + \cos 2x - \cos^2 2x - \cos^3 2x) dx \\
&= \frac{1}{8} \left[x + \frac{1}{2} \sin 2x - I - J \right] \\
I &= \int \cos^2 2x dx = \frac{1}{2} \int (1 + \cos 4x) dx \\
&= \frac{1}{2} \left(x + \frac{1}{4} \sin 4x \right)
\end{aligned}$$

$$\begin{aligned}
 J &= \int \cos^3 2x dx = \int (1 - \sin^2 2x) \cos 2x dx \\
 &= \frac{1}{2} \int (1 - u^2) du = \frac{1}{2} \left(\sin 2x - \frac{1}{3} \sin^3 2x \right)
 \end{aligned}$$

Combining and simplifying gives

$$\int \sin^2 x \cos^4 x dx = \frac{1}{16} \left(x - \frac{1}{4} \sin 4x + \frac{1}{3} \sin^3 2x \right) + C$$

Eliminating Square Roots

Example 5.10.4.

$$\begin{aligned}
 \int_0^{\pi/4} \sqrt{1 + \cos 4x} dx &= \int_0^{\pi/4} \sqrt{2 \cos^2 2x} dx = \int_0^{\pi/4} \sqrt{2} \sqrt{\cos^2 2x} dx \\
 &= \sqrt{2} \int_0^{\pi/4} |\cos 2x| dx = \sqrt{2} \int_0^{\pi/4} \cos 2x dx \\
 &= \sqrt{2} \left[\frac{\sin 2x}{2} \right]_0^{\pi/4} = \frac{\sqrt{2}}{2} [1 - 0] = \frac{\sqrt{2}}{2}
 \end{aligned}$$

Integrals of Powers of $\tan x$ and $\sec x$

Example 5.10.5.

$$\begin{aligned}
 \int \tan^4 x dx &= \int \tan^2 x \cdot \tan^2 x dx = \int \tan^2 x \cdot (\sec^2 x - 1) dx \\
 &= \int \tan^2 x \sec^2 x dx - \int \tan^2 x dx \\
 &= \int \tan^2 x \sec^2 x dx - \int (\sec^2 x - 1) dx \\
 &= \int \tan^2 x \sec^2 x dx - \int \sec^2 x dx + \int dx \\
 &= \frac{1}{3} \tan^3 x - \tan x + x + C
 \end{aligned}$$

where we used by letting $u = \tan x$ and $du = \sec^2 x$

$$I = \int \tan^2 x \sec^2 x dx = \int u^2 du = \frac{1}{3} \tan^3 x + C$$

Example 5.10.6. Find $\int \sec^3 x dx$.

Use the integration by parts

$$u = \sec x, \quad dv = \sec^2 x dx, \quad v = \tan x, \quad du = \sec x \tan x dx$$

to get

$$\begin{aligned}
 I &= \int \sec^3 x dx = \sec x \tan x - \int (\tan x)(\sec x \tan x dx) \\
 &= \sec x \tan x - \int (\sec^2 x - 1) \sec x dx \\
 &= \sec x \tan x + \int \sec x dx - \int \sec^3 x dx \\
 &= \sec x \tan x + \ln |\sec x + \tan x| - I
 \end{aligned}$$

$$I = \int \sec^3 x dx = \frac{1}{2} \sec x \tan x + \frac{1}{2} \ln |\sec x + \tan x| + C$$

Some Examples

Example 5.10.7.

$$\int \sin^3 x dx = \int \sin^2 x \sin x dx = \int (1 - \cos^2 x) \sin x dx = \int \sin x dx - \int \cos^2 x \sin x dx = -\cos x + \frac{1}{3} \cos^3 x + C$$

Example 5.10.8.

$$\begin{aligned} \int \sin^5 x dx &= \int (\sin^2 x)^2 \sin x dx = \int (1 - \cos^2 x)^2 \sin x dx = \int (1 - 2\cos^2 x + \cos^4 x) \sin x dx \\ &= \int \sin x dx - \int 2\cos^2 x \sin x dx + \int \cos^4 x \sin x dx = -\cos x + \frac{2}{3} \cos^3 x - \frac{1}{5} \cos^5 x + C \end{aligned}$$

Example 5.10.9.

$$\begin{aligned} \int \sin^3 x \cos^3 x dx &= \int \sin^3 x \cos^2 x \cos x dx = \int \sin^3 x (1 - \sin^2 x) \cos x dx = \int \sin^3 x \cos x dx - \int \sin^5 x \cos x dx \\ &= \frac{1}{4} \sin^4 x - \frac{1}{6} \sin^6 x + C \end{aligned}$$

Example 5.10.10.

$$\begin{aligned} \int \cos^2 x dx &= \int \frac{1 + \cos 2x}{2} dx = \frac{1}{2} \int (1 + \cos 2x) dx = \frac{1}{2} \int dx + \frac{1}{2} \int \cos 2x dx = \frac{1}{2} \int dx + \frac{1}{4} \int \cos 2x \cdot 2 dx \\ &= \frac{1}{2} x + \frac{1}{4} \sin 2x + C \end{aligned}$$

Example 5.10.11.

$$\int_0^\pi \sqrt{1 - \cos 2x} dx = \int_0^\pi \sqrt{2} |\sin x| dx = \int_0^\pi \sqrt{2} \sin x dx = [-\sqrt{2} \cos x]_0^\pi = \sqrt{2} + \sqrt{2} = 2\sqrt{2}$$

Example 5.10.12.

$$\int \sec^3 x \tan x dx = \int \sec^2 x \sec x \tan x dx = \frac{1}{3} \sec^3 x + C$$

where we used $u = \sec x$ substitution.

Example 5.10.13.

$$\begin{aligned} \int \sec^4 x \tan^2 x dx &= \int \sec^2 x \tan^2 x \sec^2 x dx = \int (\tan^2 x + 1) \tan^2 x \sec^2 x dx \\ &= \int \tan^4 x \sec^2 x dx + \int \tan^2 x \sec^2 x dx = \frac{1}{5} \tan^5 x + \frac{1}{3} \tan^3 x + C \end{aligned}$$

Example 5.10.14.

$$\begin{aligned} \int \sec^4 \theta d\theta &= \int (1 + \tan^2 \theta) \sec^2 \theta d\theta = \int \sec^2 \theta d\theta + \int \tan^2 \theta \sec^2 \theta d\theta = \tan \theta + \frac{1}{3} \tan^3 \theta + C \\ &= \tan \theta + \frac{1}{3} \tan \theta (\sec^2 \theta - 1) + C = \frac{1}{3} \tan \theta \sec^2 \theta + \frac{2}{3} \tan \theta + C \end{aligned}$$

Example 5.10.15.

$$\begin{aligned}\int \tan^5 x dx &= \int \tan^4 x \tan x dx = \int (\sec^2 x - 1)^2 \tan x dx = \int (\sec^4 x - 2 \sec^2 x + 1) \tan x dx \\&= \int \sec^4 x \tan x dx - 2 \int \sec^2 x \tan x dx + \int \tan x dx \\&= \int \sec^3 x \sec x \tan x dx - 2 \int \sec x \sec x \tan x dx + \int \tan x dx = \frac{1}{4} \sec^4 x - \sec^2 x + \ln |\sec x| + C \\&= \frac{1}{4} (\tan^2 x + 1)^2 - (\tan^2 x + 1) + \ln |\sec x| + C = \frac{1}{4} \tan^4 x - \frac{1}{2} \tan^2 x + \ln |\sec x| + C\end{aligned}$$

Chapter 6

Applications of Integration


6.1 Volumes Using Cross-Sections

Solids of Revolution: The Disk Method

Suppose the graph of $y = f(x)$, $a \leq x \leq b$ is revolved around x -axis. Let $a = x_0 < x_1 < \cdots x_n = b$. Approximating the volume of the revolved region by disks,

$$V \approx \sum_{k=1}^n \pi f(x_k)^2 \Delta x_k \rightarrow V = \int_a^b \pi f(x)^2 dx.$$


Example 6.1.1. Find the volume of the solid obtained by revolving the curve $y = \sqrt{x}$, $0 \leq x \leq 4$ around the x -axis.



$$V = \int_0^4 \pi (\sqrt{x})^2 dx = 8\pi$$


Example 6.1.2. Find the volume of the sphere with radius a .

The sphere is obtained by revolving the graph of $y = \sqrt{a^2 - x^2}$, $-a \leq x \leq a$ around the x -axis.



$$V = \int_{-a}^a \pi(a^2 - x^2) dx = \frac{4}{3} \pi a^3.$$

Example 6.1.3. Find the volume of the solid generated by revolving the region bounded by $y = \sqrt{x}$ and the lines $y = 1$, $x = 4$ about the line $y = 1$.



The distance to the line of revolution is $R(x) = \sqrt{x} - 1$.

$$V = \int_1^4 \pi(\sqrt{x} - 1)^2 dx = \pi \int_1^4 (x - 2\sqrt{x} + 1) dx = \frac{7\pi}{6}.$$

Example 6.1.4. Find the volume of the solid generated by revolving the region between the y -axis and the curve $x = 2/y$, $1 \leq y \leq 4$ about the y -axis.

1st method.

$$V = \int_1^4 \pi(R(y))^2 dy = \pi \int_1^4 \left(\frac{2}{y}\right)^2 dy = 3\pi.$$

2nd method. Use $x \leftrightarrow y$.

Example 6.1.5. Find the volume of the solid generated by revolving the region between the parabola $x = y^2 + 1$ and the line $x = 3$ about the line $x = 3$.

The intersection of $x = y^2 + 1$ and $x = 3$ is at $x = \pm\sqrt{2}$. $R(y) = 3 - (y^2 + 1)$.

$$V = \int_{-\sqrt{2}}^{\sqrt{2}} \pi(R(y))^2 dy = \int_{-\sqrt{2}}^{\sqrt{2}} \pi(2 - y^2)^2 dy = \frac{64\pi\sqrt{2}}{15}$$

Solids of Revolution: The Washer Method

Example 6.1.6. The region bounded by the curve $y = x^2 + 1$ and the line $y = -x + 3$ is revolved about the x -axis. Find the volume of the solid.

The volume = The volume of the outer solid - the volume of the inner solid.

$$V = \int_{-2}^1 \pi(R_{\text{outer}}(x))^2 dx - \int_{-2}^1 \pi(R_{\text{inner}}(x))^2 dx = \int_{-2}^1 \pi((-x + 3)^2 - (x^2 + 1)^2) dx = \frac{117\pi}{5}.$$

Example 6.1.7. The region bounded by the parabola $y = x^2$ and the line $y = 2x$ in the first quadrant is revolved about the y -axis to generate a solid. Find the volume.

$$V = \int_0^4 \pi\left((\sqrt{y})^2 - \left(\frac{y}{2}\right)^2\right) dy = \pi \int_0^4 \left(y - \frac{y^2}{4}\right) dy = \frac{8\pi}{3}.$$

Example 6.1.8. Find the volume of the solid generated by revolving the regions bounded by $y = 2\sqrt{x}$, $y = 2$, $x = 0$ about the x -axis.

$$\begin{aligned} r(x) = 2\sqrt{x} \text{ and } R(x) = 2 &\Rightarrow V = \int_0^1 \pi ([R(x)]^2 - [r(x)]^2) dx \\ &= \pi \int_0^1 (4 - 4x) dx = 4\pi \left[x - \frac{x^2}{2} \right]_0^1 = 4\pi \left(1 - \frac{1}{2} \right) = 2\pi \end{aligned} \quad (6.1)$$

Example 6.1.9. Find the volume of the solid generated by revolving the region enclosed by the triangle with vertices $(1, 0)$, $(2, 1)$, and $(1, 1)$ region about the y -axis.

$$\begin{aligned} r(y) = 1 \text{ and } R(y) = 1 + y &\Rightarrow V = \int_0^1 \pi ([R(y)]^2 - [r(y)]^2) dy \\ &= \pi \int_0^1 [(1 + y)^2 - 1] dy = \pi \int_0^1 (1 + 2y + y^2 - 1) dy \\ &= \pi \int_0^1 (2y + y^2) dy = \pi \left[y^2 + \frac{y^3}{3} \right]_0^1 = \pi \left(1 + \frac{1}{3} \right) = \frac{4\pi}{3} \end{aligned} \quad (6.2)$$

Exercises 6.1 from Thomas. 25, 30, 39, 42, 52

6.2 Volumes Using Cylindrical Shells

Solids of Revolution: The Disk Method

A cylindrical shell with height h , outer radius $x + \Delta x$ and inner radius x has volume

$$V_{\text{shell}} = \pi(x + \Delta x)^2 h - \pi x^2 h = \pi(2x\Delta x + (\Delta x)^2)h$$

If Δx is very small then $(\Delta x)^2$ is very small compared to Δx .

$$V_{\text{shell}} \approx 2\pi x \Delta x h$$

Suppose the graph of $y = f(x)$, $a \leq x \leq b$ is revolved around the y -axis. Let $a = x_0 < x_1 < \cdots < x_n = b$. Approximating the volume of the revolved region by cylindrical shells, the height of each shell is $f(x_k)$ and the outer radius is $x_k + \Delta x_k$ and inner radius x_k .

$$V \approx \sum_{k=1}^n 2\pi x f(x_k) \Delta x_k \xrightarrow{\Delta x_k \rightarrow 0} V = \int_a^b 2\pi x f(x) dx = \int_a^b 2\pi (\text{shell radius}) \cdot (\text{shell height}) dx.$$

Example 6.2.1. The region bounded by the curve $y = \sqrt{x}$, the x -axis, and the line $x = 4$ is revolved about the y -axis to generate a solid. Find the volume of the solid.

$$\begin{aligned} V &= \int_0^4 2\pi(x)(\sqrt{x}) dx \\ &= 2\pi \int_0^4 x^{3/2} dx = 2\pi \left[\frac{2}{5} x^{5/2} \right]_0^4 = \frac{128\pi}{5} \end{aligned} \quad (6.3)$$

Example 6.2.2. The region bounded by the curve $y = \sqrt{x}$, the x -axis, and the line $x = 4$ is revolved about the x -axis to generate a solid. Find the volume of the solid by the shell method.

$$\begin{aligned}
 V &= \int_a^b 2\pi \left(\begin{array}{c} \text{shell} \\ \text{radius} \end{array} \right) \left(\begin{array}{c} \text{shell} \\ \text{height} \end{array} \right) dy \\
 &= \int_0^2 2\pi(y)(4 - y^2) dy \\
 &= 2\pi \int_0^2 (4y - y^3) dy \\
 &= 2\pi \left[2y^2 - \frac{y^4}{4} \right]_0^2 = 8\pi
 \end{aligned} \tag{6.4}$$

Exercises 6.2 from Thomas. 15, 18, 21, 22, 33, 36.

6.3 Arc Length

Finding the length of the curve $y = f(x)$, $a \leq x \leq b$.




Figure 6.1: From Thomas.

Partition the interval $a = x_0 < x_1 < \cdots < x_n = b$. The approximation of the arc length is

$$\sum \sqrt{(\Delta x_k)^2 + (\Delta y_k)^2}$$

where $\Delta x_k = x_k - x_{k-1}$ and $\Delta y_k = y_k - y_{k-1}$. By the Mean Value Theorem there is a point c_k with $x_{k-1} < c_k < x_k$ such that

$$\Delta y_k = f'(c_k) \Delta x_k$$

So the approximation becomes

$$\sum \sqrt{(\Delta x_k)^2 + (\Delta y_k)^2} = \sum \sqrt{(\Delta x_k)^2 + (f'(c_k) \Delta x_k)^2} = \sum \sqrt{1 + (f'(c_k))^2} \Delta x_k \xrightarrow{\Delta x_k \rightarrow 0} \int_a^b \sqrt{1 + (f'(x))^2} dx$$

The arc length is defined as

$$\int_a^b \sqrt{1 + (f'(x))^2} dx$$

Example 6.3.1. Find the length of the curve

$$y = \frac{4\sqrt{2}}{3}x^{3/2} - 1, \quad 0 \leq x \leq 1 \quad (6.5)$$

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \int_0^1 \sqrt{1 + 8x} dx \\ &= \frac{2}{3} \cdot \frac{1}{8} (1 + 8x)^{3/2} \Big|_0^1 = \frac{13}{6} \approx 2.17 \end{aligned} \quad (6.6)$$

Example 6.3.2. Find the length of the graph

$$f(x) = \frac{x^3}{12} + \frac{1}{x}, \quad 1 \leq x \leq 4 \quad (6.7)$$

$$\begin{aligned} L &= \int_1^4 \sqrt{1 + [f'(x)]^2} dx = \int_1^4 \left(\frac{x^2}{4} + \frac{1}{x^2} \right) dx \\ &= \left[\frac{x^3}{12} - \frac{1}{x} \right]_1^4 = \left(\frac{64}{12} - \frac{1}{4} \right) - \left(\frac{1}{12} - 1 \right) = \frac{72}{12} = 6 \end{aligned} \quad (6.8)$$

Example 6.3.3. Find the length of the curve

$$y = \frac{1}{2}(e^x + e^{-x}), \quad 0 \leq x \leq 2 \quad (6.9)$$

$$\begin{aligned} L &= \int_0^2 \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \int_0^2 \frac{1}{2}(e^x + e^{-x}) dx \\ &= \frac{1}{2} [e^x - e^{-x}]_0^2 = \frac{1}{2}(e^2 - e^{-2}) \approx 3.63 \end{aligned} \quad (6.10)$$

The length of $x = g(y)$, $c \leq y \leq d$ is

$$L = \int_c^d \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy = \int_c^d \sqrt{1 + [g'(y)]^2} dy \quad (6.11)$$

Exercises 6.3 from Thomas. 2, 13, 14, 15, 17.

6.4 Areas Of Surfaces Of Revolution




Figure 6.2: From Thomas.

The surface are of the above shape is

$$2\pi \left(\frac{y_1 + y_2}{2} \right) L,$$

The surface are of the region obtained be revolving $y = f(x)$, $a \leq x \leq b$ around the x -axis is

$$\sum 2\pi \left(\frac{f(x_{k-1}) + f(x_k)}{2} \right) \sqrt{(\Delta x_k)^2 + (\Delta y_k)^2} \xrightarrow{\Delta x_k \rightarrow 0} \int_a^b 2\pi f(x) \sqrt{1 + (f'(x))^2} dx$$

Example 6.4.1. Find the area of the surface generated by revolving $y = 2\sqrt{x}$, $1 \leq x \leq 2$ about the x -axis.

$$\begin{aligned} \sqrt{1 + \left(\frac{dy}{dx} \right)^2} &= \sqrt{1 + \left(\frac{1}{\sqrt{x}} \right)^2} \\ &= \sqrt{1 + \frac{1}{x}} = \sqrt{\frac{x+1}{x}} = \frac{\sqrt{x+1}}{\sqrt{x}} \end{aligned} \tag{6.12}$$

$$\begin{aligned} S &= \int_1^2 2\pi \cdot 2\sqrt{x} \frac{\sqrt{x+1}}{\sqrt{x}} dx = 4\pi \int_1^2 \sqrt{x+1} dx \\ &= 4\pi \cdot \frac{2}{3} (x+1)^{3/2} \Big|_1^2 = \frac{8\pi}{3} (3\sqrt{3} - 2\sqrt{2}) \end{aligned} \tag{6.13}$$

For revolution about the y -axis, we interchange x and y . Exercises 6.4 from Thomas. 1, 9, 13, 22