

## Plan Quality Algorithm: Emami TD 50-5 [56 Max Possible] [56 Metrics] (Page 1 of 6)

1	 V (%) D(Gy)	Dose (Gy) covering 100 (%) of the BLADDER (SYMPTOMATIC BLADDER CONTRACTURE)	1	
2	 V (%) D(Gy)	Dose (Gy) covering 66.666 (%) of the BLADDER (SYMPTOMATIC BLADDER CONTRACTURE)	1	
3	 V (%) D(Gy)	Dose (Gy) covering 100 (%) of the BRACHIAL PLEXUS (CLINICALLY APPARENT NERVE DAMAGE)	1	
4	 V (%) D(Gy)	Dose (Gy) covering 66.666 (%) of the BRACHIAL PLEXUS (CLINICALLY APPARENT NERVE DAMAGE)	1	
5	 V (%) D(Gy)	Dose (Gy) covering 33.333 (%) of the BRACHIAL PLEXUS (CLINICALLY APPARENT NERVE DAMAGE)	1	
6	 V (%) D(Gy)	Dose (Gy) covering 100 (%) of the BRAIN (NECROSIS)	1	
7	 V (%) D(Gy)	Dose (Gy) covering 66.666 (%) of the BRAIN (NECROSIS)	1	
8	 V (%) D(Gy)	Dose (Gy) covering 33.333 (%) of the BRAIN (NECROSIS)	1	
9	 V (%) D(Gy)	Dose (Gy) covering 100 (%) of the BRAINSTEM (NECROSIS)	1	
10	 V (%) D(Gy)	Dose (Gy) covering 100 (%) of the CAUDA EQUINA (CLINICALLY APPARENT NERVE DAMAGE)	1	


## Plan Quality Algorithm: Emami TD 50-5 [56 Max Possible] [56 Metrics] (Page 2 of 6)

11	 V (%) D(Gy)	Dose (Gy) covering 100 (%) of the COLON (OBSTRUCTION, PERFORATION, ULCERATION)	1		$\leq 55 \geq 55.00$
12	 V (%) D(Gy)	Dose (Gy) covering 33.33 (%) of the COLON (OBSTRUCTION, PERFORATION, ULCERATION)	1		$\leq 65 \geq 65.00$
13	 V (%) D(Gy)	Dose (Gy) covering 100 (%) of the EAR (ACUTE SEROUS OTITIS)	1		$\leq 40 \geq 40.00$
14	 V (%) D(Gy)	Dose (Gy) covering 66.666 (%) of the EAR (ACUTE SEROUS OTITIS)	1		$\leq 40 \geq 40.00$
15	 V (%) D(Gy)	Dose (Gy) covering 33.333 (%) of the EAR (ACUTE SEROUS OTITIS)	1		$\leq 40 \geq 40.00$
16	 V (%) D(Gy)	Dose (Gy) covering 100 (%) of the EAR (CHRONIC SEROUS OTITIS)	1		$\leq 65 \geq 65.00$
17	 V (%) D(Gy)	Dose (Gy) covering 66.666 (%) of the EAR (CHRONIC SEROUS OTITIS)	1		$\leq 65 \geq 65.00$
18	 V (%) D(Gy)	Dose (Gy) covering 33.333 (%) of the EAR (CHRONIC SEROUS OTITIS)	1		$\leq 65 \geq 65.00$
19	 V (%) D(Gy)	Dose (Gy) covering 100 (%) of the ESOPHAGUS (STRICTURE, PERFORATION)	1		$\leq 68 \geq 68.00$
20	 V (%) D(Gy)	Dose (Gy) covering 66.666 (%) of the ESOPHAGUS (STRICTURE, PERFORATION)	1		$\leq 70 \geq 70.00$

## Plan Quality Algorithm: Emami TD 50-5 [56 Max Possible] [56 Metrics] (Page 3 of 6)

21	 V (%) D(Gy)	Dose (Gy) covering 33.333 (%) of the ESOPHAGUS (STRICTURE, PERFORATION)	1	
22	 V (%) D(Gy)	Dose (Gy) covering 100 (%) of the FEMORAL HEAD (NECROSIS)	1	
23	 V (%) D(Gy)	Dose (Gy) covering 100 (%) of the HEART (PERICARDITIS)	1	
24	 V (%) D(Gy)	Dose (Gy) covering 66.666 (%) of the HEART (PERICARDITIS)	1	
25	 V (%) D(Gy)	Dose (Gy) covering 33.333 (%) of the HEART (PERICARDITIS)	1	
26	 V (%) D(Gy)	Dose (Gy) covering 100 (%) of the KIDNEY (CLINICAL NEPHRITIS)	1	
27	 V (%) D(Gy)	Dose (Gy) covering 66.666 (%) of the KIDNEY (CLINICAL NEPHRITIS)	1	
28	 V (%) D(Gy)	Dose (Gy) covering 100 (%) of the LARYNX (NECROSIS)	1	
29	 V (%) D(Gy)	Dose (Gy) covering 66.666 (%) of the LARYNX (NECROSIS)	1	
30	 V (%) D(Gy)	Dose (Gy) covering 33.333 (%) of the LARYNX (NECROSIS)	1	












## Plan Quality Algorithm: Emami TD 50-5 [56 Max Possible] [56 Metrics] (Page 4 of 6)



## Plan Quality Algorithm: Emami TD 50-5 [56 Max Possible] [56 Metrics] (Page 5 of 6)

41	V (%)  D(Gy)	Dose (Gy) covering 100 (%) of the PAROTID GLAND (XEROSTOMIA)	1	≤ 46 ≥ 46.00
42	V (%)  D(Gy)	Dose (Gy) covering 66.666 (%) of the PAROTID GLAND (XEROSTOMIA)	1	≤ 46 ≥ 46.00
43	V (%)  D(Gy)	Dose (Gy) covering 100 (%) of the RECTUM (SEVERE PROCTITIS, NECROSIS, FISTULA, STENOSIS)	1	≤ 80 ≥ 80.00
44	V (%)  D(Gy)	Dose (Gy) covering 100 (%) of the RETINA (BLINDNESS)	1	≤ 65 ≥ 65.00
45	V (%)  D(Gy)	Dose (Gy) covering 33.333 (%) of the RIBCAGE (PATHOLOGIC FRACTURE)	1	≤ 65 ≥ 65.00
46	V (cc)  D(Gy)	Dose (Gy) covering 100 (cc) of the SKIN (NECROSIS, ULCERATION)	1	≤ 65 ≥ 65.00
47	V (cc)  D(Gy)	Dose (Gy) covering 10 (cc) of the SPINAL CORD (MYELITIS, NECROSIS)	1	≤ 70 ≥ 70.00
48	V (cc)  D(Gy)	Dose (Gy) covering 5 (cc) of the SPINAL CORD (MYELITIS, NECROSIS)	1	≤ 70 ≥ 70.00
49	V (%)  D(Gy)	Dose (Gy) covering 100 (%) of the SMALL INTESTINE (OBSTRUCTION, PERFORATION)	1	≤ 55 ≥ 55.00
50	V (%)  D(Gy)	Dose (Gy) covering 33.333 (%) of the SMALL INTESTINE (OBSTRUCTION, PERFORATION)	1	≤ 60 ≥ 60.00

## Plan Quality Algorithm: Emami TD 50-5 [56 Max Possible] [56 Metrics] (Page 6 of 6)

51	 V(%) D(Gy)	Dose (Gy) covering 100 (%) of the STOMACH (ULCERATION, PERFORATION)	1	
52	 V(cc) D(Gy)	Dose (Gy) covering 66.666 (cc) of the STOMACH (ULCERATION, PERFORATION)	1	
53	 V(%) D(Gy)	Dose (Gy) covering 33.333 (%) of the STOMACH (ULCERATION, PERFORATION)	1	
54	 V(%) D(Gy)	Dose (Gy) covering 100 (%) of the TMJ MANDIBLE (LIMITED JOINT FUNCTION)	1	
55	 V(%) D(Gy)	Dose (Gy) covering 66.666 (%) of the TMJ MANDIBLE (LIMITED JOINT FUNCTION)	1	
56	 V(%) D(Gy)	Dose (Gy) covering 33.333 (%) of the TMJ MANDIBLE (LIMITED JOINT FUNCTION)	1	