


Homework 2: Sea level variability in the Pacific Due November 3, 2020

Altimeter Regional Sea Level Trends

Sea level trends (1993-2018)



<https://uhslc.soest.hawaii.edu>

- 1) Load the file `ALT_Pacific.mat`, which contains the sea level anomaly for the Pacific from multiple satellite missions (data source: <http://marine.copernicus.eu/services-portfolio/access-to-products/>). The data are contained in "ALT" a structure array. Type `ALTreadme` for a description of the variables.
- 2) Load the file `TGdata.mat` which contains sea level from tide gauge data at San Diego, Honolulu, and Mera, Japan. The data are contained in "TG" a structure array. Type `TGreadme` for a description of the variables.
- 3) Fit a linear trend, mean, and seasonal cycle from each ALT and TG time series using `regress.m`. Plot maps (using `imagesc` or `pcolor`) of ALT sea level trends and the standard deviation of the seasonal cycle.
- 4) Form sea level anomalies by removing the trend, mean, and seasonal cycle from each ALT and TG time series.
- 5) For each TG station, compute the correlation with every ocean grid point in the ALT array. Plot a map of the correlation coefficients.
- 6) Estimate integral time scales for each of the TG stations. Base on these time scales, identify correlations on your map that are significant at the 95% level.
- 7) For each TG station, compute the regression ($ALT = b * TG$) with every ocean grid point in the ALT array. Plot a map of the regression coefficients.

8) Describe dominant patterns in the correlation and regression maps.