

Calculus Skeleton Notes

Contents

September 9: Average Rates of Change	1
September 11: 1.1 - Limits	3

Average Rates of Change

Goals: Compute average rates of change, slopes of secant lines, and motivate the idea of limits

Before Class Videos: None today, starting next class

Example 1: I left my house at time $t = 0$. 30 minutes later, I was at UVic which is 12 km away.

a) What was my average speed?

b) Sketch a possible graph

c) How could I approximate my speed at $t = 10$ minutes?

Example 2: What is average rate of change of $f(x) = x^2$ on $[2, 4]$?

Example 3: Find the equation of the line through (x_1, y_1) and (x_2, y_2) .

Example 4: Consider the graph of a function $f(x)$. How do I write an equation for the secant line through $(x_1, f(x_1))$ and $(x_2, f(x_2))$?

Big Idea: As x_2 gets closer to x_1 the slope of the secant line approaches the slope of the tangent line at x_1 . We need to investigate and define this limiting process more precisely. This DESMOS visualizes the idea: <https://www.desmos.com/calculator/oi7poyhodr>

Suggested after class problems from APEX: None today, but review the Course Outline on Brightspace

1.1 - Limits

Goals: Describe limits graphically, numerically, and algebraically.

Before Class Videos: 1.1.1 and 1.1.2

Example 1: Consider these three functions:

$$f(x) = x + 1, \quad g(x) = \frac{x^2 - 1}{x - 1} = \frac{(x + 1)(x - 1)}{x - 1}, \quad h(x) = \begin{cases} x + 1, & x \neq 1, \\ 3, & x = 1. \end{cases}$$

How are these similar or different?

Definition:

$\lim_{x \rightarrow a} f(x) = L$ if $f(x)$ can be made arbitrarily close to L for all values of x sufficiently close to a .

Example 2: Compute the limit as $x \rightarrow 1$ of $f(x)$, $g(x)$, and $h(x)$.

Example 3: Use a table of values to estimate $\lim_{x \rightarrow 1} (x + 1)$.

Example 4: Use a table of values to estimate $\lim_{x \rightarrow 0} \frac{1}{x}$.

Example 5: Investigate

$$f(x) = \begin{cases} x, & x < 1, \\ x + 1, & x \geq 1. \end{cases}$$

Definition:

$\lim_{x \rightarrow a^+} f(x)$ means $f(x)$ gets arbitrarily close to L for all x sufficiently close to a from the right,
i.e. $x > a$.

Example 6: Investigate limits of

$$f(x) = \sin\left(\frac{1}{x}\right)$$

Example 7: Graphically investigate limits of $\frac{x^2-1}{x^2-x-2}$

Suggested after class problems from APEX: Section 1.1 problems 9,11,15