Variational Bayesian Methods (in Neuroscience)

Tyler Benster & Aaron Andalman

Deisseroth (Tyler & Aaron) and Druckmann (Tyler) Labs



Computational Neuroscience Journal Club Stanford University

July 10, 2019



Roadmap



∟Roadmap

Introduction: Why care about the distribution of data?

Problem: Analyzing high dimensional data is hard

Solution: Variational Methods

Example: variational autoencoder

Discussion: Neuroscience applications

Introduction: Why care about the distribution of data?

Problem: Analyzing high dimensional data is hard

Solution: Variational Methods

Solution: Variational Methods

Example: variational autoencoder

Discussion: Neuroscience applications

Introduction: Why care about the distribution of data?



Introduction: Why care about the distribution of data?

Introduction: Why care about the distribution of data? 2019-07-15 -Introduction: Why care about the distribution of data? -Introduction: Why care about the distribution


of data?


Introduction: Why care about the distribution of data

The probability distribution revolution



- Karl Pearson (1857-1936) came with the idea that scientific measurements should be conceived as coming from probability distributions.
- Scientific measurements are just random reflections of the underlying truth that is the distribution.
- "A great book on the history of statistics" → Aaron







The power of probability distributions



Distributions allow scientists to:

- ▶ Understand scientific measurement
- ▶ Predict the probability of specific data
- ► Test specific hypothesis (p-values)
- Produce generative models
- Better conceptual understanding data.

Introduction: Why care about the distribution of data? 2019-07-15 Introduction: Why care about the distribution of data? The power of probability distributions

- Predict the probability of specific data
- Distributions allow orientists to
- Produce generative models ► Better conceptual understanding data

Introduction: Why care about the distribution of data?

Estimating distributions from data



Low-Dimensional:

Great tools to fit and understand the underlying probability distribution of data.

High-Dimensional:

- ► In some cases, classical statistical tools are insufficient.
- Problematic for modern neuroscience:
- Thousands of electrodes.
- Millions of voxels.

Introduction: Why care about the distribution of data?
Introduction: Why care about the distribution of data?
Estimating distributions from data

 Great tools to fit and understand the underlying probability distribution of

ind In some cases, class derlying statistical tools are tion of insufficient.

Problematic for m neuroscience:

Thousands of electronic

Variational Bayesian Methods



Estimate the probability distribution of high-dimensional neural data.

- ► Compute the probability of observing a particular neural state.
- ► Sample neural states/trajectory from the estimate.
- ► Generate statistics / test hypotheses
- ▶ Estimate latent factors/states that drive the observations.
- ► Reduce dimensionality (with advantages over other methods, e.g. PCA, T-SNE)

Introduction: Why care about the distribution of data?
Introduction: Why care about the distribution of data?

data?
Variational Bayesian Methods

Estimate the probability distribution of high-dimensional neural data.

- ► Compute the probability of observing a particular neural state
- Connecte statistics / test benetheres
- Generate statistics / test hypoth
- Estimate latent factors/states that drive the observations
- Reduce dimensionality (with advantages over other me e.g. PCA, T-SNE)

Problem: Analyzing high dimensional data is hard




Introduction: Why care about the distribution of data?

Problem: Analyzing high dimensional data is hard

Solution: Variational Methods

Example: variational autoencoder

Discussion: Neuroscience applications



Problem: Analyzing high dimensional data is hard

tion: Why care about the distribution of data?

Problem: Analyzing high dimensional data is hard


Solution: Variational Meth

Example: variational autoencoder

Discussion: Neuroscience applications

Approximating the area of a circle with Monte Carlo





40/50 samples in circle gives area of $4*0.8=3.2\approx\pi.\sim2\%$ error

CCO 1.0, wikimedia, MonteCarloIntegrationCircle.svg

Problem: Analyzing high dimensional data is hard

Problem: Analyzing high dimensional data is hard


nte

40/50 samplini in occide givos area of coccide givos area of

Approximating the area of a circle with Monte Carlo

Useful for analytically intractable integrals





Robert Lin - Monte Carlo Integration

Problem: Analyzing high dimensional data is hard -Problem: Analyzing high dimensional data is hard

Useful for analytically intractable integrals

Extension to unit hypercube



We can approximate a high-dimensional integral using a Monte Carlo approximation:

$$\int_0^1 \cdots \int_0^1 g(x_1, \ldots, x_n) dx_1, \ldots, dx_n \approx \frac{1}{N} \sum_{i=1}^N g(\bar{x}_i)$$

where $\bar{x}_1, \dots, \bar{x}_N \sim \mathcal{U}(0,1)$ is the ith random sample


Problem: Analyzing high dimensional data is hard \cupL$ Problem: Analyzing high dimensional data is hard

We can approximate a high-dimensional integral using a Mont Carlo approximation: $\int_0^1 \cdots \int_0^1 g(x_1, \dots, x_n) dx_1, \dots, dx_n \approx \frac{1}{N} \sum_{j=1}^N g(\bar{x}_j)$ where $\bar{x}_1, \dots, \bar{x}_N \sim U(0,1)$ is the \bar{x}^h random sample

Extension to unit hypercube

Performance scales poorly with number of dimensions





1000 samples, estimate = 4.072, actual = 4.189, \sim 3% error

http://www2.hawaii.edu/ yuxian/phys305/a6/

Problem: Analyzing high dimensional data is hard -Problem: Analyzing high dimensional data is hard

-Performance scales poorly with number of dimensions

Solution: Variational Methods



Introduction: Why care about the distribution of data?

Problem: Analyzing high dimensional data is hard

Solution: Variational Methods

Example: variational autoencoder

Discussion: Neuroscience applications

Solution: Variational Methods —Solution: Variational Methods

-Solution: Variational Methods

Introduction: Why care about the distribution of data?

Problem: Analyzing high dimensional data is hard

Solution: Variational Methods

Example: variational autoencoder

Solution: Variational Methods

Tyler Benster & Aaron Andalman, CNJC

Alternate approach: Variational Bayes












Alternate approach: Variational Bayes



Example: variational autoencoder



Introduction: Why care about the distribution of data

Problem: Analyzing high dimensional data is hard

Solution: Variational Methods

Example: variational autoencoder

Discussion: Neuroscience applications

Example: variational autoencoder

Example: variational autoencoder

-Example: variational autoencoder

troduction: Why care about the distribution of data?

Solution: Variational Methods

Example: variational autoencoder

Example: variational autoencoder

Tyler Benster & Aaron Andalman, CNJC

Demo



https://bit.ly/2LcEhow

└ Demo

https://bit.ly/2LcEhow

Summary



- ▶ Problem: Analyzing high dimensional data is hard
- ► Solution: Variational Methods
- ► Example: variational autoencoder

Example: variational autoencoder

Example: variational autoencoder

—Summary

- Problem: Analyzing high dimensional data is hard
 - lution: Variational Methods
- Example: variational autoencoder

Discussion: Neuroscience applications



Introduction: Why care about the distribution of data?

Problem: Analyzing high dimensional data is hard

Solution: Variational Methods

Example: variational autoencoder

Discussion: Neuroscience applications

Discussion: Neuroscience applications — Discussion: Neuroscience applications

-Discussion: Neuroscience applications

oduction: Why care about the distribution of data?

Colorina Visitational Markedo

Solution: Variational Metho

Discussion: Neuroscience applications

Example: LFADS



Discussion: Neuroscience applications

Discussion: Neuroscience applications

Example: LFADS