

**2025 TAMPA BAY REASONABLE
ASSURANCE COMPLIANCE ASSESSMENT
REPORT**

Table of contents

1	The Tampa Bay Nitrogen Management Consortium Partnership for Progress	3
2	2025 Tampa Bay Estuary Nutrient Management Compliance Assessment	5
3	2025 Results Summary	10
	References	22

1 The Tampa Bay Nitrogen Management Consortium Partnership for Progress



TO: Adam Blalock, FDEP
Kevin J. McOmber, US EPA Region 4

FROM: Ed Sherwood, TBEP Executive Director (NMC Facilitator)

DATE: Jan. 28, 2026

SUBJECT: 2025 Tampa Bay Nutrient Management Compliance Assessment Results

cc Ken Weaver, Jessica Mostyn, Ben Ralys, Kevin O'Donnell, Lawrence Glenn, Kenneth Hayman (FDEP Tallahassee)

Ramandeep Kaur, Jorge Perez, Lance Kautz, Jessica Pein, Erica Peck (FDEP Tampa)

Craig Hesterlee, Jeaneanne M. Gettle, Wade Lehmann, Cindy Barger, Nancy Laurson, Felicia Burks, Johnnie Purify (EPA Region 4/HQ)

Michele Duggan, Santino Provenzano (TBNMC)

Ed Sherwood, Maya Burke, Marcus Beck (TBEP)

Source content: [here](#)

On behalf of the Tampa Bay Nitrogen Management Consortium, please find attached the 2025 update on water quality and seagrass resources in the Tampa Bay estuary. This update has been developed in accordance with the compliance assessment adopted through [FDEP's](#)

[Tampa Bay Reasonable Assurance determination on December 22, 2010](#), FDEP's subsequent approval of the [2022 RA Update](#), and the [federally-recognized TMDL for Tampa Bay](#). The formal annual compliance assessment utilized by the Consortium is detailed in Section VIII.B of the [Final 2009 Reasonable Assurance Addendum: Allocation and Assessment Report](#).

Chlorophyll-a concentrations for all four major bay segments were below FDEP-approved numeric nutrient criteria thresholds in 2025. In the Remainder Lower Tampa Bay segment, concentrations were below numeric nutrient criteria thresholds in Boca Ciega Bay South and Manatee River, while the chlorophyll-a threshold was exceeded in Terra Ceia Bay. The approved chlorophyll-a thresholds were adopted as part of FDEP's 2002 Reasonable Assurance determination for Tampa Bay, and, at that time, it was determined that Tampa Bay's seagrass restoration goals could be achieved if annual, uncorrected chlorophyll-a concentrations remained below these thresholds. If a bay segment's chlorophyll-a concentration remains above thresholds for 2 concurrent years, additional compliance assessment steps are required by the Consortium. This nutrient management strategy has been consistently used by the TBEP and Consortium in their Annual Decision Matrix and Assessment reports (Beck, Burke, and Sherwood 2025).

Seagrass coverage in Tampa Bay increased by 1,407 acres between 2022 and 2024. The Southwest Florida Water Management District's (SWFWMD) 2024 baywide seagrass coverage estimate is 31,563 acres, below the baywide goal of 40,000 acres (Figure 3.4). Gains were observed in all bay segments, except Old Tampa Bay where a loss was observed. For all Tampa Bay segments, water quality remained supportive of seagrass resources in 2025, though continued seagrass losses in the Old Tampa Bay segment continue to be examined. Annual [seagrass transect surveys](#) for Old Tampa Bay show a slight increase in seagrass in recent years, although definitive changes in coverage cannot be assessed until the next update from the SWFWMD using aerial images obtained over the winter 2025-26 period. Additional research and restoration initiatives, including a recently completed assimilative capacity assessment, are being conducted in response to this localized trend. The Consortium's approved nutrient management strategy remains a necessary tool to adaptively manage and address nutrient loading to the Tampa Bay estuary.

Thank you again for your continued participation in the Consortium's process. Please contact Ed Sherwood (esherwood@tbep.org) with any questions about the Consortium's Annual Compliance Assessment.

2 2025 Tampa Bay Estuary Nutrient Management Compliance Assessment

On December 22, 2010, then FDEP Secretary Drew signed a [Final Order](#) accepting and approving the [2009 Reasonable Assurance](#) Addendum for the Tampa Bay estuary. The final order found that the Nitrogen Management Consortium (NMC) provided FDEP reasonable assurance that: 1) completed and proposed management actions in the 2009 RA Addendum will result in the continued attainment of the estuarine nutrient criteria within Tampa Bay, and 2) compliance with the allocations in the 2009 RA Addendum ensures reasonable progress towards continued attainment of the estuarine nutrient criteria and associated Class III designated uses. Furthermore, the FDEP finalized a WQBEL for the Tampa Bay estuary in accordance with the allocations developed under the 2009 RA Addendum in November 2010. The Consortium completed subsequent RA Updates in [2012](#), [2017](#), and [2022](#) maintaining allocations and expanding upon projects originally defined in the [2002 RA Submittal](#), [2007 RA Update](#), [2009 RA Addendum](#), [2012 RA Update](#), [2017 RA Update](#) and [2022 RA Update](#).

As part of the compliance assessment stipulated under the 2009 RA Addendum, the NMC committed to annually assess the water quality and seagrass conditions within Tampa Bay and annually report these to FDEP and EPA. The Consortium's assessment responsibilities are shown in green in Figure [2.1](#). It should be noted that the Consortium's reasonable assurance assessment strategy begins with the observation of water quality conditions in the bay for a particular year. As is recommended in numerous EPA guidance documents for the development of numeric nutrient criteria, the Consortium's assessment strategy attempts to apply a stressor-response rationale for the determination of nitrogen load allocation reasonable assurance in the estuary.

The framework is applied on a bay-segment basis, and is predicated on assessing annual attainment of the bay segment chlorophyll-a concentration threshold as the initial step. If the bay segment-specific chlorophyll-a threshold is met, the Consortium annually reports the results to FDEP and EPA and additional assessment steps are not required by the Consortium (by June of the following year). If annual average chlorophyll-a thresholds are not met in one or more bay segments, additional assessment steps are required by the Consortium as noted in the framework and assessment process (Figure [2.1](#), Table [2.1](#)).

Regardless of the assessment results, the Consortium will annually report (by June of the following year) whether the bay segment specific chlorophyll-a thresholds are met using the

Environmental Protection Commission of Hillsborough County (EPCHC) dataset, as traditionally assessed using the “Decision Matrix” management strategy developed by the TBEP (Janicki, Wade, and Pribble 2000) and will deliver this to FDEP and EPA (Figure 2.1; NMC Action 1 in the Framework). Additional data from Pinellas and Manatee County are used to determine if chlorophyll thresholds in the Remainder Lower Tampa Bay segment (Boca Ciega Bay South, Terra Ceia Bay, and Manatee River) are met in these areas. If an annual, individual exceedence of a bay segment chlorophyll-a threshold is observed, an addendum report outlining the anomalous event(s) or data which influenced the bay segment chlorophyll-a exceedence will be delivered to FDEP and EPA upon review by NMC participants by September of the following year (Figure 2.1; NMC Action 2 in the Framework). An evaluation of the bay segment assimilative capacity (i.e. revision to the federally-recognized TMDL) is formally considered (if not already considered by the NMC) when bay segment chlorophyll-a thresholds are not met in 2 concurrent years, and hydrologically normalized loads for those years meet the federally-recognized TMDL (Figure 2.1; NMC Action 3 in the Framework). This was the case for the Old Tampa Bay segment during the 2019-2021 period and an assimilative capacity assessment was completed in 2025. Alternatively, when bay segment chlorophyll-a thresholds are not met in 2 concurrent years and hydrologically normalized loads for those years also do not meet the federally-recognized TMDL, the Consortium will deliver a full loading report to FDEP and EPA comparing the observed, combined entity/source annual or multiple year loadings to the sources’ 5-yr annual average allocation by September of the following year. This report will identify any exceedences among combined entity/source load categories after taking into consideration “set allocation” sources and hydrologically-normalized sources, and if necessary, whether exceedences were observed for individual MS4 or unpermitted (LA) sources (Figure 2.1; NMC Action 4 in the Framework). It is noted that FDEP will independently assess individual entities for compliance with their allocations.

Table 2.1: Assessment steps linked to the Nitrogen Management Consortium’s decision framework, as depicted in the flowchart above.

Assessment Step	Result	Action
I. Determine annual bay segment specific chlorophyll-a FDEP threshold attainment as traditionally assessed using the Decision Matrix management strategy developed by the TBEP (Janicki, Wade, and Pribble 2000).	Yes	NMC Action 1

	No	NMC Action 1
II. Review data and determine if an anomalous event(s) influenced non-attainment of the bay segment specific chlorophyll-a threshold.	Yes	NMC Action 2
	No	Go to III
III. Determine if the chlorophyll-a thresholds have been exceeded for <2 consecutive years.	Yes	NMC Action 2
	No	Go to IV
IV. Determine if the bay segment specific federally-recognized TMDL has been achieved using the hydrologically-adjusted compliance assessment outlined in NMC Decision Memo #11 (Appendix 2-11).	Yes	NMC Action 3
	No	Go to V
V. For a given year or for multiple years, compile and report entity-specific combined source loads in comparison to 5-yr annual average reasonable assurance allocation.	Compile & Report	NMC Action 4

NMC actions outlined in Figure 2.1 and Table 2.1 performed during RA Implementation Period (2022-2026) are as follows:

- NMC Action 1 - A report assessing attainment of bay segment specific chlorophyll-a thresholds using the EPCHC dataset, as traditionally assessed using the Decision Matrix management strategy developed by the TBEP (Janicki, Wade, and Pribble 2000) will be delivered to FDEP and EPA (this report).
- NMC Action 2 - A report of the anomalous event(s) or data which influenced the bay segment chlorophyll-a exceedence will be delivered to FDEP and EPA, upon review by NMC participants (this report).
- NMC Action 3 - Consider re-evaluation of the bay segment assimilative capacity based on nonattainment of bay segment chlorophyll-a threshold while meeting federally-recognized TMDL.
- NMC Action 4 - If federally-recognized TMDL not achieved, compile results of hydrologic evaluation for FDEP's review and identify potential further actions needed to achieve reasonable assurance for bay segment allocations.

3 2025 Results Summary

Results from 2025 indicate that all RA bay segments met chlorophyll-a thresholds accepted by the FDEP to maintain FDEP Reasonable Assurance for Tampa Bay and to comply with the EPA TMDL (Figure 3.1) and estuarine numeric nutrient criteria for Tampa Bay (EPA Amended Approval Letter Jun. 28, 2013). A water quality dashboard (<https://shiny.tbep.org/wq-dash>) is available to further synthesize available data, assess additional water quality metrics (phytoplankton counts), and inform Consortium participants and other resource managers on the status of water quality in Tampa Bay. Chlorophyll-a concentrations were notably lower than the previous year for many bay segments likely because of below average rainfall in 2025 (Figure 3.3). Blooms of harmful algal species (e.g., *Karenia brevis*, *Pyrodinium bahamense*) were not observed in any of the bay segments. Although understanding and mitigating blooms of *Pyrodinium bahamense* in OTB continues to be a focus of research efforts, summer concentrations in Old Tampa Bay were lower compared to prior years and the chlorophyll-a criteria was met for the fourth consecutive year of the RA period. Recommendations from the assimilative capacity assessment for Old Tampa Bay to address recurring water quality issues in that bay segment were presented to the Tampa Bay Nitrogen Management Consortium (TB-NMC) and Old Tampa Bay working group (OTBWG) in 2025 (Stantec Consulting Services, Inc. 2024, 2025a, 2025b).

Key findings from the Old Tampa Bay Assimilative Capacity Assessment project include: 1) Relationships among nitrogen, chlorophyll, water clarity, and seagrass within OTB vary over time and space within OTB sub-segments, and now differ from the broader-scale relationships documented in Tampa Bay as a whole; 2) Although linear pathways between nitrogen, chlorophyll, light attenuation, and seagrass remain valid within OTB, the existing paradigm is likely too simplistic to account for potentially synergistic or compounding stressors now being observed in the bay segment; 3) While varying hydrologic inputs and resulting residence times still appear influential to chlorophyll-a NNC attainment, the nitrogen delivery ratios for Old Tampa Bay are consistently discordant with chlorophyll-a NNC attainment; 4) Absolute TN loading assessments were more robustly linked to annual chlorophyll-a NNC attainment compared to normalized loads and alternative hydrologic normalization processes did not produce more robust links to chlorophyll-a attainment; 5) Assessing attainment of seasonal chlorophyll-a management targets for the NW and CW sub-segments of OTB may provide resource managers an earlier warning sign and more proactive opportunity to adaptively manage any subsequent chlorophyll-a NNC exceedance at the annual time-scale; 6) In addition to permitted load allocations, alternative management interventions such as oyster reef restoration or reductions in managed freshwater releases from Lake Tarpon may support better attainment

of water quality standards in OTB; and 7) CASM-OTB model results corroborated empirical analyses from Task 2 and other mechanistic modeling efforts (i.e., Integrated Ecosystem Model, WASP) that suggested more consistent attainment of the chlorophyll-a NNC relative to an absolute TN loading threshold of 486 tons/yr. These findings and continued reporting on completed, ongoing, and planned nitrogen load reduction projects will be incorporated into the next RA update.

Baywide coverage of seagrass is estimated to be 31,563 acres. The next SWFWMD seagrass coverage estimate will be developed from aerial photographs acquired over the winter 2025-2026 period. Between 2022 and 2024, gains were observed in all bay segments, excluding Old Tampa Bay. Annual transect surveys conducted by TBEP and its partners (shown in the [seagrass transect dashboard](#)) are used to corroborate areal trends. In 2025, slight gains in frequency occurrence in Old Tampa Bay suggest seagrasses may have increased in the last year. Additional research is being pursued by the [Southwest Florida Seagrass Working Group](#) to better understand the potential influence of long-term temperature increases and salinity reductions on seagrass declines (Beck et al. 2024).

Detailed results for the 2022-2026 RA implementation period are provided in Tables [3.2](#), [3.3](#), [3.4](#), [3.5](#), and [3.6](#) for each bay segment. Results for the Remainder Lower Tampa Bay segment (Boca Ciega Bay South, Terra Ceia Bay, Manatee River) are also included in this annual assessment (reporting began with the 2023 annual assessment). As of the 2025 reporting period, NMC Actions 2-5 are not necessary based upon observed water quality conditions within Tampa Bay, though work is being continued by the TBEP and TBNMC to further understand drivers of seagrass change and Old Tampa Bay's current assimilative capacity. Individual annual reports of the bay's conditions from 2025 can be found on the TBEP website, as specified in the following link (Beck, Burke, and Sherwood 2025) and the [water quality dashboard](#). A summary of historic attainment of the regulatory chlorophyll-a thresholds for each of the bay segments is depicted in Figure [3.5](#).

Lastly, annual hydrologic conditions among all the major bay segments in 2025 were below 1992-1994 levels for all segments except Old Tampa Bay (Table [3.1](#)). Therefore, hydrologic adjustments for evaluating compliance with individual entity load allocations/permitting targets should be applied for each major segment except Old Tampa Bay (Janicki Environmental, Inc. 2012, 2016). The estimated hydrologic loads for each bay segment relative to observed 1992-1994 levels are indicated in the table below. The estimated compliance load adjustment factors (if applicable) are also specified. A tool to calculate the hydrologic estimates and adjustment factors by bay segment is available online through an interactive dashboard and automatically updated as provisional hydrologic estimates are approved by monitoring agencies (https://shiny.tbep.org/tbnmc_hydrologic_estimates/).

Table 3.1: Hydrologic load estimates in 2025 relative to 1992-1994 levels and estimated compliance load adjustment factors for the major bay segments.

Bay Segment	1992 - 1994 Hydrology (95% Prediction Interval, million m3)	Hydrology Estimate (million m3)	Compliance Load Adjustment Factor
Old Tampa Bay	383 - 548	471.14	
Hillsborough Bay	753-1110	682.75	0.75
Middle Tampa Bay	524-756	488.38	0.76
Lower Tampa Bay	312-402	296.93	0.82

Table 3.2: Demonstration of reasonable assurance assessment steps for Old Tampa Bay. Green and red squares indicate outcomes of decision points outlined in the Consortium's reasonable assurance assessment framework.

Bay Segment Reasonable Assurance Assessment Steps	DATA USED TO ASSESS ANNUAL REASONABLE ASSURANCE					OUTCOME
	Year 1 (2022)	Year 2 (2023)	Year 3 (2024)	Year 4 (2025)	Year 5 (2026)	
NMC Action 1: Determine if observed chlorophyll-a exceeds FDEP threshold of 9.3 ug/L	No (7.1)	No (6.2)	No (8.8)	No (6.6)		All years below threshold so far, not necessary for NMC Actions 2-5
NMC Action 2: Determine if any observed chlorophyll-a exceedences occurred for 2 consecutive years	No	No	No	No		All years met threshold, not necessary for NMC Actions 3-5

NMC Action 3: Determine if observed hydrologically-normalized total load exceeds federally-recognized TMDL of 486 tons/year	N/A	N/A	N/A	N/A		Not necessary due to observed water quality and seagrass conditions in the bay segment
NMC Actions 4-5: Determine if any entity/source/facility specific exceedences of 5-yr average allocation occurred during implementation period						Not necessary when chlorophyll-a threshold met

Table 3.3: Demonstration of reasonable assurance assessment steps for Hillsborough Bay. Green and red squares indicate outcomes of decision points outlined in the Consortium's reasonable assurance assessment framework.

Bay Segment Reasonable Assurance Assessment Steps	DATA USED TO ASSESS ANNUAL REASONABLE ASSURANCE					OUTCOME
	Year 1 (2022)	Year 2 (2023)	Year 3 (2024)	Year 4 (2025)	Year 5 (2026)	
NMC Action 1: Determine if observed chlorophyll-a exceeds FDEP threshold of 15 ug/L	No (8.9)	No (6.9)	No (10.7)	No (10)		All years below threshold so far, not necessary for NMC Actions 2-5
NMC Action 2: Determine if any observed chlorophyll-a exceedences occurred for 2 consecutive years	No	No	No	No		All years met threshold, not necessary for NMC Actions 3-5
NMC Action 3: Determine if observed hydrologically-normalized total load exceeds federally-recognized TMDL of 1451 tons/year	N/A	N/A	N/A	N/A		Not necessary due to observed water quality and seagrass conditions in the bay segment

NMC Actions 4-5: Determine if any entity/source/facility specific exceedences of 5-yr average allocation occurred during implementation period	Not necessary when chlorophyll-a threshold met
---	--

Table 3.4: Demonstration of reasonable assurance assessment steps for Middle Tampa Bay. Green and red squares indicate outcomes of decision points outlined in the Consortium's reasonable assurance assessment framework.

Bay Segment Reasonable Assurance Assessment Steps	DATA USED TO ASSESS ANNUAL REASONABLE ASSURANCE					OUTCOME
	Year 1 (2022)	Year 2 (2023)	Year 3 (2024)	Year 4 (2025)	Year 5 (2026)	
NMC Action 1: Determine if observed chlorophyll-a exceeds FDEP threshold of 8.5 ug/L	No (5)	No (3.7)	No (7.8)	No (5.5)		All years below threshold so far, not necessary for NMC Actions 2-5
NMC Action 2: Determine if any observed chlorophyll-a exceedences occurred for 2 consecutive years	No	No	No	No		All years met threshold, not necessary for NMC Actions 3-5
NMC Action 3: Determine if observed hydrologically-normalized total load exceeds federally-recognized TMDL of 799 tons/year	N/A	N/A	N/A	N/A		Not necessary due to observed water quality and seagrass conditions in the bay segment
NMC Actions 4-5: Determine if any entity/source/facility specific exceedences of 5-yr average allocation occurred during implementation period						Not necessary when chlorophyll-a threshold met




Figure 3.1: Historic chlorophyll-a annual averages for the four major bay segments of Tampa Bay and those that include the Remainder Lower Tampa Bay segment (Boca Ciega Bay South, Terra Ceia Bay, Manatee River). Annual averages in 2025 were below the regulatory thresholds developed under the Tampa Nitrogen Management Consortium’s nutrient management strategy in all bay segments. Vertical grey bars indicate the portion of the 2022-2026 Reasonable Assurance compliance assessment period covered by the results.




Figure 3.2: Map depicting individual station chlorophyll-a exceedences in Tampa Bay relative to FDEP regulatory thresholds for chlorophyll-a in 2025. Note individual station exceedences do not indicate failed compliance at the bay segment scale.




Figure 3.3: 2025 monthly chlorophyll-a bay segment means (red dots) compared to monthly distributions from prior years (box plots and black dots). Prior years extend to 1975 for Old Tampa Bay, Hillsborough Bay, Middle Tampa Bay, and Lower Tampa Bay, 1991 for Boca Ciega Bay South, 1989 for Terra Ceia Bay, and 1990 for Manatee River. Note that Pinellas and Manatee County data are used for Boca Ciega Bay South, Terra Ceia Bay, and Manatee River and has less frequent sampling intervals than data from the Environmental Protection Commission of Hillsborough County used for the other bay segments. Boxes encompass the 25th and 75th percentiles, while whiskers bound the interquartile range. Dots beyond the whiskers represent outliers throughout the 1975-2024 sample period.




Figure 3.4: Historic seagrass coverage estimates for Tampa Bay. The target coverage of 38,000 acres was changed to 40,000 acres in 2020 to reflect programmatic goals in the 2020 Habitat Master Plan Update ([TBEP #07-20](#)). Data source: TBEP & SWFWMD.

Table 3.5: Demonstration of reasonable assurance assessment steps for Lower Tampa Bay. Green and red squares indicate outcomes of decision points outlined in the Consortium’s reasonable assurance assessment framework.

Bay Segment Reasonable Assurance Assessment Steps	DATA USED TO ASSESS ANNUAL REASONABLE ASSURANCE					OUTCOME
	Year 1 (2022)	Year 2 (2023)	Year 3 (2024)	Year 4 (2025)	Year 5 (2026)	
NMC Action 1: Determine if observed chlorophyll-a exceeds FDEP threshold of 5.1 ug/L	No (3.6)	No (2.6)	No (4.6)	No (3)		All years below threshold so far, not necessary for NMC Actions 2-5
NMC Action 2: Determine if any observed chlorophyll-a exceedences occurred for 2 consecutive years	No	No	No	No		All years met threshold, not necessary for NMC Actions 3-5

NMC Action 3: Determine if observed hydrologically-normalized total load exceeds federally-recognized TMDL of 349 tons/year	N/A	N/A	N/A	N/A		Not necessary due to observed water quality and seagrass conditions in the bay segment
NMC Actions 4-5: Determine if any entity/source/facility specific exceedences of 5-yr average allocation occurred during implementation period						Not necessary when chlorophyll-a threshold met

Table 3.6: Demonstration of reasonable assurance assessment steps for Remainder Lower Tampa Bay. Green and red squares indicate outcomes of decision points outlined in the Consortium's reasonable assurance assessment framework.

Bay Segment Reasonable Assurance Assessment Steps	DATA USED TO ASSESS ANNUAL REASONABLE ASSURANCE					OUTCOME
	Year 1 (2022)	Year 2 (2023)	Year 3 (2024)	Year 4 (2025)	Year 5 (2026)	
NMC Action 1: Determine if observed chlorophyll-a exceeds FDEP threshold of 5.1 ug/L	No (3.6)	No (2.6)	No (4.6)	No (3)		All years below threshold so far, not necessary for NMC Actions 2-5
NMC Action 2: Determine if any observed chlorophyll-a exceedences occurred for 2 consecutive years	No	No	No	No		All years met threshold, not necessary for NMC Actions 3-5
NMC Action 3: Determine if observed hydrologically-normalized total load exceeds federally-recognized TMDL of 349 tons/year	N/A	N/A	N/A	N/A		Not necessary due to observed water quality and seagrass conditions in the bay segment

NMC Actions 4-5: Determine if any entity/source/facility specific exceedences of 5-yr average allocation occurred during implementation period	Not necessary when chlorophyll-a threshold met
---	--

	OTB	HB	MTB	LTB	RALTB
1975	R	R	R	G	
1976	R	R	R	G	
1977	R	R	R	R	
1978	R	R	R	G	
1979	R	R	R	R	
1980	R	R	R	R	
1981	R	R	R	R	
1982	R	R	R	R	
1983	R	R	R	R	
1984	G	G	R	G	
1985	R	R	R	G	
1986	R	R	G	G	
1987	R	G	R	G	
1988	G	G	G	G	
1989	R	G	G	G	
1990	R	G	G	G	
1991	G	G	G	G	G
1992	G	G	G	G	G
1993	G	G	G	G	G
1994	R	R	R	R	R
1995	R	R	R	G	G
1996	G	G	G	G	G
1997	G	G	G	G	G
1998	R	R	R	R	R
1999	G	G	G	G	G
2000	G	G	G	G	G
2001	G	G	G	G	G
2002	G	G	G	G	G
2003	R	G	G	G	G
2004	R	G	G	G	G
2005	G	G	G	R	R
2006	G	G	G	G	G
2007	G	G	G	G	G
2008	G	G	G	G	G
2009	R	G	G	G	G
2010	G	G	G	G	G
2011	R	G	G	G	G
2012	G	G	G	G	G
2013	G	G	G	G	G
2014	G	G	G	G	G
2015	R	G	G	G	G
2016	G	G	G	G	G
2017	R	G	G	G	G
2018	G	G	G	G	G
2019	R	G	G	G	G
2020	R	G	G	G	G
2021	R	G	G	G	G
2022	G	G	G	G	G
2023	G	G	G	G	G
2024	G	G	G	G	G
2025	G	G	G	G	G

Figure 3.5: Attainment of adopted chlorophyll-a thresholds (1975 - 2025) in the four major bay segments and Remainder Lower Tampa Bay segment (Boca Ciega Bay South, Terra Ceia Bay, Manatee River). Green (yes) indicates that average annual chlorophyll-a thresholds were met; red (no) indicates that threshold levels were not met. Grey line is the beginning of the current Reasonable Assurance implementation period.

References

- Beck, M. W., M. C. Burke, and E. T. Sherwood. 2025. “2024 Tampa Bay Water Quality Assessment.” 03-25. St. Petersburg, Florida: Tampa Bay Estuary Program. <https://drive.google.com/file/d/1jZwjt6c0LdiGL3Arl6uFVIn25Iji4ITx/view?usp=sharing>.
- Beck, M. W., K. E. Flaherty-Walia, S. Scolaro, M. C. Burke, B. T. Furman, D. J. Karlen, C. Pratt, C. J. Anastasiou, and E. T. Sherwood. 2024. “Hot and Fresh: Evidence of Climate-Related Suboptimal Water Conditions for Seagrass in a Large Gulf Coast Estuary.” *Estuaries & Coasts* 47: 1475–97. <https://doi.org/https://doi.org/10.1007/s12237-024-01385-0>.
- Janicki, A., D. Wade, and J. R. Pribble. 2000. “Developing and Establishing a Process to Track the Status of Chlorophyll-a Concentrations and Light Attenuation to Support Seagrass Restoration Goals in Tampa Bay.” 04-00. St. Petersburg, Florida: Tampa Bay Estuary Program. https://drive.google.com/file/d/1XMULU8w4syWcSv_ciOUOhnC_G4xt6GIF/view?usp=drivesdk.
- Janicki Environmental, Inc. 2012. “Development of a Screening Level Tool for Estimating Annual Hydrologic Loadings to Tampa Bay.” 05-12. St. Petersburg, Florida: Tampa Bay Estuary Program. <https://drive.google.com/file/d/1C6Arwat9IxYs8jeTZpcmBB6WBTpeQ2nw/view?usp=drivesdk>.
- . 2016. “Update on the Development of a Screening Level Tool for Estimating Annual Hydrologic Loadings to Tampa Bay.” 03-16. St. Petersburg, Florida: Tampa Bay Estuary Program. <https://drive.google.com/file/d/11NT0NQ2WbPO6pVZaD7P7Z6qjcwO1jxHw/view?usp=drivesdk>.
- Stantec Consulting Services, Inc. 2024. “Old Tampa Bay (OTB) Assimilative Capacity Assessment Project: Task 1 Evaluating Existing Management Paradigm and Assessing Alternatives.” 09-24. St. Petersburg, Florida: Tampa Bay Estuary Program. https://drive.google.com/file/d/1IfvUTLE20wzVgzULuyeRGWapEFT3DhQ0/view?usp=drive_link.
- . 2025a. “Old Tampa Bay Assimilative Capacity Study: Task 2 Evaluation of Eutrophication Indicator Targets, Thresholds, and Numeric Nutrient Criteria and Proposed Adjustments.” 05-25. St. Petersburg, Florida: Tampa Bay Estuary Program. <https://drive.google.com/file/d/12xYfSs-4bxhLa1ZVQbmhQSv5Jo4Jf-oY/view?usp=sharing>.
- . 2025b. “Old Tampa Bay Assimilative Capacity Study: Task 3 Management Intervention Assessment and Recommendations.” 17-25. St. Petersburg, Florida: Tampa Bay Estuary Program. <https://drive.google.com/file/d/1uVeMIpSylz1Mull6WMwVDc7KVInzosqy/view?usp=sharing>.